Motion in a Pl ane



1. State, for each of the following physical quantities, if it is a scalar or a vector. Volume, mass, speed, acceleration, density, number of moles, velocity, angular frequency, displacement, angular velocity.

Ans. Scalar. Mass, volume, density, angular frequency, number of moles, speed.

Vector. Acceleration, angular velocity, velocity, displacement.

A scalar quantity is specified by its magnitude. Mass, volume, density, angular frequency, number of moles, speed are some of the scalar physical quantities.

A vector quantity is specified by its magnitude and the direction associated with it.

Acceleration, angular velocity, velocity, displacement belong to this category.

2. Pick out the two scalar quantities in the following list.

Force, angular momentum, work, current, linear momentum, electric field, average velocity, magnetic moment, relative velocity.

Ans. Work and current are examples of scalar quantities.

Work done is said to be the dot product of force and displacement. As the dot product of two quantities is always a scalar, work is considered as a scalar physical quantity.

Current is described by its magnitude. Its direction is not considered.

Thus, it is a scalar quantity.

3. Pick out the only vector quantity in the following list. Temperature, pressure, impulse, time, power, total path length, energy, gravitational potential, coefficient of friction, charge.

Ans: Impulse

It is given by the product of force and time. As force is a vector quantity, its product with time gives a vector quantity.

4. State with reasons, whether the following algebraic operations with scalar and vector physical quantities are meaningful.

(a) Adding any two scalars.

Ans: Not meaningful.

The addition of two scalar quantities will be meaningful only if they both represent the same physical quantity.

(b) Adding a scalar to a vector of the same dimensions.

Ans: Not meaningful.

The addition of a vector quantity with a scalar quantity is considered not meaningful.

(c) Multiplying any vector by any scalar.

Ans: Meaningful.

A scalar can be multiplied with a vector. Force is multiplied with time to give impulse.

(d) Multiplying any two scalars.

Ans: Meaningful.

A scalar, respective of the physical quantity, can be multiplied with another scalar having the same or different dimensions.

(e) Adding any two vectors.

Ans: Not meaningful.

The addition of two vector quantities is considered meaningful only if they both represent the same physical quantity.

(f) Adding a component of a vector to the same vector.

Ans: Meaningful

A component of a vector can be added to the same vector as both of them have the same dimensions.

5. Read each statement below carefully and state with reasons, if it is true or false.

(a) The magnitude of a vector is always a scalar.

Ans: True.

The magnitude of a vector is a number. So, it is a scalar.

(b) Each component of a vector is always a scalar.

Ans: False.

Each component of a vector is a vector.

(c) The total path length is always equal to the magnitude of the displacement vector of a particle.

Ans: False.

The total path length is scalar, whereas displacement is a vector quantity.

So, the total path length is greater than the magnitude of displacement. It is equal to the magnitude of displacement only when a particle is moving in a straight line.

(d) The average speed of a particle (defined as total path length divided by the time taken to cover the path) is either greater or equal to the magnitude of average velocity of the particle over the same interval of time.

Ans: True.

It is because the total path length is always greater than or equal to the magnitude of displacement of a particle.

(e) Three vectors not lying m a plane can never add up to give a null vector.

Ans: True.

Three vectors, which do not lie in a plane, can't be represented by the sides of a triangle taken in the same order.

6. Establish the following vector inequalities geometrically or otherwise.

- (a) $|a+b| \le |a|+|b|$
- **(b)** $|a+b| \ge |a|-|b|$
- (c) $|a-b| \le |a|-|b|$
- (d) $|a-b| \ge |a|-|b|$

When does the equality sign above apply?

Ans: Let a and b be represented by the adjacent sides of a parallelogram *OMNP* as shown below.



 $|\overrightarrow{OM}| = |\overrightarrow{a}|$...(i)

 $|\overrightarrow{\mathrm{MN}}| = |\overrightarrow{\mathrm{OP}}| = |\overrightarrow{\mathrm{b}}| \qquad ...(\mathrm{ii})$

 $|\overrightarrow{ON}| = |\overrightarrow{a} + \overrightarrow{b}|$...(iii)

As each side is smaller than the sum of the other two sides in a triangle,

In ΔOMN ,

ON < (OM + MN)

$$|\vec{a} + \vec{b}| < |\vec{a}| + |\vec{b}|$$
 ...(iv)

If \vec{a} and \vec{b} act along a straight line in the same direction, then.

$$|\vec{a} + \vec{b}| = |\vec{a}| + |\vec{b}|$$
(v)

Combine equations (iv) and (v)

$$|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}|$$

Let \vec{a} and \vec{b} be represented by the adjacent sides of a parallelogram \vec{b} , as shown below.



$$|\overrightarrow{OM}| = |\vec{a}| \qquad \dots(i)$$
$$|\overrightarrow{MN}| = |\overrightarrow{OP}| = |\vec{b}| \qquad \dots(ii)$$
$$|\overrightarrow{ON}| = |\vec{a} + \vec{b}| \qquad \dots(iii)$$

As each side is smaller than the sum of the other two sides in a triangle,

In ΔOMN ,

ON + MN > OM

ON + OM > MN

 $|ON| > |\overrightarrow{OM} - \overrightarrow{OM}|$ (: OP = MN)

$$|a+b| > ||a| - |b|| \dots (iv)$$

If \vec{a} and \vec{b} act along a straight line in the same direction, then.

 $|\vec{a} + \vec{b}| = ||\vec{a}| - |\vec{b}||$ (v)

Combine equations (iv) and (v)

$$\mid \stackrel{\rightarrow}{a} + \stackrel{\rightarrow}{b} \mid \geq \mid \mid \stackrel{\rightarrow}{a} \mid - \mid \stackrel{\rightarrow}{b} \mid \mid$$

Let \vec{a} and \vec{b} be represented by the adjacent sides of a parallelogram *PQRS*

$$|\overrightarrow{OR}| = |\overrightarrow{PS}| = |\overrightarrow{b}|$$
 ...(i)
 $|\overrightarrow{OP}| = |\overrightarrow{a}|$...(ii)

As each side is smaller than the sum of the other two sides in a triangle,

In $\triangle OPS$,

OS < OP + PS

$$|\vec{a} - \vec{b}| < |\vec{a}| + |-\vec{b}|$$

$$|\vec{a} - \vec{b}| < |\vec{a}| + |\vec{b}|$$
 ...(iii)

If the two vectors act in a straight line but in opposite directions, then.

$$|\vec{a} - \vec{b}| = |\vec{a}| + |\vec{b}|$$

Combine equations (iii) and (iv)

$$|\vec{\mathbf{a}} - \vec{\mathbf{b}}| \leq |\vec{\mathbf{a}}| + |\vec{\mathbf{b}}|$$

Let \vec{a} and \vec{b} be represented by the adjacent sides of a parallelogram *PQRS*

$$OS + PS > OP$$
 ...(i)
 $OS > OP-PS$...(ii)

$$|\vec{a} - \vec{b}| > |\vec{a}| - |\vec{b}|$$
 ...(iii)

The L.H.S is always positive and R.H.S can be positive or negative.

To make both quantities positive, take modulus on both sides.

$$\left\| \vec{a} - \vec{b} \right\| < \left\| \vec{a} \right\| - \left\| \vec{b} \right\|$$
$$\left\| \vec{a} - \vec{b} \right\| > \left\| \vec{a} \right\| - \left\| \vec{b} \right\| \qquad \dots (iv)$$

If the two vectors act in a straight line but in the same direction.

$$\left|\vec{a} - \vec{b}\right| = \left\|\vec{a}\right| - \left|\vec{b}\right|$$
 ...(v)

Combine equation ^(iv) and equation^(v).

$$\left|\vec{a} - \vec{b}\right| \ge \left\|\vec{a}\right| - \left|\vec{b}\right\|$$

7. Given a+b+c+d=0, which of the following statements are correct.

(a) ^{a,b,c}, and ^d must each be a null vector.

Ans: Incorrect

To make $\vec{a} + \vec{b} + \vec{c} + \vec{d} = 0$, it is not necessary to have all four vectors as null vectors. There are many other combinations which will give the sum zero.

(b) The magnitude of (a+c) equals the magnitude of (b+d).

Ans: Correct

 $\vec{a} + \vec{b} + \vec{c} + \vec{d} = 0$

$$\vec{a} + \vec{c} = -(\vec{b} + \vec{d})$$

Take modulus on both sides.

$$|\vec{a} + \vec{c}| = |-(\vec{b} + \vec{d})| = |(\vec{b} + \vec{d})|$$

So, the magnitude of $(\vec{a} + \vec{c})$ is the same as the magnitude of $(\vec{b} + \vec{d})$

(c) The magnitude of a can never be greater than the sum of the magnitudes of b, c, and d.

Ans: Correct

 $\vec{a} + \vec{b} + \vec{c} + \vec{d} = 0$ $\vec{a} = -(\vec{b} + \vec{c} + \vec{d})$

Take modulus on both sides.

$$\vec{a} = |(\vec{b} + \vec{c} + \vec{d})|$$

 $\vec{a} \le |\vec{b}| + |\vec{c}| + |\vec{d}|$ (i)

 $(\vec{b} + \vec{c} + \vec{d})$ is the sum of vectors \vec{b} , \vec{c} and \vec{d} . The magnitude of $(\vec{b} + \vec{c} + \vec{d})$ is less than, or equal to the sum of the magnitudes of \vec{b} , \vec{c} and \vec{d} . So, the magnitude of \vec{a} cannot be greater than the sum of the magnitudes of \vec{b} , \vec{c} and \vec{d} . Equation (i) shows that the magnitude of \vec{a} is equal to or less than the sum of the magnitudes of \vec{b} , \vec{c} and \vec{d}

(d) b+c must lie in the plane of a and d if a and d are not collinear, and in the line of a and d. If they are collinear?

Ans: Correct

For,
$$\vec{a} + \vec{b} + \vec{c} + \vec{d} = 0$$

 $\vec{a} + (\vec{b} + \vec{c}) + \vec{d} = 0$

The resultant sum of the vectors \vec{a} , $(\vec{b} + \vec{c})$ and \vec{d} is zero only if $(\vec{b} + \vec{c})$ lie in the same plane as \vec{a} and \vec{d}

If \vec{a} and \vec{d} are collinear, then $(\vec{b} + \vec{c})$ is in the line of \vec{a} and \vec{d} . This is true in this case and the vector sum of all the vectors will be zero.

8. Three girls skating on a circular ice ground of radius ^{200m} start from a point ^P on the edge of the ground and reach a point ^Q diametrically opposite to ^P following different paths as shown in Fig. 4.20. What is the magnitude of

the displacement vector for each? For which girl is this equal to the actual length of the path skated?



Ans: The magnitudes of displacements are equal to the diameter of the ground.

Radius of the ground = 200 m

Diameter of the ground $= 2 \times 200 = 400 \text{ m}$

So, the magnitude of the displacement for each girl is 400 m which is equal to the actual length of the path skated by girl B.

9. A cyclist starts from the centre O of a circular park of radius 1 km, reaches the edge P of the park, then cycles along the circumference, and returns to the centre along QO as shown in the following figure. If the round trip takes min, what is the

(a) Net displacement.

- **Ans:** The cyclist comes to the starting point after cycling for ¹⁰ minutes. So, his net displacement is zero.
 - (b) Average velocity

Ans: Average velocity

= Net displacement Total time Total time

As the net displacement of the cyclist is zero, his average velocity is also zero.

(c) Average speed of the cyclist?

Ans:

Average speed = $\frac{\text{Total path length}}{\text{Total time}}$

Total path length = OP + PQ + QO

$$= 1 + \frac{1}{4}(2\pi \times 1) + 1$$
$$= 2 + \frac{1}{2}\pi 3.570 \,\mathrm{km}$$

Time taken =10min

$$= \frac{10}{60}$$
$$= \frac{1}{6}h$$
$$\therefore \text{ Average speed}$$

$$=\frac{3.570}{1}=21.42\,\mathrm{km/h}$$

10. On an open ground, a motorist follows a track that turns to his left by an angle of 60° after every 500m. Starting from a given turn, specify the displacement of the total at the third, sixth and eighth turn. Compare the magnitude of the displacement with the total path length covered by the motorist in each case.

Ans: The path is a regular hexagon with side 500m.

Let the motorist start from P.

The motorist takes the third turn at S.

 \therefore Magnitude of displacement = PS

= PV + VS

=500+500

 $=1000 \,\mathrm{m}$

Total path length = PQ + QR + RS

=500+500+500

 $=1500 \,\mathrm{m}$

The motorist takes the sixth turn at P, which is the starting point.

 \therefore Magnitude of displacement =0

Total path length = PQ + QR + RS + ST + TU + UP

=500+500+500+500+500+500

 $= 3000 \,\mathrm{m}$

The motorist takes the eight turn at R.

 \therefore Magnitude of displacement = PR

$$=\sqrt{PQ^2+QR^2+2(PQ)\cdot(QR)\cos 60^\circ}$$

$$=\sqrt{500^{2}+500^{2}+(2\times500\times500\times\cos{60^{\circ}})}$$

$$=\sqrt{250000+250000+\left(500000\times\frac{1}{2}\right)}$$

 $= 866.03 \,\mathrm{m}$

$$\beta = \tan^{-1} \left(\frac{500 \sin 60^{\circ}}{500 + 500 \cos 60^{\circ}} \right)$$

 $\beta = 30^{\circ}$

Thus, the magnitude of displacement is 866.03 m at an angle of 30° with PR.

Total path length = Circumference of the hexagon +PQ + QR

 $= 6 \times 500 + 500 + 500$

 $=4000\,\mathrm{m}$

Turn	Magnitude of displacement	Total path length
Third	1000	1500
Sixth	0	3000
Eighth	866.03; 30°	4000

11. A passenger arriving in a new town wishes to go from the station to a hotel located 10 km away on a straight road from the station. A dishonest cabman takes him along a circuitous path 23 km long and reaches the hotel in 28 min. What is

(a) the average speed of the taxi.

Ans: Total distance travelled = 23 km

Total time taken,

 $= 28 \min = \frac{28}{60} h$

Average speed of the taxi,

= Total distance travelled Total time taken

(b) the magnitude of average velocity? Are the two equal?

Ans: Distance between the hotel and the station =10km = Displacement of the car

$$\therefore \text{ Average velocity } = \frac{10}{\frac{28}{60}}$$

 $= 21.43 \, \text{km} / \text{h}$

12. The ceiling of a long hall is ²⁵m high. What is the maximum horizontal distance that a ball thrown with a spectrum of ⁻¹ can go without hitting the ceiling of the hall?

Ans: Speed of the ball, 40 ms^{-1}

Maximum height, h = 25 m

In projectile motion, the maximum height reached, by a body projected at an angle θ is.

$$h = \frac{u^2 \sin^2 \theta}{2g}$$
$$25 = \frac{(40)^2 \sin^2 \theta}{2 \times 9.8}$$
$$\sin^2 \theta = 0.30625$$
$$\sin \theta = 0.5534$$
$$\theta = \sin^{-1}(0.5534)$$

θ=33.60°

The horizontal range is $R = \frac{u^{2} \sin 2\theta}{g}$ $R = \frac{(40)^{2} \times \sin 2 \times 33.60}{9.8}$ $R = \frac{1600 \times \sin 67.2}{9.8}$ $R = \frac{1600 \times 0.922}{9.8}$ R = 150.53m

13. A cricketer can throw a ball to a maximum horizontal distance of 100m .How much high above the ground can the cricketer throw the same ball?

Ans: Maximum horizontal distance, R=100m

The cricketer will throw the ball to the maximum horizontal distance when the angle of projection is 45° , i.e., $\theta = 45^{\circ}$

The horizontal range for a projection velocity v, is.

$$R = \frac{u^2 \sin 2\theta}{g}$$
$$100 = \frac{u^2 \sin 90^\circ}{g}$$
$$\frac{u^2}{g} = 100 \qquad \dots(i)$$

The ball will reach the maximum height when it is thrown vertically upward. For this type of motion, the final velocity is zero at the maximum height H.

Acceleration, a = -g

Use the third equation of motion.

$$v^{2} - u^{2} = -2gH$$
$$H = \frac{1}{2} \times \frac{u^{2}}{g}$$
$$H = \frac{1}{2} \times 100$$
$$H = 50 \,\mathrm{m}$$

14. A stone tied to the end of a string ⁸⁰cm long is whirled in a horizontal circle with a constant speed. If the stone makes ¹⁴ revolutions in ²⁵s, what is the magnitude and direction of acceleration of the stone?

Ans: Length of the string, = 80 cm = 0.8 m

Number of revolutions =14

Time taken = 25s

Frequency,

v =
$$\frac{\text{Number of revolutions}}{\text{Time taken}} = \frac{14}{25} \text{Hz}$$

Angular frequency, $\omega = 2\pi v$

$$= 2 \times \frac{22}{7} \times \frac{14}{25}$$
$$= \frac{88}{7} \operatorname{rads}^{-1}$$

$$=\frac{66}{25}$$
 rads⁻¹

Centripetal acceleration, $a_e = \omega^2 r$

$$= \left(\frac{88}{25}\right)^2 \times 0.8$$
$$= 9.91 \,\mathrm{ms}^{-2}$$

The direction of centripetal acceleration is always along the string, towards the center, at all points.

15. An aircraft executes a horizontal loop of radius 1.00 km with a steady speed of 900 km/h. Compare its centripetal acceleration with the acceleration due to gravity.

Ans: Radius of the loop, r = 1 km = 1000 m

Speed of the aircraft, $v = 900 \text{ kmh}^{-1}$

$$=900\times\frac{5}{18}$$

 $= 250 \, ms^{-1}$

Centripetal acceleration,

$$a_e = \frac{v^2}{r}$$

= $\frac{(250)^2}{1000}$
= 62.5 ms⁻²

Acceleration due to gravity, $g = 9.8 \text{ ms}^{-2}$

 $\frac{a_c}{g} = \frac{6.25}{1000}$ $a_c = 6.38g$

The Centripetal acceleration is 6.38 times the acceleration due to gravity.

16. Read each statement below carefully and state, with reasons, if it is true or false.

(a) The net acceleration of a particle m circular motion is always along the radius of the circle towards the centre.

Ans: False

In circular motion, the net acceleration of a particle is not always directed along the radius of the circle toward the centre. It happens only in the case of uniform circular motion.

(b) The velocity vector of a particle at a point is always along the tangent to the path of the particle at that point.

Ans: True

At a point on a circular path, a particle appears to move tangentially to the circular path.

Thus, the velocity vector of the particle is always along the tangent at a point.

(c) The acceleration vector of a particle in uniform circular motion averaged over one cycle is a null vector.

Ans: True

In uniform circular motion, the acceleration vector points towards the centre of the circle. The average of these vectors over one cycle is a null vector.

17. The position of a particle is given by $r = 3.0t\hat{i} - 2.0t\hat{j} + 4.0$ km

Where t is in seconds and the coefficients have the proper units for r to be in metres.

(a) Find the v and a of the particle?

Ans: $v(t) = (3.0\hat{i} - 4.0\hat{j}) : a = -4.0\hat{j}$

The position of the particle is.

 $\vec{r} = 3.0\hat{i} - 2.0t^2\hat{j} + 4.0k$

Velocity, \vec{v} of the particle is.

$$\vec{v} = \frac{d\vec{r}}{dt}$$
$$\vec{v} = \frac{d}{dt} (3.0t\hat{i} - 2 \cdot Ot^2\hat{j} + 4.0k)$$
$$\therefore \vec{v} = 3.0\hat{i} - 4.0t$$

Acceleration of the particle is.

$$\vec{a} = \frac{d\vec{v}}{dt}$$
$$\vec{a} = \frac{d}{dt}(3.0\hat{i} - 4.0\hat{j})$$
$$\therefore \vec{a} = -4.0\hat{j}$$

The velocity vector, $\vec{v} = 3.0\hat{i} - 4.0t\hat{j}$

At
$$t = 2.0s$$
.
 $\vec{v} = 3.0\hat{i} - 8.0\hat{j}$

(b) What is the magnitude and direction of velocity of the particle at t = 2.0s?

Ans: The magnitude of velocity is.

$$|\vec{v}| = \sqrt{3^2 + (-8)^2}$$
$$|\vec{v}| = \sqrt{73}$$
$$|\vec{v}| = 8.54 \text{ m/s}$$
$$And \qquad \theta = \tan^{-1}\left(\frac{v_y}{v_x}\right)$$
$$= \tan^{-1}\left(\frac{-8}{3}\right)$$
$$= -\tan^{-1}(2.667)$$

=-69.45°

The negative sign indicates that the direction of velocity is 8.54 ms^{-1} , 69.45° below the *x*-axis.

18. A particle starts from the origin at t = 0s with a velocity of 10.07 and moves in the x-y plane with a constant acceleration of $(8.0\hat{i}+2.0\hat{j})ms^{-2}$

(a) At what time is the x- coordinate of the particle 16 m? What is the y- coordinate of the particle at that time?

Ans: Velocity of the particle, $\vec{v} = 10.0 \hat{j} \text{ ms}^{-1}$

Acceleration of the particle $\vec{a} = (8.0\hat{i} + 2.0\hat{j})$

But, $\vec{a} = \frac{d\vec{v}}{dt} = 8.0\hat{i} + 2.0\hat{j}$

 $\vec{dv} = (8.0\hat{i} + 2.0\hat{j})dt$

Integrate both sides.

$$\vec{v}(t) = 8.0\hat{i} + 2.0\hat{j} + \vec{u}$$

Where,

 \vec{u} = Velocity vector of the particle at t=0

 \vec{v} = Velocity vector of the particle at time ϕ

$$\vec{v} = \frac{dr}{dt}$$

 $d\vec{r} = \vec{v}dt = (8.0t\hat{i} + 2.0t\hat{j} + u)dt$

Integrate the equations with. at t=0; r=0 and at t=t; r=r

$$\vec{r} = \vec{u}t + \frac{1}{2}8.0t^{2}\hat{i} + \frac{1}{2}2.0t^{2}\hat{j}$$
$$= \vec{u}t + 4 \times 0t^{2}\hat{i} + t^{2}\hat{j}$$
$$= (10.0j)t + 4.0t^{2}\hat{i} + i^{2}\hat{j}$$
$$\hat{i} + y = 4.0t^{2}\hat{i} + (10t + t^{2})\hat{j}$$

Equate the coefficients of \hat{i} and $\hat{j}.$

$$x = 4t^{2}$$
$$t = \left(\frac{x}{4}\right)^{\frac{1}{2}}$$

And $y=10t+t^2$

When x = 16m.

$$t = \left(\frac{16}{4}\right)^{\frac{1}{2}}$$

$$t = 2s$$

 $\therefore y = 10 \times 2 + (2)^2 = 24m$

(b) What is the speed of the particle at the time?

Ans:

Velocity of the particle is.

$$\vec{v}(t) = 8.0t\hat{i} + 2.0t\hat{i} + \vec{u}$$

At $t = 2s$
 $\vec{v}(t) = 8.0 \times 2\hat{i} + 2.0 \times 2\hat{j} + 10\hat{j}$
 $= 16\hat{i} + 14\hat{j}$

$$\therefore$$
 Speed of the particle

$$|\vec{v}| = \sqrt{(16)^2 + (14)^2}$$

= $\sqrt{256 + 196}$
= $\sqrt{452}$
= 21.26 ms⁻¹

19. \hat{i} and \hat{j} are unit vectors along x- and y-axis respectively. What is the magnitude and direction of the vectors $\hat{i}+\hat{j}$ and $\hat{i}.\hat{j}$? What are the components of a vector $a = 2\hat{i}+3\hat{j}$ along the directions of $\hat{i}+\hat{j}$ and $\hat{i}-\hat{j}$?

Ans: Consider a vector \vec{P}

$$\vec{P} = \hat{i} + \hat{j}$$
$$P_x \hat{i} + P_y \hat{j} = \hat{i} + \hat{j}$$

Compare the components on both sides.

$$P_{x} = P_{y} = 1$$
$$\left| \vec{P} \right| = \sqrt{P_{x}^{2} + P_{y}^{2}}$$
$$\left| \vec{P} \right| = \sqrt{1^{2} + 1^{2}}$$
$$\left| \vec{P} \right| = \sqrt{2} \qquad \dots(i)$$

So, the magnitude of the vector $\hat{i} + \hat{j}$ is $\sqrt{2}$

Let θ be the angle made by \vec{P} , with the x-axis.

So,

$$\tan \theta = \left(\frac{P_x}{P_y}\right)$$
$$\theta = \tan^{-1}\left(\frac{1}{1}\right)$$
$$\theta = 45^{\circ} \dots(ii)$$

 $\theta = 45^{\circ}$

So, the vector $\hat{i} + \hat{j}$ makes an angle of 45° with the x-axis

Let θ be the angle made by \vec{Q} , with the x-axis.

$$\vec{Q} = \hat{i} - \hat{j}$$

$$Q_x \hat{i} - Q_y \hat{j} = \hat{i} - \hat{j}$$

$$Q_x + Q_y = 1$$

$$|\vec{Q}| = \sqrt{Q_x^2 + Q_y^2}$$

$$|\vec{Q}| = \sqrt{2}$$

So, the magnitude of the vector $\hat{i} - \hat{j}$ is $\sqrt{2}$.

Let θ be the angle made by the vector \vec{Q} , with the x-axis.

$$\therefore \tan \theta = \left(\frac{Q_y}{Q_x}\right)$$
$$\theta = -\tan^{-1}\left(-\frac{1}{1}\right)$$
$$\theta = -45^\circ$$

So, the vector $\hat{i} - \hat{j}$ makes and angle of -45° with the axis. Compare the coefficients of \hat{i} and \hat{j}

$$\vec{A} = 2\hat{i} + 3\hat{j}$$
$$A_x\hat{i} + A_y\hat{j} = 2\hat{i} + 3\hat{j}$$

Let \overline{A} make an angle θ with the x-axis

 $\therefore \tan \theta = \left(\frac{A_x}{A_y}\right)$ $\theta = \tan^{-1}\left(\frac{3}{2}\right)$ $= \tan^{-1}(1.5)$ $= 56.31^\circ$

Angle between $(2\hat{i}+3\hat{j})$ and $(\hat{i}+\hat{j})$

 $\theta = 56.31 - 45 = 11.31^{\circ}$

 \vec{A} , along the direction of \vec{P} making an angle θ'

 $\therefore \tan \theta = \left(\frac{A_x}{A_y}\right)$

 $\tan\theta = (A\cos\theta)P$

 $\tan \theta = (A \cos 11.31) \frac{(\hat{i} + \hat{j})}{\sqrt{2}}$ $\tan \theta = \sqrt{13} \times \frac{0.9806}{\sqrt{2}} (\hat{i} + \hat{j})$ $\tan \theta = 2.5(\hat{i} + \hat{j})$ $\tan \theta = \frac{25}{10} \times \sqrt{2}$ $\tan \theta = \frac{5}{\sqrt{2}} \dots (v)$

Let θ be the angle between $(2\hat{i}+3\hat{j})$ and $(\hat{i}+\hat{j})$

$$\theta'' = 45 + 56.31 = 101.31^{\circ}$$

Component of vector A, along the direction of Q, making an angle θ

$$= (A\cos\theta)\vec{Q}$$
$$= (A\cos\theta)\frac{(\hat{i}-\hat{j})}{\sqrt{2}}$$
$$= \sqrt{13}\cos\left(901.31^\circ\right)\frac{(\hat{i}+\hat{j})}{\sqrt{2}}$$
$$= -\sqrt{\frac{13}{2}}\sin 11.30^\circ(\hat{i}-\hat{j})$$
$$= -2.550 \times 0.1961(\hat{i}-\hat{j})$$
$$= -0.5(\hat{i}-j)$$
$$= -\frac{5}{10} \times \sqrt{2}$$
$$= -\frac{1}{\sqrt{2}}$$

20. For any arbitrary motion in space, which of the following relations are true.

(a)
$$v_{average} = \left(\frac{1}{2}\right) \left(v(t_1) + v(t_2)\right)$$

Ans: False. As the motion of the particle is arbitrary, the average velocity of the particle cannot be given by this equation.

(b)
$$v_{average} = \frac{[r(t_2)-r(t_1)]}{(t_2-t_1)}$$

Ans: True. The arbitrary motion of the particle can be represented by the given equation.

(c) v(t)=v(0)+at

Ans: False. The motion of the particle is arbitrary. The acceleration of the particle may also be non-uniform. So, this equation cannot represent the motion of the particle in space.

(**d**)
$$r(t)=r(0)+v(0)t+\left(\frac{1}{2}\right)at^{2}$$

Ans: False. The motion of the particle is arbitrary, acceleration of the particle may also be non-uniform. So, this equation cannot represent the motion of particle in space.

(e)
$$a_{average} = \frac{[v(t_2)-v(t_1)]}{(t_2-t_1)}$$

(The 'average' stands for average of the quantity over the time interval t_1 to t_2)

- **Ans:** True. The arbitrary motion of the particle can be represented by the given equation.
- 21. Read each statement below carefully and state, with reasons and examples, if it is true or false.

A scalar quantity is one that.

- (a) Is conserved in a process.
- Ans: False. Energy is not conserved in inelastic collisions.

(b) Can never take negative values.

Ans: False. Temperature can take negative values.

(c) Must be dimensionless.

Ans: False. Total path length is a scalar quantity and has the dimension of length.

(d) Does not vary from one point to another in space.

Ans: False. A scalar quantity like gravitational potential can vary from one point to another in space.

(e) Has the same value for observers with different orientations of axes

- **Ans:** True. The value of a scalar does not change for observers with different orientations of axes.
- 22. An aircraft is flying at a height of 3400m above the ground. If the angle subtended at a ground observation point by the aircraft positions 10.0s apart is 30°, what is the speed of the aircraft?

Ans: The positions of the observer and the aircraft are shown below.



Height of the aircraft from ground, OR = 3400 m

Angle subtended between the positions, $\angle POQ = 30^{\circ}$

Time =10s

In $\triangle PRO$:

 $\tan 15^{\circ} = \frac{PR}{OR}$

 $PR = OR \tan 15^{\circ}$

 $PR = 3400 \times \tan 15^{\circ}$

 \triangle PRO is similar to $\triangle RQO$

 \therefore PR = RQ

PQ = PR + RQ

PQ = 2PR

 $PQ = 2 \times 3400 \tan 15^{\circ}$

 $PQ = 6800 \times 0.268$

 $PQ = 1822.4 \,\mathrm{m}$

 $\therefore \text{ Speed of the aircraft} = \frac{1822.4}{10} = 182.24 \text{ m/s}$