5 Chapter

Exercise 5.1

- 1. Prove that f(x)=5x-3 is a continuous function at x=0, x=-3 and x=5.
- Ans: The given function is f(x)=5x-3.

continuity and differentiabilituy

At x=0, $f(0)=5\times 0-3=3$.

Taking limit as $x \rightarrow 0$ both sides of the function give

$$\lim_{x \to 0} f(x) = \lim_{x \to 0} (5x-3) = 5 \times 0-3 = 3$$

$$\therefore \lim_{\mathbf{x}\to 0} \mathbf{f}(\mathbf{x}) = \mathbf{f}(0).$$

Thus, f satisfies continuity at x=0.

Again, at x=-3, $f(-3)=5\times(-3)-3=-18$.

Now, taking limit as $x \rightarrow 3$ both sides of the function give

$$\lim_{x \to 3} f(x) = \lim_{x \to 3} f(5x-3) = 5 \times (-3) - 3 = -18$$

$$\therefore \lim_{x \to 3} f(x) = f(-3).$$

Therefore, f satisfies continuity at x=-3.

Also, at
$$x=5, f(x)=f(5)=5\times 5-3=25-3=22$$
.

Taking limit as $x \rightarrow 5$ both sides of the function give

$$\lim_{x \to 5} f(x) = \lim_{x \to 5} (5x-3) = 5 \times 5 - 3 = 22$$
$$\therefore \lim_{x \to 5} f(x) = f(5).$$

Hence, f satisfies continuity at x=5.

2. Verify whether the function $f(x)=2x^2-1$ is continuous at x=3.

Ans: The given function is $f = 2(x^2)1$.

Now, at x=3, $f(3)=2\times 3^2-1=17$.

Taking limit as $x \rightarrow 3$ both sides of the function give

$$\lim_{x \to 3} f(x) = \lim_{x \to 3} (2x^2 - 1) = 2 \times 3^2 - 1 = 17$$

$$\therefore \lim_{x \to 3} f(x) = f = (3).$$

Hence, f satifies continuity at x=3.

3. Verify whether the following functions are continuous.

(a)
$$f(x) = x-5$$

Ans: The given function is f(x) = x-5.

It is assured that for every real number k, f is defined and its value at k is k-5. Also, it can be noted that

$$\lim_{x \to k} f(x) = \lim_{x \to k} f(x-5) = k = k-5 = f(k).$$

$$\therefore \lim_{x \to k} f(x) = f(k)$$

Hence, f satisfies continuity at every real number and so, it is a continuous function.

(b)
$$f(x) = \frac{1}{x-5}, x \neq 5$$

Ans: The given function is

$$f(x) = \frac{1}{x-5}.$$

Let $k \neq 5$ is any real number, then taking limit as $x \rightarrow k$ both sides of the function give

$$\lim_{x \to k} f(x) = \lim_{x \to k} \frac{1}{x \cdot 5} = \frac{1}{k \cdot 5}$$

Also, $f(k) = \frac{1}{k \cdot 5}$, since $k \neq 5$
 $\therefore \lim_{x \to k} f(x) = f(k)$

Therefore, f satisfies continuity at every point in the domain of f and so, it is a continuous function.

(c)
$$f(x) = \frac{x^2 - 25}{x + 5}, x \neq 5$$

Ans: The given function is

$$f(x) = \frac{x^2 - 25}{x + 5}, x \neq 5$$

Now let $c \neq -5$ be any real number, then taking limit as $x \rightarrow c$ on both sides of the function give

$$\lim_{x \to c} f(x) = \lim_{x \to c} \frac{x^2 - 25}{x + 5} = \lim_{x \to c} \frac{(x + 5)(x - 5)}{x + 5} = \lim_{x \to c} (x - 5) = (c - 5)$$

Again, $f(c) = \frac{(c + 5)(c - 5)}{c + 5} = c(c - 5)$, since $c \neq 5$.

Hence, f satisfies continuity at every point in the domain of f and so it is a continuous function.

(d) f(x) = |x - 5|

Ans: The given function is $f(x) = |x-5| = \begin{cases} 5-x, & \text{if } x < 5 \\ x-5, & \text{if } x > 5 \end{cases}$.

Note that, f is defined at all points in the real line. So, let assume c be a point on a real line.

Then, we have c < 5 or c = 5 or c > 5.

Now, let discuss these three cases one by one.

Case (i): c<5

Then, the function becomes f(c)=5-c.

Now,
$$\lim_{x\to c} f(x) = \lim_{x\to c} (5-x) = 5-c$$

$$\therefore \lim_{x\to c} f(x) = f(c).$$

Therefore, f is continuous at all real numbers which are less than .5..

Case (ii): c=5Then, f(c)=f(5)=(5-5)=0.

Now,

 $\lim_{x \to 5^{-}} f(x) = \lim_{x \to 5} (5 - x) = (5 - 5) = 0 \text{ and}$ $\lim_{x \to 5^{+}} f(x) = \lim_{x \to 5} (x - 5) = 0.$

Therefore, we have

 $\lim_{x\to c^{-}} f(x) = \lim_{x\to c^{+}} f(x) = f(c).$

Thus, f satisfies continuity at x=5, and so f is continuous at x=5.

Case (iii): c>5

Then we have, f(c)=f(5)=c-5.

Now,

$$\lim_{x\to c} f(x) = \lim_{x\to c} (x-5) = c-5.$$

Therefore,

 $\lim_{x\to c} f(x) = f(c).$

So, f is continuous at all real numbers that are greater than 5.

Thus, f satisfies continuity at every real number and hence, it is a continuous function.

4. Prove that $f(x)=x^n$ is continuous at x=n, where n is a positive integer.

Ans: The given function is $f(x)=x^n$.

We noticed that the function f is defined at all positive integers n and also its value at x=n is n^n .

Therefore,
$$\lim_{x\to n} f(n) = \lim_{x\to n} f(x^n) = n^n$$

So, $\lim_{x\to n} f(x) = f(n)$.

Thus, the function $f(x)=x^n$ is continuous at x=n, where n is a positive integer.

5. Verify whether the following function f is continuous at x=0, x=1 and at x=2.

$$f(x) = \begin{cases} x, \text{ if } x \le 1 \\ 5, \text{ if } x > 1 \end{cases}$$

Ans: The given function is $f(x) = \begin{cases} x, & \text{if } x \le 1 \\ 5, & \text{if } x > 1 \end{cases}$.

It is obvious that the function f is defined at x=0 and its value at x=0 is 0. Now, $\lim_{x\to 0} f(x) = \lim_{x\to 0} x=0$. So, $\lim_{x\to 0} f(x) = f(0)$.

Hence, the function f satisfies continuity at x=0.

It can be observed that f is defined at x=1 and its value at this point is 1.

Now, the left-hand limit of the function f at x=1 is

$$\lim_{x\to 1^{-}} f(x) = \lim_{x\to 1^{-}} x = 1$$

Also, the right-hand limit of the function f at x=1 is

 $\lim_{x \to l^+} f(x) = \lim_{x \to l^+} f(5)$

Therefore, $\lim_{x\to 1^-} f(x) \neq \lim_{x\to 1^+} f(x)$.

Thus, f is not continuous at x=1

It can be found that f is defined at x=2 and its value at this point is 5.

```
That is, \lim_{x \to 2} f(x) = \lim_{x \to 2} f(5) = 5.
```

Therefore,
$$\lim_{x\to 2} f(x) = f(2)$$

Hence, f satisfies continuity at x=2.

6. Locate all the discontnuity points for the function f, where f is given by

$$f(x) = \begin{cases} 2x+3, & \text{if } x \le 2 \\ 2x-3, & \text{if } x > 2 \end{cases}.$$

Ans: The given function is $f(x) = \begin{cases} 2x+3, & \text{if } x \le 2\\ 2x-3, & \text{if } x > 2 \end{cases}$.

It can be observed that the function f is defined at all the points in the real line. Let consider c be a point on the real line. Then, three cases may arise.

- I. c<2
- II. c>2
- III. c=2

Case (i): When c<2

Then, we have $\lim_{x\to c} f(x) = \lim_{x\to\infty} (2x+3) = 2c+3$.

Therefore,

 $\lim_{x\to c} f(x) = f(c).$

Hence, f attains continuity at all points x, where x < 2.

Case (ii): When c>2

```
Then, we have f(c)=2c-3.
```

So,

 $\lim_{x\to c} f(x) = \lim_{x\to\infty} (2x-3) = 2c-3.$

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Hence, f satisfies continuity at all points x , where x>2.

Case(iii): When c=2

Then, the left-hand limit of the function f at x=2 is

$$\lim_{x \to 2^{-}} f(x) = \lim_{x \to 2^{-}} (2x+3) = 2 \times 2 + 3 = 7 \text{ and}$$

the right-hand limit of the function f at x=2 is,

 $\lim_{x \to 2^+} f(x) = \lim_{x \to 2^+} (2x+3) = 2 \times 2 - 3 = 1.$

Thus, at x=2, $\lim_{x\to 2^{-}} f(x) \neq \lim_{x\to 2^{+}} f(x)$.

So, the function f does not satisfy continuity at x=2.

Hence, x=2 is the only point of discontinuity of the function f(x).

7. Locate all the discontinuity points for the function f, where f is given by

$$f(x) = \begin{cases} |x|+3, & \text{if } x \le -3 \\ -2x, & \text{if } -3 < x < 3 \\ 6x+2, & \text{if } x \ge 3 \end{cases}.$$

Ans: The given function is
$$f(x) = \begin{cases} |x|+3, & \text{if } x \le -3 \\ -2x, & \text{if } -3 < x < 3 \\ 6x+2, & \text{if } x \ge 3 \end{cases}$$
.

Observe that, f is defined at all the points in the real line.

Now, let assume c as a point on the real line.

Then five cases may arise. Either c<-3, or c=-3 or -3< c<3, or c=3, or c>3.

Let's discuss the five cases one by one.

Case I: When c<-3

Then,
$$f(c) = -c+3$$
 and $\lim_{x \to c} f(x) = \lim_{x \to c} (-x+3) = -c+3$.

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Hence, f satisfies continuity at all points x, where x < -3.

Case II: When c=-3

Then, f(-3) = -(-3) + 3 = 6.

Also, the left-hand limit

$$\lim_{x\to 3^{-}} f(x) = \lim_{x\to 3^{-}} (-x+3) = -(-3) + 3 = 6.$$

and the right-hand limit

 $\lim_{x\to 3^{+}} f(x) = \lim_{x\to 3^{+}} f(-2x) = 2x(-3) = 6.$

Therefore, $\lim_{x\to 3} f(x) = f(-3)$.

Hence, f satisfies continuity at x=-3.

Case III: When -3<c<3

Then, f(c) = -2c and also $\lim_{x \to c} f(x) = \lim_{x \to 3c} (-2x) = -2c$.

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Hence, f satisfies continuity at x, where -3 < x < 3.

Case IV: When c=3

Then, the left-hand limit of the function f at x=3 is

 $\lim_{x \to 3^{-}} f(x) = \lim_{x \to 3^{-}} f(-2x) = -2 \times 3 = 6 \text{ and}$

the right-hand limit of the function f at x=3 is

 $\lim_{x\to 3^{+}} f(x) = \lim_{x\to 3^{+}} f(6x+2) = 6 \times 3 + 2 = 20.$

Thus, at x=3, $\lim_{x\to 3^{-}} f(x) \neq \lim_{x\to 3^{+}} f(x)$.

Hence, f does not satisfy continuity at x=3.

Case V: When c>3.

Then f(c)=6c+2 and also

$$\lim_{x\to c} f(x) = \lim_{x\to c} (6x+2) = 6c+2.$$

Therefore, $\lim_{x\to c} f(x) = f(c)$.

So, f satisfies continuity at all points x, when x>3.

Thus, x=3 is the only point of discontinuity of the function f.

8. Locate all the discontnuity points for the function f, where f is given by

$$f(x) = \begin{cases} \frac{|x|}{x}, & \text{if } x \neq 0\\ 0, & \text{if } x=0 \end{cases}.$$

Ans: The given function is $f(x) = \begin{cases} \frac{|x|}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$.

Now, f(x) can be rewritten as

$$f(x) = \begin{cases} \frac{|x|}{x} = \frac{-x}{x} = -1 \text{ if } x \neq 0\\ 0, \text{ if } x = 0\\ \frac{|x|}{x} = \frac{x}{x} = 1 \text{ if } x > 0 \end{cases}$$

It can be noted that the function f is defined at all points of the real line.

Now, let assume c as a point on the real line.

Then three cases may arise, either c<0, or c=0, or c>0.

Let discuss three cases one by one.

Case I: When c<0.

Then, f(c) = -1 and

 $\lim_{x\to c} f(x) = \lim_{x\to c} (-1) = -1.$

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Hence, f satisfies continuity at all the points x where x < 0.

Case II: When c=0.

Then, the left-hand limit of the function f at x=0 is

$$\lim_{x \to 0^{-}} f(x) = \lim_{x \to 0^{-}} (-1) = -1 \text{ and}$$

the right-hand limit of the function f at x=0 is

$$\lim_{x \to 0^{+}} f(x) = \lim_{x \to 0^{+}} (1) = 1.$$

At x=0,
$$\lim_{x \to 0^{-}} f(x) \neq \lim_{x \to 0^{+}} f(x).$$

Hence, the function f does not satisfy continuity at x=0.

Case III: When c>0. Then f(c)=1 and also $\lim_{x\to c} f(x) = \lim_{x\to c} (1)=1.$ Therefore, $\lim_{x\to c} f(x) = f(c).$

So, the function f is continuous at all the points x, for x>0.

Thus, x=0 is the only point of discontinuity for the function f.

9. Locate all the discontinuity points for the function f, where f is given by $f(x) = \left\{ \frac{x}{|x|}, \text{ if } x < 0 \\ -1, \text{ if } x \ge 0 \right\}.$

Ans: The given function is $f(x) = \begin{cases} \frac{x}{|x|}, & \text{if } x < 0 \\ -1, & \text{if } x \ge 0 \end{cases}$.

Now, we know that, if x < 0, then |x| = -x.

Therefore, the f(x) can be written as

$$f(x) = \begin{cases} \frac{x}{|x|}, & \text{if } x < 0 \\ -1, & \text{if } x \ge 0 \end{cases}.$$

 \Rightarrow f(x)=-1 for all positive real numbers.

Now, let assume c as any real number.

Then, we have $\lim_{x\to c} f(x) = \lim_{x\to c} (-1) = -1$ and

 $f(c) = -1 = \lim_{x \to e} f(x).$

Therefore, the function f(x) is a continuous function.

Thus, there does not exist any point of discontinuity.

10. Locate all the discontinuity points for the function f, where f is given by $f(x) = \begin{cases} x+1, & \text{if } x \ge 1 \\ x^2+1, & \text{if } x < 1 \end{cases}.$

Ans: The given function is

$$f(x) = \begin{cases} x+1, \text{ if } x^3 \\ x^2+1, \text{ if } x < 1 \end{cases}$$

Note that, f(x) is defined at all the points of the real line.

Now, let assume c as a point on the real line.

Then three cases may arise, either c<1, or c=1, or c>1.

Let discuss the three cases one by one.

Then,
$$f(c)=c^2+1$$
 and also

$$\lim_{x\to c} f(x) = \lim_{x\to c} f(x^2+1) = c^2+1.$$

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Hence, f satisfies continuity at all the points x, where x < 1.

Case II: When c=1.

Then, we have f(c)=f(1)=1+1=2.

Now, the left-hand limit of f at x=1 is

 $\lim_{x \to 1^{+}} f(x) = \lim_{x \to 1^{+}} (x^{2}+1) = 1^{2}+1=2 \text{ and the right-hand limit of } f \text{ at } x=1 \text{ is,}$ $\lim_{x \to 1^{+}} f(x) = \lim_{x \to 1^{+}} (x^{2}+1) = 1^{2}+1=2.$ Therefore, $\lim_{x \to 1} f(x) = f(c).$ Hence, f satisfies continuity at x=1. Case III: When c>1. Then, we have f(c)=c+1 and $\lim_{x \to c} f(x) = \lim_{x \to c} (x+1)=c+1.$ Therefore, $\lim_{x \to c} f(x) = f(c).$ So, f satisfies continuity at all the points x, where x>1. Hence, there does not exist any discontnuity points.

- 11. Locate all the discontnuity points for the function f, where f is given by $f(x) = \begin{cases} x^3 - 3, & \text{if } x \le 2 \\ x^2 + 1, & \text{if } x > 2 \end{cases}.$
- Ans: The given function is $f(x) = \begin{cases} x^3 3, & \text{if } x \le 2 \\ x^2 + 1, & \text{if } x > 2 \end{cases}$.

Observe that, the function f is defined at all points in the real line.

Now, let assume c as a point on the real line.

Case I: When c<2.

Then, we have $f(c)=c^3-3$ and also $\lim_{x\to c} f(x)=\lim_{x\to c} (x^3-3)=c^3-3$.

Therefore, the function f attains continuity at all the points x, where x < 2.

Case II: When c=2.

Then, we have $f(c)=f(2)=2^{3}-3=5$.

Now the left-hand limit of the function is

 $\lim_{x \to 2^{-}} f(x) = \lim_{x \to 2^{-}} (x^{3} - 3) = 2^{3} - 3 = 5 \text{ and the right-hand limit is}$ $\lim_{x \to 2^{+}} f(x) = \lim_{x \to 2^{+}} (x^{2} + 1) = 2^{2} + 1 = 5.$ Therefore, $\lim_{x \to 2} f(x) = f(2).$ Hence, the function f is continuous at x=2. Case III: When c>2. Then, f(c)=c^{2} + 1 and $\lim_{x \to c} f(x) = \lim_{x \to c} (x^{2} + 1) = c^{2} + 1.$ Therefore, $\lim_{x \to c} f(x) = f(c).$

So, f attains continuity at all the points x, where x > 2.

Thus, the function f is continuous at all the points on the real line.

Hence, f does not have any point of discontinuity.

12. Locate all the discontinuity points for the function f, where f is given by $f(x)\begin{cases} x^{10}-1, & \text{if } x \le 1 \\ x^2, & \text{if } x > 1 \end{cases}.$

Ans: The given function is
$$f(x) \begin{cases} x^{10}-1, \text{ if } x \leq 1 \\ x^2, \text{ if } x > 1 \end{cases}$$
.

Observe that, the function f is defined at every point of the real line.

Now, let assume c as a point on the real number line.

Case I: When c<1. Then $f(c)=c^{10}-1$. Also, $\lim_{x\to c} f(x) = \lim_{x\to c} (x^{10}-1)=c^{10}-1$ Therefore, $\lim_{x\to c} f(x)=f(c)$.

Hence, the function f attains continuity at every point x, for x<1. Case II: When c=1.

Then the left-hand limit of the function f(x) at x=1 is

$$\lim_{x \to 1^{-}} f(x) = \lim_{x \to 1^{-}} (x^{10}-1) = 10^{10}-1 = 1-1 = 0 \text{ and}$$

the right-hand limit of the function f at x=1 is

$$\lim_{x \to 1^{+}} f(x) = \lim_{x \to 1^{+}} (x^{2}) = 1^{2} = 1.$$

So, we can notice that, $\lim_{x\to l^-} f(x) \neq \lim_{x\to l^+} f(x)$.

Hence, the function f does not satisfy continuity at x=1.

Case III: When
$$c>1$$
.

Then, $f(c)=c^2$.

Also,
$$\lim_{x\to c} f(x) = \lim_{x\to c} (x^2) = c^2$$
.

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Thus, the function f attains continuity at every point x, for x>1.

Hence, we can conclude that x>1 is the only point of discontinuity for the function f.

13. Verify whether the function $f(x) = \begin{cases} x+5, & \text{if } x \le 1 \\ x-5, & \text{if } x > 1 \end{cases}$ is continuous.

Ans: The given function is $f(x) = \begin{cases} x+5, & \text{if } x \le 1 \\ x-5, & \text{if } x > 1 \end{cases}$.

It can be noted that the function f is defined at every point on the real line.

Now, let assume c as a point on the real line.

Case I: When c<1.

Then, $f(c) = c^{10} - 1$.

Also, $\lim_{x\to c} f(x) = \lim_{x\to c} (x^{10} - 1) = c^{10} - 1.$

Hence, f satisfies continuity at every point x, for x < 1.

Case II: When c=1.

Then, f(1)=1+5=6.

Now, the left-hand limit of the function f at x=1 is

 $\lim_{x \to 1^{-}} f(x) = \lim_{x \to 1^{-}} (x+5) = 1+5=6 \text{ and}$

the right-hand limit of the function at x=1 is $\lim_{x \to 1^+} f(x) = \lim_{x \to 1^+} (x-5) = 1-5=4$.

Thus, it is seen that, $\lim_{x\to 1^-} f(x) \neq \lim_{x\to 1^+} f(x)$.

Hnece, f does not attain continuity at x=1.

Case III: When c>1.

Then f(c)=c-5.

Also, $\lim_{x\to c} f(x) = \lim_{x\to c} (x-5) = c-5$.

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Thus, the function f is continuous at every point x, for x>1.

Hence, we can conclude that x=1 is the only point of discontinuity for the function f.

14. Verify whether the following function **f** is continuous.

$$f(x) = \begin{cases} 3, \text{ if } 0 \le x \le 1 \\ 4, \text{ if } 1 < x < 3 \\ 5, \text{ if } 3 \le x \le 10 \end{cases}$$

Ans: The given function is $f(x) = \begin{cases} 3, \text{ if } 0 \le x \le 1 \\ 4, \text{ if } 1 < x < 3 \\ 5, \text{ if } 3 \le x \le 10 \end{cases}$.

Therefore, f is defined in the interval [0,10].

Now let assume c as a point in the interval [0,10].

Then there may arise five cases.

Case I: When $0 \le c < 1$.

Then f(c)=3.

```
Also, \lim_{x\to c} f(x) = \lim_{x\to c} (3) = 3.
```

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Hence, the function f attains continuity at the interval [0,1].

```
Case II: When c=1.
```

Then f(3)=3.

Also, the left-hand-limit of the function at x=1 is

 $\lim_{x \to 1^{-}} f(x) = \lim_{x \to 1^{-}} (3) = 3 \text{ and the right-hand-limit of the function at } x=1 \text{ is}$ $\lim_{x \to 1^{+}} f(x) = \lim_{x \to 1^{+}} (4) = 4.$

Thus, it is noticed that $\lim_{x\to 1^-} f(x) \neq \lim_{x\to 1^+} f(x)$.

Hence, the function f does not satisfy continuity at x=1.

Case III: When 1 < c < 3.

Then f(c)=4. Also, $\lim_{x\to c} f(x) = \lim_{x\to c} (4)=4$. Thus, $\lim_{x\to c} f(x) = f(c)$.

Hence, the function f attains continuity at every point in the interval [1,3]. Case IV: When c=3.

Then f(c)=5.

Now, the left-hand-limit of the function f at x=3 is

 $\lim_{x \to 3^{-}} f(x) = \lim_{x \to 3^{-}} (4) = 4 \text{ and the right-hand-limit of the function } f \text{ at } x = 3 \text{ is}$ $\lim_{x \to 3^{+}} f(x) = \lim_{x \to 3^{+}} (5) = 5.$

Therefore, it is noted that $\lim_{x\to 3^{+}} f(x) \neq \lim_{x\to 3^{+}} f(x)$.

Hence, the function f is not continuous at x=3.

Case V: When $3 < c \le 10$.

Then f(c)=5.

Also, $\lim_{x\to c} f(x) = \lim_{x\to c} (5) = 5$.

Therefore,
$$\lim_{x\to c} f(x) = f(c)$$
.

So, the function f attains continuity at every point in the interval [3,10].

Hence, the function f is not continuous at x=1 and x=3.

15. Verify whe ther the following function f is continuous. f such that $f(x) = \begin{cases} 2x, & \text{if } x < 0 \\ 0, & \text{if } 0 \le x \ne 1 \\ 4x, & \text{if } x > 1 \end{cases}$

Ans: The given function is $f(x) = \begin{cases} 2x, & \text{if } x < 0 \\ 0, & \text{if } 0 \le x \ne 1 \\ 4x, & \text{if } x > 1 \end{cases}$.

Now, let consider c be a point on the real number line.

Then, five cases may arrive.

Case I: When c<0.

Then, f(c)=2c.

Also,
$$\lim_{x \to c} f(x) = \lim_{x \to c} (2x) = 2c$$
.
Therefore, $\lim_{x \to c} f(x) = f(c)$.

Hence, the function f attains continuity at every point x whenever x < 0.

Case II: When
$$c = 0$$
.

Then,
$$f(c) = f(0) = 0$$
.

Now, the left-hand-limit of the function f at x = 0 is

 $\lim_{x\to 0^{-}} f(x) = \lim_{x\to 0^{-}} (2x) = 0 \text{ and the right-hand limit of the function } f \text{ at } x = 0 \text{ is,}$ $\lim_{x\to 0^{+}} (x) = \lim_{x\to 0^{+}} (0) = 0.$ Therefore, $\lim_{x\to 0} f(x) = f(0).$ Thus, the function f attains continuity at x = 0. Case II: When 0<c<1

Then, f(x)=0.

Also, $\lim_{x\to c} f(x) = \lim_{x\to c} (0) = 0$.

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Hence, f attains continuity at every point in the interval (0,1).

Case IV: When c = 1.

Then, f(c)=f(1)=0.

Now, the left-hand-limit at x = 1 is

 $\lim_{x\to l^-} f(x) = \lim_{x\to l^-} (0) = 0 \text{ and the right-hand-limit at } x = 1 \text{ is}$

 $\lim_{x \to 1^+} f(x) = \lim_{x \to 1^+} (4x) = 4 \times 1 = 4.$

Thus, it is noticed that, $\lim_{x\to l^-} f(x) \neq \lim_{x\to l^+} f(x)$.

Hence, the function f is not continuous at x = 1.

Case V: When c < 1.

Then, f(c)=f(1)=0.

Also, $\lim_{x\to c} f(x) = \lim_{x\to c} (4x) = 4c$

Therefore, $\lim_{x\to c} f(x) = f(c)$.

So, the function f attains continuity at every point x, for x>1.

Hence, the function f is discontinuous only at x = 1.

16. Verify whether the function f is continuous. Provided that f is defined by $f(x) = \begin{cases} -2, & \text{if } x \le -1 \\ 2x, & \text{if } -1 < x \le 1 \\ 2, & \text{if } x > 1 \end{cases}$. Ans: The given function is $f(x) = \begin{cases} -2, \text{ if } x \leq -1 \\ 2x, \text{ if } -1 < x \leq 1 \\ 2, \text{ if } x > 1 \end{cases}$.

Note that, f is defined at every point in the interval $[-1,\infty)$.

Now, let assume c is a point on the real number line.

Case I: When c<-1.

Then, f(c)=-2.

Also, $\lim_{x\to c} f(x) = \lim_{x\to c} (-2) = -2$.

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Hence, the function f attains continuity at every point x , for x < -1.

Case II: When c=-1.

Then, f(c)=f(-1)=-2.

Now, the left-hand-limit of the function at x=-1 is

 $\lim_{x \to 1^{-}} f(x) = \lim_{x \to 1^{-}} (-2) = -2 \text{ and the right-hand-limit at } x = -1 \text{ is}$ $\lim_{x \to 1^{+}} (x) = \lim_{x \to 1^{+}} = 2 \times (-1) = -2.$ Therefore, $\lim_{x \to -1} f(x) = f(-1).$ Hence, the function f satisfies continuity at x=-1. Case III: When -1 < c < 1. Then, f(c) = 2c and $\lim_{x \to c} f(x) = \lim_{x \to c} (2x) = 2c.$ Therefore, $\lim_{x \to c} f(x) = f(c).$

Hence, the function f attains continuity at every point in the interval (-1,1). Case IV: When c=1. Then, $f(c)=f(1)=2\times 1=2$

Now, the left-hand-limit of the function at x = 1 is $\lim_{x \to 1^{+}} f(x) = \lim_{x \to 1^{+}} (2x) = 2 \times 1 = 2$ and the right-hand-limit at x = 1 is $\lim_{x \to 1^{+}} f(x) = \lim_{x \to 1^{+}} 2 = 2.$ Therefore, $\lim_{x \to 1} f(x) = f(c)$. Thus, the function f attains continuity at x = 2. Case V: When c > 1. Then f(c) = 2. Also, $\lim_{x \to 2} f(x) = \lim_{x \to 2} (2) = 2$. Therefore, $\lim_{x \to 2} f(x) = f(c)$.

Hence, the function f is continuous at every point x, for x>1.

17. Formulate a relationship between a and b so that the function f defined by $f(x) = \begin{cases} ax+1, & if \ x \le 3 \\ bx+3, & if \ x > 3 \end{cases}$ is continuous at x=3.

Ans: The given function is $f(x) = \begin{cases} ax+1, & \text{if } x \le 3 \\ bx+3, & \text{if } x > 3 \end{cases}$.

The function f will be continuous at x = 3 if

 $\lim_{x \to 3^{-}} f(x) = \lim_{x \to 3^{-}} f(x) = f(3), \qquad \dots \dots (1)$ $\lim_{x \to 3^{-}} f(x) = \lim_{x \to 3^{+}} f(ax+1) = 3a+1,$ $\lim_{x \to 3^{+}} f(x) = \lim_{x \to 3^{+}} f(bx+1) = 3b+3, \qquad \dots \dots (2)$ and

$$f(3)=3a+1.$$
(3)

Therefore, from the equation (1), (2), and (3) gives

3a+1=3b+3=3a+1 $\Rightarrow 3a+1=3b+3$ $\Rightarrow 3a=3b+2$ $\Rightarrow a=b+\frac{2}{3}$

Hence, the required relationship between a and b is given by $a=b+\frac{2}{3}$.

18. Determine the value of λ for which the function defined by $f(x) = \begin{cases} \lambda(x^2 - 2x), & \text{if } x \le 0 \\ 4x + 1, & \text{if } x > 0 \end{cases}$ is continuous at x = 0. Also discuss the continuity of f at x = 1?

Ans: The given function is $f(x) = \begin{cases} \lambda(x^2-2x), \text{ if } x \le 0 \\ 4x+1, \text{ if } x > 0 \end{cases}$.

Now the function will be continuous at x = 0 if

$$\lim_{x \to 0^{-}} f(x) = \lim_{x \to 0^{-}} f(x) = f(0)$$
$$\Rightarrow \lim_{x \to 0^{-}} \lambda(x^2 - 2x) = \lim_{x \to 0^{-}} (4x + 1) = \lambda(0^2 - 2 \times 0)$$
$$\Rightarrow \lambda(0^2 - 2 \times 0) = 4 \times 0 + 1 = 0$$

 $\Rightarrow 0=1=0$, which is impossible.

Thus, there does not exist any value of λ for which f is continuous at x = 0.

Now, at
$$x = 1$$
,

 $f(1)=4x+1=4\times 1+1=5$ and

 $\lim_{x\to 1} (4x+1) = 4 \times 1 + 1 = 5$.

Therefore, $\lim_{x\to 1} f(x) = f(1)$.

Hence, the function f is continuous at x = 1, for all values of λ .

- 19. Prove that the function g(x)=x-[x] is not continuous at any integral point, where [x] denotes the greatest integer value of x that are less than or equal to x.
- **Ans:** The given function is g(x)=x-[x].

Note that, the function is defined at every integral point.

Now, let assume that n be an integer.

Then, g(n)=n-[n]=n-n=0.

Now taking left-hand-limit as $x \rightarrow n$ to the function g gives

 $\lim_{x \to n} g(x) = \lim_{x \to n} [x-[x]] = \lim_{x \to n} (x) - \lim_{x \to n-} [x] = n - (n-1) = 1.$

Again, the right-hand-limit on the function at x=n is

$$\lim_{x \to n^+} g(x) = \lim_{x \to n^+} [x-[x]] = \lim_{x \to n^+} (x) - \lim_{x \to n^+} [x] = n-n=0.$$

Note that, $\lim_{x\to n^-} g(x) \neq \lim_{x\to n^+} g(x)$.

Thus, the function f is cannot be continuous at x=n,

Hence, the function g is not continuous at any integral point.

20. Verify whether the function $f(x)=x^2-\sin x+5$ is continuous at $x = \pi$?

Ans: The given function is $f(x)=x^2-\sin x+5$.

Now, at $x=\pi$,

$$f(x)=f(\pi)=\pi^2-\sin\pi+5=\pi^2-0+5=\pi^2+5$$
.

Taking limit as $x \rightarrow \pi$ on the function f(x) gives

$$\lim_{x\to\pi} f(x) = \lim_{x\to\pi} (x^2 \operatorname{-sinx} + 5).$$

Now substitute $x=\pi+h$ into the function f(x).

When
$$x \rightarrow \pi$$
, then $h \rightarrow 0$.
Therefore,

$$\lim_{x \rightarrow \pi} f(x) = \lim_{x \rightarrow \pi} (x^2 - \sin x) + 5.$$

$$= \lim_{h \rightarrow 0} [(\pi + h^2) - \sin(\pi + h) + 5]$$

$$= \lim_{h \rightarrow 0} (\pi + h)^2 - \lim_{h \rightarrow 0} \sin(\pi + h) + \lim_{h \rightarrow 0} 5$$

$$= (\pi + 0)^2 - \lim_{h \rightarrow 0} [\sin\pi \cosh + \cos\pi \sinh] + 5$$

$$= \pi^2 - \lim_{h \rightarrow 0} \sin\pi \cosh - \lim_{h \rightarrow 0} \cos\pi \sinh + 5$$

$$= \pi^2 - \sin\pi \cos 0 - \cos\pi \sin 0 + 5$$

$$= \pi^2 - 0 \times 1 - (-1) \times 0 + 5 = \pi^2 + 5.$$
So, $\lim_{x \rightarrow x} f(x) = f(\pi).$

Hence, it is concluded that the function f is continuous at x=n.

21. Determine whether the following functions are continuous.

(a) $f(x) = \sin x + \cos x$ (b) $f(x) = \sin x - \cos x$ (c) $f(x) = \sin x \times \cos x$.

Ans: It is known that if two functions g and h are continuous, then g+h, g-h and g,h are also continuous.

So, let us assume that, g(x)=sinx and h(x)=cosx are two continuous functions.

Now, as g(x)=sinx is defined for every real number, so let c be a real number. Substitute x=c+h into the function g.

When $x \rightarrow c$, then $h \rightarrow 0$.

So, g(c)=sinc.

Also,

```
\lim_{x \to c} g(x) = \limsup_{x \to c} sinx
= \lim_{h \to 0} sin(c+h)
= \lim_{h \to 0} [sinccosh+coscsinh]
= \lim_{h \to 0} (sinccosh) + \lim_{h \to 0} (coscsinh)
= sinccos0 + coscsin0
= sinc+0
= sinc
```

```
Therefore, \lim_{x\to c} g(x) = g(c).
```

Hence, the function g is a continuous.

Again, let us assume that h(x) = cosx.

Note that, the function $h(x)=\cos x$ is defined for every real number.

Now, let c be a real number.

Substitute x=c+h into the function.

When $x \rightarrow c$, then $h \rightarrow 0$.

So, h(c)=cosc and

$$\lim_{x \to c} h(x) = \lim_{x \to c} \cos x$$

=
$$\lim_{h \to 0} \cos(c+h)$$

=
$$\lim_{h \to 0} [\cos \cosh - \sin \sinh]$$

=
$$\lim_{h \to 0} \cos \cosh - \lim_{h \to 0} \sin \cosh h$$

=
$$\cos \cos \theta - \sin \theta$$

=
$$\cos c \times 1 - \sin c \times \theta$$

=
$$\cos c$$

Therefore, $\lim_{h\to 0} h(x) = h(c)$.

Thus, the function h is continuous.

Hence, we conclude that all the following functions are continuous.

(a)
$$f(x)=g(x)+h(x)=sinx+cosx$$
.

(b)
$$f(x)=g(x)-h(x)=sinx-cosx$$
.

(c) $f(x)=g(x) \times h(x)=sinx \times cosx$.

22. Verify whether the following trigonometric functions are continuous. sine, cosine, cosecant, secant and cotangent.

Ans: We know that if two functions say g and h are continuous, then

i.
$$\frac{h(x)}{g(x)}, g(x) \neq 0$$
 is continuous.

ii.
$$\frac{1}{g(x)}$$
, $g(x) \neq 0$ is continuous.

iii.
$$\frac{1}{h(x)}$$
, $h(x) \neq 0$ is continuous.

It can be observed that the function g(x)=sinx is defined for all real numbers.

Now, let consider c be a real number and substitute x=c+h into the function g.

```
When, x \rightarrow c, then h \rightarrow 0.

So, g(c)=sinc and

\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} sinx
= \lim_{h \rightarrow 0} sin(c+h)
= \lim_{h \rightarrow 0} [sinccosh+coscsinh]
= \lim_{h \rightarrow 0} (sinccosh) + \lim_{h \rightarrow 0} (coscsinh)
= sinccos0 + coscsin0
= sinc+0
= sinc
```

```
Therefore, \lim_{x\to c} g(x) = g(c).
```

Thus, the function g(x)=sinx is continuous.

Again, let h(x) = cosx.

Then, h(c) = cosc and

It can be noted that $h(x)=\cos x$ is defined for all real numbers.

Now, let consider c be a real number and substitute x=c+h into the function h.

```
\lim_{x \to c} h(x) = \lim_{x \to c} \cos x
= \lim_{h \to 0} \cos(c+h)
= \lim_{h \to 0} [\cos \cosh - \sin \sinh]
= \lim_{h \to 0} \cos \cosh - \lim_{h \to 0} \sin \cosh h
= \cos \cosh - \sin \sin \theta
= \cos \cos \theta - \sin \theta
```

Therefore, $\lim_{h\to 0} h(x) = h(c)$.

Thus, the function $h(x)=\cos x$ is continuous.

Now note that,

cosec $x = \frac{1}{\sin x}$, and $\sin x \neq 0$ is a continuous function.

 \Rightarrow cosec x, x \neq n π (n \in Z) is also a continuous function.

Also, secant function is continuous except at $x=(2n+1)\frac{\pi}{2}$, $n \in \mathbb{Z}$.

Therefore, $\sec x = \frac{1}{\cos x}$, $\cos x \neq 0$ is continuous.

 \Rightarrow secx, $x \neq (2n+1)\frac{\pi}{2}$, $n \in \mathbb{Z}$ is a continuous function.

Thus, secant function is also continuous except at $x=(2n+1)\frac{\pi}{2}$, $n \in \mathbb{Z}$.

And the cotangent function is

 $\cot x = \frac{\cos x}{\sin x}$, and where $\sin x \neq 0$ is a continuous function.

 \Rightarrow cotx, x \neq n π , n $\in \mathbb{Z}$ is a continuous function.

Hence, the cotangent function is continuous except at $x=n\pi$, $n \in \mathbb{Z}$.

23. Determine all the discontinuity points for the following function f defined by $f(x) = \left\{ \frac{\sin x}{x}, \text{ if } x < 0 \right\}.$

by
$$f(x) = \left\{ \begin{array}{c} x \\ x \\ x+1, \text{ if } x \ge 0 \end{array} \right\}.$$

Ans: The given function is
$$f(x) = \begin{cases} \frac{\sin x}{x}, & \text{if } x < 0 \\ x+1, & \text{if } x \ge 0 \end{cases}$$
.

Note that, the function f is defined at every point on the real number line. Now, let consider c be a real number. Then there may arise three cases, either c<0, or c>0, or c=0.

Let us discuss one after another.

Case I: When c<0.

Then,
$$f(c) = \frac{\sin c}{c}$$
.
Also, $\lim_{x \to c} f(x) \left(\frac{\sin x}{x} \right) = \frac{\sin c}{c}$.
Therefore, $\lim_{x \to c} f(x) = f(c)$.

Hence, the function f is continuous at every point x, for x < 0.

Case II: When c>0.

Then f(c)=c+1.

Also,
$$\lim_{x \to c} f(x) = \lim_{x \to c} (x+1) = c+1.$$

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Hence, the function f is continuous at every point, where x>0.

Case III: When c = 0.

Then f(c)=f(0)=0+1=1.

Now, the left-hand-limit of the function f at x=0 is

 $\lim_{x \to 0^{\circ}} f(x) = \lim_{x \to 0^{\circ}} \frac{\sin x}{x} = 1$ and the right-hand-limit is

 $\lim_{x \to 0^{-}} f(x) = \lim_{x \to 0^{-}} (x+1) = 1$

Therefore, $\lim_{x\to 0^{-}} f(x) = \lim_{x\to 0^{-}} f(x) = f(0)$.

So, the function f is continuous at x = 0. Thus, the function f is continuous at every real point.

Hence, the function f does not have any point of discontinuity.

24. Discuss the continuity of the function f defined by

$$\mathbf{f}(\mathbf{x}) = \begin{cases} \mathbf{x}^2 \sin \frac{1}{\mathbf{x}}, & \text{if } \mathbf{x} \neq \mathbf{0} \\ \mathbf{0}, & \text{if } \mathbf{x} = \mathbf{0} \end{cases}.$$

Ans: The given function is
$$f(x) = \begin{cases} x^2 \sin \frac{1}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$
.

We can observe that the function f is defined at every point on the real number line.

Now, let consider c be a real number.

Then, there may arise two cases, either $c \neq 0$ or c=0.

Let us discuss the cases one after another.

Case I: When $c \neq 0$.

Then
$$f(c)=c^2\sin\frac{1}{c}$$
.

Also,

$$\lim_{x\to c} f(x) = \lim_{x\to c} \left(x^2 \sin \frac{1}{x} \right) = \left(\lim_{x\to c} x^2 \right) \left(\limsup_{x\to c} x^2 - \frac{1}{x} \right) = c^2 \sin \frac{1}{c}.$$

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Hence, the function f is continuous at every point $x \neq 0$.

Case II: When c = 0.

Then f(0)=0 and also

$$\lim_{x \to 0^{-}} f(x) = \lim_{x \to 0^{-}} \left(x^{2} \sin \frac{1}{x} \right) = \lim_{x \to 0} \left(x^{2} \sin \frac{1}{2} \right).$$

Now, we know that,

$$-1 \le \sin \frac{1}{x} \le 1, \ x \ne 0.$$

$$\Rightarrow -x^{2} \le \sin \frac{1}{x} \le x^{2}$$

$$\Rightarrow \lim_{x \to 0} (-x^{2}) \le \lim_{x \to 0} \left(x^{2} \sin \frac{1}{x} \right) \le 0$$

$$\Rightarrow 0 \le \lim_{x \to 0} \left(x^{2} \sin \frac{1}{x} \right) \le 0$$

$$\Rightarrow \lim_{x \to 0} \left(x^{2} \sin \frac{1}{x} \right) = 0$$

Therefore, $\lim_{x\to 0^-} f(x) = 0$.

Similarly, we have,

$$\lim_{x \to 0^+} f(x) = \lim_{x \to 0^+} \left(x^2 \sin \frac{1}{x} \right) = \lim_{x \to 0} \left(x^2 \sin \frac{1}{x} \right) = 0$$

Therefore, $\lim_{x \to 0^-} f(x) = f(0) = \lim_{x \to 0^+} f(x)$.

Thus, the function f is continuous at the point x = 0.

So, the function f is continuous at all real points.

Hence, the function f is continuous.

25. Determine whether the following function **f** is continuous.

f such that $f(x) = \begin{cases} sinx-cosx, if x \neq 0 \\ 1 & if x=0 \end{cases}$.

Ans: The given function is $f(x) = \begin{cases} \sin x - \cos x, & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$.

It can be observed that the function f is defined at every point on the real number line.

Now, let consider c be a real number.

Then, there may arise two cases, either $c \neq 0$ or c=0.

Let us discuss the cases one after another.

Case I: When $c \neq 0$.

Then, f(c)=sinc-cosc.

Also, $\lim_{x\to c} f(x) = \lim_{x\to c} (\sin x - \cos x) = \operatorname{sinc-cosc}$.

Therefore, $\lim_{x\to c} f(x) = f(c)$.

Hence, the function f is continuous at every point x for $x \neq 0$.

Case II: When c = 0.

Then, f(0) = -1.

Now the left-hand-limit of the function f at x=0 is

 $\lim_{x \to 0^{-}} f(x) = \lim_{x \to 0^{-}} (\sin x - \cos x) = \sin 0 - \cos 0 = 0 - 1 = -1 \text{ and the right-hand-limit is}$ $\lim_{x \to 0^{+}} f(x) = \lim_{x \to 0^{+}} (\sin x - \cos x) = \sin 0 - \cos 0 = 0 - 1 = -1.$ Therefore, $\lim_{x \to 0^{-}} f(x) = \lim_{x \to 0^{+}} f(x) = f(0).$

So, the function f is continuous at x = 0.

Thus, the function f is continuous at all real points.

Hence, the function f is continuous.

26. Calculate the values of k for which the function f attains continuity at the given points.

$$f(x) \begin{cases} \frac{k\cos x}{\pi - 2x}, & \text{if } x \neq \frac{\pi}{2} \\ 3, & \text{if } x = \frac{\pi}{2} \end{cases}$$

Ans: The given function is
$$f(x)$$
 $\begin{cases} \frac{k\cos x}{\pi - 2x}, & \text{if } x \neq \frac{\pi}{2} \\ 3, & \text{if } x = \frac{\pi}{2} \end{cases}$.

Observe that, f is defined and continuous at $x = \frac{\pi}{2}$, since the value of the f at $x = \frac{\pi}{2}$ is equal with the limiting value of f at $x = \frac{\pi}{2}$. Since f is defined at $x = \frac{\pi}{2}$ and $f\left(\frac{\pi}{2}\right) = 3$, so

Since, f is defined at $x = \frac{\pi}{2}$ and $f\left(\frac{\pi}{2}\right) = 3$, so

$$\lim_{x \to \frac{\pi}{2}} f(x) = \lim_{x \to \frac{\pi}{2}} \frac{k \cos x}{\pi - 2x}.$$

Substitute $x = \frac{\pi}{2} + h$ into the function f(x).

So, we have,
$$x \rightarrow \frac{\pi}{2} \Rightarrow h \rightarrow 0$$
.

Then,

$$\lim_{x \to \frac{\pi}{2}} f(x) = \lim_{x \to \frac{\pi}{2}} \frac{k \cos x}{\pi - 2x} = \lim_{h \to 0} \frac{k \cos \left(\frac{\pi}{2} + h\right)}{\pi - 2\left(\frac{\pi}{2} + h\right)}.$$

$$\Rightarrow k \lim_{h \to 0} \frac{-\sinh}{-2h} = \frac{k}{2} \lim_{h \to 0} \frac{\sinh}{h} = \frac{k}{2}.1 = \frac{k}{2}.$$

Therefore,
$$\lim_{x \to \frac{\pi}{2}} f(x) = f\left(\frac{\pi}{2}\right)$$

$$\Rightarrow \frac{k}{2} = 3$$

$$\Rightarrow k = 6$$

Hence, the value of k is 6 for which the function f is continuous.

27. Determine the values of k for which the following function f satisfies continuity at the given points.

$$\mathbf{f}(\mathbf{x}) = \begin{cases} \mathbf{k}\mathbf{x}^2, & \text{if } \mathbf{x} \le 2\\ 3, & \text{if } \mathbf{x} > 2 \end{cases} \text{ at } \mathbf{x} = 2.$$

Ans: The given function is $f(x) = \begin{cases} kx^2, \text{ if } x \le 2\\ 3, \text{ if } x > 2 \end{cases}$.

Note that, f is continuous at x = 2 only if f is defined at x=2 and if the value of f at x = 2 is equal with the limiting value of f at x = 2.

Since, it is provided that f is defined at x=2 and $f(2)=k(2)^2=4k$, so

$$\lim_{x \to 2^{+}} f(x) = \lim_{x \to 2^{+}} f(x) = f(2)$$

$$\Rightarrow \lim_{x \to 2^{-}} (kx^{2}) = \lim_{x \to 2^{+}} (3) = 4k$$

$$\Rightarrow k \times 2^{2} = 3 = 4k$$

$$\Rightarrow 4k = 3 = 4k$$

$$\Rightarrow 4k = 3$$

$$\Rightarrow k = \frac{3}{4}$$

Hence, the value of k is $\frac{3}{4}$ for which the function f is continuous.

28. Determine the values of k for which the following function f attains continuity at the given point.

$$\mathbf{f}(\mathbf{x}) = \begin{cases} \mathbf{k}\mathbf{x} + 1 & \text{, if } \mathbf{x} \le \pi \\ \cos \mathbf{x}, & \text{if } \mathbf{x} > \pi \end{cases} \quad \text{at } \mathbf{x} = \pi.$$

- Ans: For k(x) to be continuous at π , $k\pi + 1 = \cos \pi$ $k\pi = -1 - 1$ $k = -2/\pi$
- 29. Determine the values of k continuity at the provided point. for which the following function f attains

$$f(x) = \begin{cases} kx+1, \text{ if } x \le 5\\ 3x-5, \text{ if } x > 5 \end{cases} \text{ at } x = 5$$

Ans: The given function is $f(x) = \begin{cases} kx+1, \text{ if } x \le 5\\ 3x-5, \text{ if } x > 5 \end{cases}$.

1

Now, note that, the function f is continuous at x = 5 only if the value of f at x = 5 is equal to the limiting value of f at x = 5.

Since it is given that, the function f is defined at x = 5 and f(5)=kx+1=5k+1, so

$$\lim_{x \to 5} f(x) = \lim_{x \to 5^+} (3x-5) = 5k + 1$$
$$\Rightarrow 5k + 1 = 15 - 5 = 5k + 1$$
$$\Rightarrow 5k + 1 = 10$$
$$\Rightarrow 5k = 9$$
$$\Rightarrow k = \frac{9}{5}$$

Hence, the value of k is $\frac{9}{5}$ for which the function f is continuous at x=5.

30. Determine the values of constants a and b for which the following function f is continuous.

$$\mathbf{f} \text{ such that } \mathbf{f}(\mathbf{x}) = \begin{cases} \mathbf{5}, & \text{if } \mathbf{x} \le 2\\ \mathbf{ax+b}, & \text{if } 2 < \mathbf{x} < 10\\ \mathbf{21}, & \text{if } \mathbf{x} \ge 10 \end{cases}.$$

$$\mathbf{Ans:} \text{ The given function is } \mathbf{f}(\mathbf{x}) = \begin{cases} \mathbf{5}, & \text{if } \mathbf{x} \le 2\\ \mathbf{ax+b}, & \text{if } 2 < \mathbf{x} < 10\\ \mathbf{21}, & \text{if } \mathbf{x} \ge 10 \end{cases}.$$

Note that, f is defined at every point on the real number line.

Now, realise that if the function f is continuous then f is continuous at every real number.

So, let f satisfies continuity at x=2 and x=10.

Then, since f is continuous at x=2, so

$$\lim_{x \to 2^{-}} f(x) = \lim_{x \to 2^{+}} f(x) = f(2)$$
$$\Rightarrow \lim_{x \to 2^{-}} (5) = \lim_{x \to 2^{+}} (ax+b) = 5$$

 $\Rightarrow 5=2a+b=5$ $\Rightarrow 2a+b=5 \qquad \dots \dots (1)$

Again, since f attains continuity at x=10, so

 $\lim_{x \to 10^{\circ}} f(x) = \lim_{x \to 10^{+}} f(x) = f(10)$ $\Rightarrow \lim_{x \to 10^{\circ}} (ax+b) = \lim_{x \to 10^{+}} (21) = 21$ $\Rightarrow 10a+b-21 = 21$ $\Rightarrow 10a+b=21$ (2)

Subtracting the equation (1) from the equation (2), gives

```
8a=16 \Rightarrow a=2
```

Substituting a=2 in the equation (1), gives

$$2 \times 2 + b = 5$$
$$\Rightarrow 4 + b = 5 \Rightarrow b = 1$$

Hence, the values of a and b are 2 and 1 respectively for which f is a continuous function.

31. Prove that the following function is continuous.

 $f(x)=cos(x^2)$

Ans: The given function is $f(x)=cos(x^2)$.

Note that, f is defined for all real numbers and so f can be expressed as the composition of two functions as, $f=g \circ h$, where $g(x)=\cos x$ and $h(x)=x^2$.

 $[\therefore(goh)(x)=g(h(x))=g(x^2)=cos(x^2)=f(x)]$

Now, it is to be Proven that, the functions $g(x)=\cos x$ and $h(x)=x^2$ are continuous.

Since the function g is defined for all the real numbers, so let consider c be a real number.

Then, g(c)=cosc.

Substitute x=c+h into the function g.

When, $x \rightarrow c$, then $h \rightarrow 0$.

Then we have,

$$\lim_{x \to c} g(x) = \lim_{x \to c} \cos x$$

=
$$\lim_{h \to 0} \cos(c+h)$$

=
$$\lim_{h \to 0} [\cos \cosh - \sin \cosh h]$$

=
$$\lim_{h \to 0} \cos \cosh - \lim_{h \to 0} \sin \cosh h$$

=
$$\cos \cosh - \sin \cosh \theta$$

=
$$\cos \cos \theta - \sin \theta$$

Therefore,
$$\lim_{x\to c} g(x) = g(c)$$
.

Hence, the function $g(x)=\cos x$ is continuous.

Again, $h(x)=x^2$ is defined for every real point.

So, let consider k be a real number, then $h(k)=k^2$ and

```
\lim_{x\to k} h(x) = \lim_{x\to k} x^2 = k^2.
```

Therefore, $\lim_{x\to k} h(x) = h(k)$.

Hence, the function h is continuous.

Now, remember that for real valued functions g and h, such that $(g \circ h)$ is defined at c, if g is continuous at c and f is continuous at g(c), then $(f \circ h)$ is continuous at c.

Hence, the function $f(x)=(g \circ h)(x)=\cos(x^3)$ is continuous.

32. Prove that the following function is continuous.

 $f(x) = |\cos x|$

Ans: The given function is $f(x) = |\cos x|$.

Note that, the function f is defined for all real numbers. So, the function f can be expressed as the composition of two functions as, $f=g \circ h$, where g(x)=|x| and h(x)=cosx.

[::(goh)(x)=g(h(x))=g(cosx)=|cosx|=f(x)]

Now, it is to be proved that the functions g(x)=|x| and $h(x)=\cos x$ are continuous.

Remember that, g(x) = |x|, can be written as

$$g(x) = \begin{cases} -x, \text{ if } x < 0 \\ x, \text{ if } x \ge 0 \end{cases}.$$

Now, since the function g is defined for every real number, so let consider c be a real number.

Then there may arise three cases, either c<0, or c>0, or c=0.

Let discuss the cases one after another.

```
Case I: When c<0.
```

Then, g(c) = -c.

Also, $\lim_{x\to c} g(x) = \lim_{x\to c} (-x) = -c$.

```
Therefore, \lim_{x\to c} g(x) = g(c).
```

Hence, the function g is continuous at every point x, for x < 0.

Case II: When c>0.

Then, g(c)=c.

Also, $\lim_{x\to c} g(x) = \lim_{x\to c} x = c$.

Therefore, $\lim_{x\to c} g(x) = g(c)$.

Hence, the function g is continuous at every point x for x>0.

Case III: When c=0.

Then, g(c)=g(0)=0.

Now, the left-hand-limit of the function g at x=0 is

 $\lim_{x \to 0^{-}} g(x) = \lim_{x \to 0^{+}} (-x) = 0 \text{ and the right-hand-limit is}$ $\lim_{x \to 0^{+}} g(x) = \lim_{x \to 0^{+}} (x) = 0.$

Therefore, $\lim_{x\to 0^-} g(x) = \lim_{x\to 0^+} g(x) = g(0)$.

Hence, the function g is continuous at x=0.

By observing the above three discussions, we can conclude that the function g is continuous at every real points.

Now, since the function $h(x)=\cos x$ is defined for all real numbers, so let consider c be a real number. Then, substitute x=c+h into the function h.

So, when $x \rightarrow c$, then $h \rightarrow 0$.

Then, we have

$$h(c) = cosc \text{ and}$$

$$\lim_{x \to c} h(x) = \lim_{x \to c} cosx$$

$$= \lim_{h \to 0} cos(c+h)$$

$$= \lim_{h \to 0} [cosccosh-sincsinh]$$

$$= \lim_{h \to 0} cosccosh-\lim_{h \to 0} sincsinh$$

$$= cosccos0-sincsin0$$

$$= cosc \times 1-sinc \times 0$$

$$= cosc$$

Therefore, $\lim_{x\to c} h(x) = h(c)$.

Hence, the function $h(x) = \cos x$ is continuous.

Now remember that, for real valued functions g and h, such that (g h) is defined at x=c only if g is continuous at c and f is continuous at g(c), then the composition functions $(f \circ g)$ is continuous at x=c.

Thus, the function f(x)=(goh)(x)=g(h(x))=g(cosx)=|cosx| is continuous.

33. Examine that sin | x | is a continuous function.

Ans: First suppose that, $f(x) = \sin|x|$.

Now, note that the function f is defined for all real numbers and so f can be expressed as the composition of functions as, $f=g \circ h$, where g(x)=|x| and h(x)=sinx.

$$\left[(g \circ h)(x) = g(h(x)) = g(\sin x) = |\sin x| = f(x)\right]$$

So, it is to be proved that the functions g(x)=|x| and $h(x)=\sin x$ are continuous. Now, remember that, the function g(x)=|x| can be written as

$$g(x) \begin{cases} -x, \text{ if } x < 0 \\ x, \text{ if } x \ge 0 \end{cases}.$$

Note that, the function g is defined for every real number, and so let consider c be a real number.

Then, there may arise three cases, either c<0, or c>0, or c=0.

Let us discuss the cases one after another.

Case I: When c<0. Then g(c)=-c. Also, $\lim_{x \to c} (-x) = \lim_{x \to c} x = -c$. Therefore, $\lim_{x \to c} g(x) = g(c)$.

Hence, the function g is continuous at every point x for x < 0.

Case II: When c > 0.

Then, g(c)=c

Also, $\lim_{x \to c} (-x) = \lim_{x \to c} x = c$.

Therefore, $\lim_{x \to c} g(x) = g(c)$.

Thus, the function g is continuous at every point x for x>0.

Case III: When c = 0.

Then, g(c)=g(0)=0.

Also, the left-hand-limit of the function g at x=0 is

 $\lim_{x \to 0^{-}} g(x) = \lim_{x \to 0^{-}} (-x) = 0 \text{ and the right-hand -limit is}$ $\lim_{x \to 0^{+}} g(x) = \lim_{x \to 0^{+}} (x) = 0.$ Therefore, $\lim_{x \to 0^{-}} g(x) = \lim_{x \to 0^{+}} (x) = g(0).$

Thus, the function g is continuous at x = 0.

By observing the above three discussions, we can conclude that that the function g is continuous at every points.

Again, since the function h(x)=sinx is defined for all real numbers, so let consider c be a real number and substitute x=c+k into the function.

Now, when x $\uparrow c$ then k $\uparrow 0$.

Then, we have

h(c)=sinc.

Also,

$$\lim_{x \to c} h(x) = \lim_{x \to c} \sin x$$

=
$$\lim_{k \to 0} \sin(c+k)$$

=
$$\lim_{k \to 0} [\operatorname{sinccosk} + \operatorname{coscsink}]$$

=
$$\lim_{k \to 0} (\operatorname{sinccosk}) + \lim_{h \to 0} (\operatorname{coscsink})$$

=
$$\operatorname{sinccos0} + \operatorname{coscsin0}$$

=
$$\operatorname{sinc} + 0$$

=
$$\operatorname{sinc}$$

Therefore, $\lim_{x \to c} h(x) = g(c)$.

Hence, the function h is continuous.

Now, remember that, for any two real valued functions g and h, such that the cmposition of functions $g \circ h$ is defined at c, if g is continuous at c and f is continuous at g(c), then the composition function $f \circ h$ is continuous at c.

Thus, the function f(x)=(goh)(x)=g(h(x))=g(sinx)=|sinx| is a continuous.

- 34. Determine all the discontinuity points of the following function f defined by f (x) = | x | | x + 1 |.
- **Ans:** The given function is $f(x) = |x| |\sin x|$.

Let consider two functions

g(x) = |x| and h(x) = |x+1|.

Then we get, f=g-h.

Now, the function g(x) = |x| can be written as

$$g(x) = \begin{cases} -x, \text{ if } x < 0 \\ x, \text{ if } x \ge 0 \end{cases}.$$

Note that, the function g is defined for every real number and so let consider c be a real number.

Then there may arise three cases, either c<0, or c>0, or c=0.

Let us discuss the cases one after another.

Case I: When c<0.

Then, g(c)=g(0)=-c.

Also, $\lim_{x \to c} g(x) = \lim_{x \to c} (-x) = -c$.

Therefore, $\lim_{x \to c} g(x) = g(c)$.

Hence, the function g is continuous at every point x for x < 0.

Case II: When c>0.

Then g(c)=c.

Also, $\lim_{x \to c} g(x) = \lim_{x \to c} x = c$.

Therefore, $\lim_{x \to c} g(x) = g(c)$.

Hence, the function g is continuous at every point x, where x>0.

Case III: When c = 0.

Then g(c)=g(0)=0.

Also, the left-hand-limit of the function g at x=0 is

 $\lim_{x \to 0^{-}} g(x) = \lim_{x \to 0^{-}} (-x) = 0 \text{ and the right-hand-limit is}$ $\lim_{x \to 0^{+}} g(x) = \lim_{x \to 0^{+}} (x) = 0.$ Therefore, $\lim_{x \to 0^{-}} g(x) = \lim_{x \to 0^{+}} (x) = g(0).$

Hence, the function g is continuous at x = 0.

Thus, we can conclude by observing the above three discussions that g is continuous at every real point.

Now, remember that, the function h(x)=|x+1| can be written as

$$h(x) = \begin{cases} -x(x+1), \text{ if, } x < -1 \\ x+1, \text{ if, } x \ge -1 \end{cases}.$$

Note that, the function h is defined for all real numbers, and so let consider c be a real number.

Case I: When c<-1.

Then h(c) = -(c+1).

```
Also, \lim_{x \to c} [-(x+1)] = -(c+1).
```

Therefore, $\lim_{x \to c} h(x) = h(c)$.

Hence, the function h attains continuity at every real point x, where x < -1.

Case II: When c>-1.

Then, h(c)=c+1.

Also, $\lim_{x \to c} h(x) = \lim_{x \to c} (x+1) = (c+1)$.

Therefore, $\lim_{x \to c} h(x) = h(c)$.

Hence, the function h satisfies continuity at every real point x for x>-1.

Case III: When c = -1.

Then, h(c)=h(-1)=-1+1=0.

Also, the left-hand-limit of the function h at x=1 is

 $\lim_{x \to 1^{-}} h(x) = \lim_{x \to 1^{-}} [-(x+1)] = -(-1+1) = 0 \text{ and the right-hand-limit is}$ $\lim_{x \to 1^{+}} h(x) = \lim_{x \to 1^{+}} (x+1) = (-1+1) = 0.$

Therefore, $\lim_{x \to 1^-} h = \lim_{x \to 1^+} h(x) = h(-1)$.

Thus, the function h satisfies continuity at x=-1.

Hence, by observing the above three discussions, we can conclude that the function h is continuous for every real point.

Now, since the functions g and h are both continuous, so the function f=g-h is also continuous.

Hence, the function f does not have any discontinuity points.

Exercise 5.2

1. Compute the derivative of the function $f(x) = \sin(x^2+5)$ with respect to x.

Ans: Let $f(x)=sin(x^2+5)$, $u(x)=x^2+5$, and v(t)=sint

Then, $(v \circ u)(x) = v(u(x)) = v(x^2+5) = tan(x^2+5) = f(x)$

Therefore, f is a composition of two functions u and v.

Substitute $t=u(x)=x^2+5$.

Then, it gives

$$\frac{dv}{dt} = \frac{d}{dt}(\sin t) = \cos t = \cos(x^2 + 5)$$
$$\frac{dt}{dx} = \frac{d}{dx}(x^2 + 5) = \frac{d}{dx}(x^2) + \frac{d}{dx}(5) = 2x + 0 = 2x$$

Applying the chain rule of derivatives gives

 $\frac{df}{dx} = \frac{dv}{dt} \cdot \frac{dt}{dx} = \cos(x^2 + 5) \times 2x = 2x\cos(x^2 + 5)$

An alternate method:

 $\frac{df}{dx} = \frac{dv}{dt} \cdot \frac{dt}{dx} = \cos(x^2 + 5) \cdot \frac{d}{dx} (x^2 + 5)$

$$=\cos(x^{2}+5)\cdot\left[\frac{d}{dx}(x^{2})+\frac{d}{dx}(5)\right]$$
$$=\cos(x^{2}+5)\cdot[2x+0]$$
$$=2x\cos(x^{2}+5)$$

Hence, the derivative of the function $f(x) = sin(x^2 + 5)$ is $2xcos(x^2+5)$.

2. Compute the derivative of the function $f(x) = \cos(\sin x)$ with respect to x.

Ans: Let suppose that, $f(x)=\cos(\sin x)$, $u(x)=\sin x$, and $v(t)=\cos t$

Then, $(v \circ u)(x)=v(u(x))=v(sinx)=cos(sinx)=f(x)$

Therefore, it is observed that f is the composition of two functions u and v.

Now, substitute t=u(x)=sinx.

Then,

$$\frac{dv}{dt} = \frac{d}{dt}(\cos t) = -\sin(\sin x)$$
 and
dt d

$$\frac{dt}{dx} = \frac{d}{dx}(\sin x) = \cos x$$
.

Applying the chain rule of derivatives gives

 $\frac{df}{dx} = \frac{dv}{dt} \cdot \frac{dt}{dx} = \sin(\sin x) \cdot \cos x = -\cos x \sin(\sin x)$

An alternate method:

$$\frac{d}{dx} [\cos(\sin x)] = -\sin(\sin x) \cdot \frac{d}{dx} (\sin x) = -\sin(\sin x) \cdot \cos x = -\cos x \sin(\sin x) \cdot \cos x = -\cos x \sin x - \cos x \sin x - \cos x \sin x - \cos x = -\cos x \sin x - \cos x \sin x - \cos x = -\cos x \sin x - \cos x \sin x - \cos x = -\cos x \sin x - \cos x \sin x - \cos x = -\cos x \sin x - \cos x \sin x - \cos x = -\cos x \sin x - \cos x - \cos x \sin x - \cos x \sin x - \cos x \sin x - \cos x - \cos x - \cos x \sin x - \cos x -$$

Hence, the derivative of the function f(x) = cos(sinx) is -cosxsin(sinx).

3. Compute the derivative of the function f(x) = sin(ax+b) with respect to x

Ans: Let suppose that, f(x)=sin(ax+b), u(x)=ax+b, and v(t)=sint

Then we get, $(v \circ u)(x)=v(u(x))=v(ax+b)=sin(ax+b)=f(x)$.

It is observed that the function $f\,$ is the composition of two functions $\,u\,$ and $\,v\,$.

Now, substitute t=u(x)=ax+b.

Therefore,

$$\frac{dv}{dt} = \frac{d}{dt}(sint) = cost = cos(ax+b)$$
 and

$$\frac{dt}{dx} = \frac{d}{dx}(ax+b) = \frac{dt}{dx}(ax) + \frac{d}{dx}(b) = a + 0 = a$$

Applying the chain rule deriavtives, gives

 $\frac{df}{dx} = \frac{dv}{dt} \cdot \frac{dt}{dx} = \cos(ax+b) \cdot a = a\cos(ax+b) \cdot a$

Alternate method

$$\frac{d}{dx} [\sin(ax+b)] = \cos(ax+b) \cdot \frac{d}{dx} (ax+b)$$
$$= \cos(ax+b) \times \left[\frac{d}{dx} (ax) + \frac{d}{dt} (b) \right]$$
$$= \cos(ax+b) \times (a+0)$$
$$= a\cos(ax+b)$$

Hence, the derivative of the function f(x) = sin(ax+b) is acos(ax+b).

4. Compute the derivative of the function $f(x) = \sec(\tan(\sqrt{x}))$ with resepcet to x.

Ans: Let suppose that, $f(x)=\sec(\tan(\sqrt{x}))$, $u(x)=\sqrt{x}$, $v(t)=\tan t$, and $w(s)=\sec s$

Then, we get, $(w \circ v \circ u)(x) = w[v(u(x))] = w[v(\sqrt{x})] = w(\tan\sqrt{x}) = f(x)$.

It is observed that the function $g\,$ is the composition of three functions $\,u\,$, $\,v\,$ and $\,w\,.$

Now, substitute s=v(t) and t=u(x)= \sqrt{x} .

Then, we get

$$\frac{dw}{ds} = \frac{d}{ds} (secs) = secs = sec(tant) \times tan(tant) \qquad [s=tant]$$
$$= sec(tan\sqrt{x}) \times tan(tan\sqrt{x}) \qquad [t=\sqrt{x}]$$

Thus, applying the chain rule of derivatives gives

$$\frac{dt}{dx} = \frac{dw}{ds} \cdot \frac{ds}{dt} \times \frac{dt}{dx}$$

$$= \sec(\tan(\sqrt{x}) \times (\tan(\sqrt{x}) \times \sec^{2}\sqrt{x} \times \frac{1}{2\sqrt{x}})$$

$$= \frac{1}{2\sqrt{x}} \sec^{2}\sqrt{x} (\tan\sqrt{x}) \tan(\tan\sqrt{x})$$

$$= \frac{\sec^{2}\sqrt{x} \sec(\tan\sqrt{x}) \tan(\tan\sqrt{x})}{2\sqrt{2}}$$

An alternate method:

$$\frac{d}{dx} \left[\sec(\tan(\sqrt{x})) \right] = \sec(\tan(\sqrt{x}).(\tan(\sqrt{x})).\frac{d}{dx}(\tan(\sqrt{x})))$$
$$= \sec(\tan(\sqrt{x})) \times \tan(\tan(\sqrt{x})) \times \sec^{2}(\sqrt{x})) \times \frac{d}{dx}(\sqrt{x})$$
$$= \sec(\tan(\sqrt{x})) \times \tan(\tan(\sqrt{x})) \times \sec^{2}(\sqrt{x})) \times \frac{1}{2\sqrt{x}}$$
$$= \frac{\sec(\tan(\sqrt{x})) \times \tan(\tan(\sqrt{x})) \times \sec^{2}(\sqrt{x}))}{2\sqrt{x}}$$

Hence, the derivative of the function $f(x) = \sec(\tan(\sqrt{x}))$ is

$$\frac{\sec^2\sqrt{x}\sec(\tan\sqrt{x})\tan(\tan\sqrt{x})}{2\sqrt{2}}$$

5. Compute the derivative of the function $f(x) = \frac{\sin(ax+b)}{\cos(cx+d)}$ with respect to x.

Ans: The given function is $f(x) = \frac{\sin(ax+b)}{\cos(cx+d)}$.

Now, let g(x)=sin(ax+b) and h(x)=cos(cx+d).

Here we will use the divide formula of derivatives $f' = \frac{g'h-gh'}{h^2}$ (1)

First, consider the function g(x)=sin(ax+b).

Let assume u(x)=ax+b, and v(t)=sint.

Then, we get $(v \circ u)(x)=v(u(x))=v(ax+b)=sin(ax+b)=g(x)$.

Therefore, we observe that the function g is the composition of two functions, u and v.

So, substitute t=u(x)=ax+b.

Then,

 $\frac{dv}{dt} = \frac{d}{dt}(sint) = cost = cos(ax+b)$ and

$$\frac{dt}{dx} = \frac{d}{dx}(ax+b) = \frac{dt}{dx}(ax) + \frac{d}{dx}(b) = a + 0 = a$$

Therefore, applying the chain rule of derivatives gives

$$g' = \frac{dg}{dx} = \frac{dv}{dt} \cdot \frac{dt}{dx} = \cos(ax+b) \cdot a = a\cos(ax+b) \cdot a$$

Now, consider the function h(x)=cos(cx+d).

Let suppose p(x)=cx+d, and q(t)=cosy.

Then, we have $(q \circ p)(x)=q(p(x))=q(cx+d)=cos(cx+d)=h(x)$.

Therefore, the function h is the composition of two functions p and q.

Now, substitute y=p(x)=cx+d.

Then we have,

$$\frac{dq}{dy} = \frac{d}{dy}(\cos y) = -\sin y = -\sin(cx+d)$$
 and

$$\frac{dy}{dx} = \frac{d}{dx}(cx+d) = \frac{d}{dx}(cx) + \frac{d}{dx}(d) = c.$$

Therefore, applying the chain rule of derivatives gives

$$h' = \frac{dh}{dx} = \frac{dq}{dy} \cdot \frac{dy}{dx} = -\sin(cx+d) \times c = -c\sin(cx+d)$$
.

Now, substituting all the obtained derivatives into the formula (1) gives

$$f' = \frac{a\cos(ax+b) \times \cos(cx+d) - \sin(ax+b)(-c\sin(cx+d))}{[\cos(cx+d)]^2}$$
$$= \frac{a\cos(ax+b)}{\cos(cx+d)} + c\sin(ax+b) \times \frac{\sin(cx+d)}{\cos(cx+d)} \times \frac{1}{\cos(cx+d)}$$
$$= a\cos(ax+b)\sec(cx+d) + c\sin(ax+b)\tan(cx+d)\sec(cx+d)$$
Hence, the derivative of the function $f(x) = \frac{\sin(ax+b)}{\cos(cx+d)}$ is

acos(ax+b)sec(cx+d)+csin(ax+b)tan(cx+d)sec(cx+d).

6. Compute the derivative of the function $f(x) = cos(x^3) \times sin^2(x^5)$ with resepct to x.

Ans: The given function is $f(x) = \cos(x^3) \times \sin^2(x^5)$.

Then,

$$\frac{d}{dx}[\cos x^{3} \times \sin^{2}(x^{5})] = \sin^{2}(x^{5}) \times \frac{d}{dx}(\cos x^{3}) + \cos x^{3}x \frac{d}{dx}[\sin^{2}(x^{5})]$$

$$= \sin^{2}(x^{5}) \times (-\sin x^{3}) \times \frac{d}{dx}(x^{3}) + \cos x^{3} + 2\sin(x^{5}) \times \frac{d}{dx}[\sin x^{5}]$$

$$= \sin^{2}(x^{5}) \times (-\sin x^{3}) \times \frac{d}{dx}(x^{3}) + \cos x^{3} + 2\sin(x^{5}) \times \frac{d}{dx}[\sin x^{5}]$$

$$= \sin^{2}(x^{5}) \times (-\sin x^{3}) \times \frac{d}{dx}(x^{3}) + \cos x^{3} + 2\sin(x^{5}) \times \frac{d}{dx}[\sin x^{5}]$$

$$= \sin^{2}(x^{5}) \times (-\sin x^{3}) \times \frac{d}{dx}(x^{3}) + \cos x^{3} + 2\sin(x^{5}) \times \frac{d}{dx}[\sin x^{5}]$$

$$= \sin^{2}(x^{5}) \times (-\sin x^{3}) \times \frac{d}{dx}(x^{3}) + \cos x^{3} + 2\sin(x^{5}) \times \frac{d}{dx}[\sin x^{5}]$$

$$= \sin^{3}(x^{5}) \times (-\sin x^{3}) \times \frac{d}{dx}(x^{3}) + \cos x^{3} + 2\sin(x^{5}) \times \frac{d}{dx}[\sin x^{5}]$$

$$= \sin^{3}(x^{5}) \times (-\sin x^{3}) \times \frac{d}{dx}(x^{3}) + \cos x^{3} + 2\sin(x^{5}) \times \frac{d}{dx}[\sin x^{5}]$$

$$= \sin^{3}(x^{5}) \times (3x^{2} + 2\sin x^{5}\cos x^{3} \times \cos x^{5} \times \frac{d}{dx}(x^{5})$$

$$= 3x^{2}\sin x^{3} \times \sin^{3}(x^{5}) + 2\sin x^{5}\cos x^{3} \times 5x^{4}$$

$$= 10x^{4}\sin x^{5}\cos x^{5}\cos x^{3} - 3x^{2}\sin x^{3}\sin^{2}(x^{5})$$

Hence, the derivative of the function $f(x) = \cos(x^3) \times \sin^2(x^5)$ is $10x^4 \sin x^5 \cos x^5 - 3x^2 \sin x^3 \sin^2(x^5)$.

7. Compute the derivative of the function $f(x) = \sqrt[2]{\cot(x^2)}$ with respect to x

Ans: The given function is $f(x) = \sqrt[2]{\cot(x^2)}$.

Then,

$$\frac{d}{dx} \left[\sqrt[2]{\cot(x^2)} \right]$$
$$= 2 \times \frac{1}{\sqrt[2]{\cot(x^2)}} \times \frac{d}{dx} \left[\cot(x^2) \right]$$
$$= \sqrt{\frac{\sin(x^2)}{\cot(x^2)}} \times \csc^2(x^2) \times \frac{d}{dx}(x^2)$$

$$= \sqrt{\frac{\sin(x^2)}{\cot(x^2)}} \times \frac{1}{\sin^2(x^2)} \times (2x)$$
$$= \frac{-2\sqrt{2x}}{\sqrt{\cos x^2}\sqrt{\sin x^2 \sin x^2}}$$
$$= \frac{-2\sqrt{2x}}{\sqrt{2\sin x^2 \cos x^2} \sin x^2}$$
$$= \frac{-2\sqrt{2x}}{\sin x^2\sqrt{\sin 2x^2}}$$

Hence, the derivative of the function $f(x) = \sqrt[2]{\cot(x^2)}$ is $\frac{-2\sqrt{2x}}{\sin^2\sqrt{\sin 2x^2}}$.

8. Compute the derivative of the function $f(x) = \cos(\sqrt{x})$ with respect to x.

Ans: The given function is $f(x) = \cos(\sqrt{x})$

Now, let $u(x) = \sqrt{x}$ and v(t) = cost.

Then, we have, $(v \circ u)(x)=v(u(x))=v(\sqrt{x})=\cos\sqrt{x}=f(x)$.

It is observed that the function f is the composition of two functions u and v. So, let $t=u(x)=\sqrt{x}$.

Then,

$$\frac{dt}{dx} = \frac{d}{dx}(\sqrt{x}) = \frac{d}{dx}\left(x^{\frac{1}{2}}\right) = \frac{1}{2}x^{\frac{1}{2}} = \frac{1}{2\sqrt{x}}.$$

Also,

$$\frac{\mathrm{d}v}{\mathrm{d}t} = \frac{\mathrm{d}}{\mathrm{d}t}(\cos t) = -\sin t = \sin(\sqrt{x}).$$

Now, by applying the chain rule of derivatives, gives

$$\frac{dt}{dx} = \frac{dv}{dt} \times \frac{dt}{dx}$$
$$= -\sin(\sqrt{x}) \times \frac{1}{2\sqrt{x}}$$
$$= -\frac{1}{2\sqrt{x}}\sin(\sqrt{x})$$
$$= -\frac{\sin(\sqrt{x})}{2\sqrt{x}}$$

An alternate method:

$$\frac{d}{dx} \left[\cos(\sqrt{x}) \right]$$

=-sin(\sqrt{x}). $\frac{d}{dx}(\sqrt{x})$
=-sin(\sqrt{x})× $\frac{d}{dx} \left(x^{\frac{1}{2}} \right)$
=-sin \sqrt{x} × $\frac{1}{2}x^{\frac{1}{2}}$
= $\frac{-\sin\sqrt{x}}{2\sqrt{x}}$

Hence, the derivative of the function $f(x) = \cos(\sqrt{x})$ is $-\frac{\sin(\sqrt{x})}{2\sqrt{x}}$.

9. Prove that the function $f(x)=|x-1|, x \in \mathbb{R}$ is not differentiable at x=1.

Ans: The given function is $f(x)=|x-1|, x \in \mathbb{R}$.

We know that a function f is called differentiable at a point x=c in its domain if both the $\lim_{k \to 0^{-}} \frac{f(c+h)-f(c)}{h}$ and $\lim_{h \to 0^{+}} \frac{f(c+h)-f(c)}{h}$ are finite and equal.

Now verify the differentiability for the function f at the point x = 1.

First, the left-hand-derivative is

$$\lim_{h \to 0^{-}} \frac{f(1+h)-f(1)}{h} = \lim_{h \to 0^{-}} \frac{f\left|1+h-1\right|\left|1-1\right|}{h}$$
$$\lim_{h \to 0^{-}} \frac{f\left|h\right|-0}{h} = \lim_{h \to 0^{+}} \frac{-h}{h} = 1, \text{ since } h < 0 \Longrightarrow |h| = -h.$$

Now the right-hand-derivative is

$$\lim_{h \to 0^{+}} \frac{f(1+h)-f(1)}{h} = \lim_{h \to 0^{+}} \frac{f\left|1+h-1\right|\left|1-1\right|}{h}$$
$$\lim_{h \to 0^{+}} \frac{f\left|h\right|-0}{h} = \lim_{h \to 0^{+}} \frac{-h}{h} = -1, \text{ since } h > 0 \Longrightarrow |h| = h.$$

From the above, it is noted that
$$\lim_{h \to 0^{-}} \frac{f(1+h)-f(1)}{h} \neq \lim_{h \to 0^{+}} \frac{f(1+h)-f(1)}{h}.$$

Hence, the function $f(x)=|x-1|, x \in R$ is not differentiable at the point x=1.

10. Prove that f(x)=[x], 0<x<3, the greatest integer function is not differentiable at the points x = 1 and x = 2.

Ans: The given function is f(x)=[x], 0 < x < 3.

Remember that a function f is called differentiable at a point x=c in its domain if both the limits, $\lim_{h \to 0^-} \frac{f(c+h)-f(c)}{h}$ and $\lim_{h \to 0^+} \frac{f(c+h)-f(c)}{h}$ are finite and equal.

First, take the left-hand-derivative of the function f at x=1 such that

$$\lim_{h \to 0^{\circ}} \frac{f(1+h)-f(1)}{h} = \lim_{h \to 0^{\circ}} \frac{[1+h]-[1]}{h} = \lim_{h \to 0^{\circ}} \frac{(0-1)}{h} = \lim_{h \to 0^{\circ}} \frac{-h}{h} = \infty.$$

Now, take the right-hand-derivative of the function f at x=1 such that

$$\lim_{h \to 0^+} \frac{f(1+h)-f(1)}{h} = \lim_{h \to 0^+} \frac{[1+h][1]}{h} = \lim_{h \to 0^+} \frac{1-1}{h} = \lim_{h \to 0^+} 0 = 0.$$

Therefore, it is being noticed that, $\lim_{h \to 0^{-}} \frac{f(1+h)-f(1)}{h} \neq \lim_{h \to 0^{+}} \frac{f(1+h)-f(1)}{h}.$ Thus, the function f is not differentiable at x=1. Now, justify the differentiability of the function f at x=2. First, take the left-hand-derivative of the function f at x=2, such that

$$\lim_{h \to 0^{-}} \frac{f(2+h)-f(2)}{h} = \lim_{h \to 0^{-}} \frac{[2+h]-[2]}{h} = \lim_{h \to 0^{-}} \frac{(1-2)}{h} = \lim_{h \to 0^{+}} \frac{-1}{h} = \infty$$

Now, take the right-hand-derivative of the function f at x=2, such that

$$\lim_{h \to 0^+} \frac{f(2+h)-f(2)}{h} = \lim_{h \to 0^+} \frac{[2+h][2]}{h} = \lim_{h \to 0^+} \frac{1-2}{h} = \lim_{h \to 0^+} 0 = 0$$

It is observed from the above discussion that, $\lim_{h \to 0^{-}} \frac{f(2+h)-f(2)}{h} \neq \lim_{h \to 0^{+}} \frac{f(2+h)-f(2)}{h}.$

Thus, the function f is not differentiable at the point x=2.

Exercise 5.3

1. Determine $\frac{dy}{dx}$ from equation 2x+3y=sinx.

Ans: The given equation is 2x+3y=sinx.

$$\frac{d}{dy}(2x+3y) = \frac{d}{dx}(\sin x)$$

$$\Rightarrow \frac{d}{dx}(2x) + \frac{d}{dx}(3y) = \cos x, \text{ applying the addition rule of derivatives}$$

$$\Rightarrow 2+3\frac{dy}{dx} = \cos x$$

$$\Rightarrow 3\frac{dy}{dx} = \cos x-2$$

Therefore,
$$\frac{dy}{dx} = \frac{\cos x - 2}{3}$$
.

2. Determine $\frac{dy}{dx}$ from the equation 2x+3y=siny.

Ans: The given equation is 2x+3y=siny.

Differentiating both sides of the equation with respect to x, gives

$$\frac{d}{dx}(2x) + \frac{d}{dx}(3y) = \frac{d}{dx}(\sin y)$$

$$\Rightarrow 2 + 3\frac{dy}{dx} = \cos y \frac{dy}{dx} , \text{ applying the chain rule of derivatives}$$

$$\Rightarrow 2 = (\cos y - 3)\frac{dy}{dx}$$

Therefore,
$$\frac{dy}{dx} = \frac{2}{\cos y - 3}$$
.

3. Determine $\frac{dy}{dx}$ from the equation $ax+by^2=cosy$.

Ans: The given function is $ax+by^2=cosy$.

Differentiating both sides of the equation with respect to x, gives $\frac{d}{dx}(ax) + \frac{d}{dx}(by^2) = \frac{d}{dx}(cosy)$ $\Rightarrow a \cdot 1 + b\frac{d}{dy}(y^2)\frac{dy}{dx} = \frac{d}{dy}(cosy)\frac{dy}{dx}$, applying the chain rule of derivatives. $\Rightarrow a + b \times 2y\frac{dy}{dx} = -\sin y\frac{dy}{dx}$

$$\Rightarrow (2by+siny) \frac{d}{dx} = a$$

Therefore, $\frac{dy}{dx} = \frac{-a}{dy^2by+siny}$.

4. Determine $\frac{dy}{dx}$ from the equation $xy+y^2=tanx+y$.

Ans: The given equation is $xy+y^2=tanx+y$.

Differentiating both sides of the equation with respect to x, gives

$$\frac{d}{dx}(xy+y^2) = \frac{d}{dx}(\tan x+y)$$

$$\Rightarrow \frac{d}{dx}(xy) + \frac{dy}{dx}(y^2) = \frac{d}{dx}(\tan y) + \frac{d}{dx}$$

$$\Rightarrow \left[y \times \frac{d}{dx}(x) + x \times \frac{dy}{dx} \right] + 2y \frac{d}{dx} = \sec^2 x + \frac{dy}{dx}, \text{ applying chain rule of derivatives.}$$

$$\Rightarrow y \times 1 + x \frac{dy}{dx} + 2y \frac{dy}{dx} = \sec^2 x + \frac{dy}{dx}$$
Therefore, $\frac{dy}{dx} = \frac{\sec^2 x - y}{(x+2y-1)}.$

5. Determine $\frac{dy}{dx}$ from the equation $x^2+xy+y^2=100$.

Ans: The given equation is $x^2+xy+y^2=100$.

Differentiating both sides of the equation with respect to x, gives $\frac{dy}{dx}(x^2+xy+y^2) = \frac{d}{dx}100$ $\Rightarrow \frac{dy}{dx}(x^2) + \frac{dy}{dx}(xy) + \frac{dy}{dx}(y^2) = 0$

$$\Rightarrow 2x + \left[y \times \frac{d}{dx}(x) + x \times \frac{dy}{dx} \right] + 2y \frac{dy}{dx} = 0 \text{, applying the chain rule of derivatives.}$$
$$\Rightarrow 2x + y \times 1 + x \times \frac{dy}{dx} + 2y \frac{dy}{dx} = 0$$
$$\Rightarrow 2x + y + (x + 2y) \frac{dy}{dx} = 0$$
Therefore, $\frac{dy}{dx} = -\frac{2x + y}{x + 2y}$.

6. Determine $\frac{dy}{dx}$ from the equation $x^2+x^2y+xy^2+y^3=81$.

Ans: The given equation is $x^2+x^2y+xy^2+y^3=81$.

Differentiating both sides of the equation with respect to x, gives

$$\frac{dy}{dx}(x^2+x^2y+xy^2+y^3) = \frac{d}{dx}(81)$$

$$\Rightarrow \frac{dy}{dx}(x^2) + \frac{dy}{dx}(x^2y) + \frac{dy}{dx}(xy^2) + \frac{dy}{dx}(y^3) = 0$$

$$\Rightarrow 3x^2 + \left[y\frac{d}{dx}(x^2) + x^2\frac{dy}{dx}\right] + \left[y^2\frac{d}{dx}(x) + x\frac{d}{dx}(y^2)\right] + 3y^2\frac{dy}{dx} = 0$$

$$\Rightarrow 3x^2 + \left[y \times 2 + x^2\frac{dy}{dx}\right] + \left[y^2 \times 1 + x \times 2y \times \frac{dy}{dx}\right] + 3y^2\frac{dy}{dx} = 0, \text{ applying chain rule.}$$

$$\Rightarrow (x^2 + 2xy + 3y^2)\frac{dy}{dx} + (3x^2 + 2xy + y^2) = 0$$
Therefore, $\frac{dy}{dx} = \frac{-(3x^2 + 2xy + 3y^2)}{(x^2 + 2xy + 3y^2)}.$

7. Determine $\frac{dx}{dy}$ from the equation $\sin^2 y + \cos xy = \pi$.

Ans: The given equation is $\sin^2 y + \cos xy = \pi$.

Differentiating both sides of the equation with respect to x, gives

$$\frac{d}{dx}(\sin^2 y + \cos xy) = \frac{d}{dx}\pi$$

$$\Rightarrow \frac{d}{dx}(\sin^2 y) + \frac{d}{dx}(\cos xy) = 0$$
....(1)

Applying the chain rule of derivatives gives

$$\frac{d}{dx}(\sin^2 y) = 2\sin y \frac{d}{dx}(\sin y) = 2\sin y \cos y \frac{dy}{dx} \qquad \dots (2)$$

$$\Rightarrow \frac{d}{dx}(\cos xy) = -\sin xy \frac{d}{dx}(xy) = -\sin xy \left[y \frac{d}{dx}(x) + x \frac{dy}{dx} \right] = -y \sin xy - x \sin xy \frac{dy}{dx} \dots (3)$$

From (1), (2) and (3), we obtain

$$2 \operatorname{sinycosy} \frac{dy}{dx} \operatorname{-ysinxy-xsinxy} \frac{dy}{dx} = 0$$
$$\Rightarrow (2 \operatorname{sinycosy-xsinxy}) \frac{dy}{dx} = \operatorname{ysinxy}$$
$$\Rightarrow (\sin 2y \operatorname{-xsinxy}) \frac{dx}{dy} = \operatorname{ysinxy}$$

Therefore,
$$\frac{dx}{dy} = \frac{ysinxy}{sin2y-xsinxy}$$
.

8. Determine $\frac{dy}{dx}$ from the equation $\sin^2 x + \cos^2 y = 1$.

Ans: The given equation is sin2x+cos2y=1.

$$\frac{dy}{dx}(\sin^2 x + \cos^2 y) = \frac{d}{dx}(1)$$

$$\Rightarrow \frac{d}{dx}(\sin^2 x) + \frac{d}{dx}(\cos^2 y) = 0$$

$$\Rightarrow 2\sin x \times \frac{d}{dx}(\sin x) + 2\cos y \times \frac{d}{dx}(\cos y) = 0$$

$$\Rightarrow 2\sin x \cos x + 2\cos y(-\sin y) \times \frac{dy}{dx} = 0$$

$$\Rightarrow \sin 2x - \sin 2y \frac{dy}{dx} = 0$$

$$\therefore \frac{dx}{dy} = \frac{\sin 2x}{\sin 2y}$$

9. Determine
$$\frac{dy}{dx}$$
 from the equation $y = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$.

Ans: The given equation is $y=\sin^{-1}\left(\frac{2x}{1+x^2}\right)$.

Now,
$$y = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$$

 $\Rightarrow \sin y = \frac{2x}{1+x^2}.$

Differentiating both sides of the equation with respect to x, gives

$$\frac{d}{dx}(\sin y) = \frac{d}{dx} \left(\frac{2x}{1+x^2} \right)$$
$$\Rightarrow \cos y \frac{dy}{dx} = \frac{d}{dx} \left(\frac{2x}{1+x^2} \right) \qquad \dots \dots (1)$$

Now, the function $\frac{2x}{1+x^2}$ is of the form of $\frac{u}{v}$.

Applying the quotient rule, gives

$$\frac{d}{dx}\left(\frac{2x}{1+x^{2}}\right) = \frac{(1+x^{2})\frac{d}{dx}(2x)-2x\times\frac{d}{dx}(1+x^{2})}{(1+x^{2})^{2}}$$
$$= \frac{(1+x^{2})\times2-2x\times[0+2x]}{(1+x^{2})^{2}} = \frac{2+2x^{2}-4x^{2}}{(1+x^{2})^{2}}$$
Therefore, $\frac{d}{dx}\left(\frac{2x}{1+x^{2}}\right) = \frac{2(1-x^{2})}{(1+x^{2})^{2}}$ (2)

It is given that,

$$siny = \frac{2x}{1+x^{2}}$$

$$\Rightarrow cosy = \sqrt{1-sin^{2}y} = \sqrt{1-\left(\frac{2x}{1+x^{2}}\right)^{2}} = \sqrt{\frac{\left(1+x^{2}\right)^{2}-4x^{2}}{(1+x^{2})^{2}}}$$

$$\Rightarrow cosy = \sqrt{\frac{\left(1-x^{2}\right)^{2}}{\left(1-x^{2}\right)^{2}}} = \frac{1-x^{2}}{1+x^{2}} \qquad \dots (3)$$

From the equation (1), (2) and (3), gives

$$\frac{1-x^{2}}{1+x^{2}}\frac{dy}{dx} = \frac{2(1-x^{2})}{(1+x^{2})^{2}}$$

Therefore, $\frac{dy}{dx} = \frac{2}{1+x^{2}}$.

10. Determine
$$\frac{dx}{dy}$$
 from the equation $y = \tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right)$, $-\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$.

Ans: The given function is $y=\tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right)$.

Now,
$$y=\tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right)$$

$$\Rightarrow \tan y = \frac{3x-x^3}{1-3x^2} \qquad \dots \dots (1)$$

According to the trigonometric formulas,

$$\tan y = \frac{3\tan \frac{y}{3} - \tan^3 \frac{y}{3}}{1 - 3\tan^2 \frac{y}{3}} \qquad \dots \dots (2)$$

By comparing the equations (1) and (2), gives

$$x = \tan \frac{y}{3}.$$
 (3)

$$\frac{d}{dx}(x) = \frac{d}{dx} \left(\tan \frac{y}{3} \right)$$

$$\Rightarrow 1 = \sec^2 \frac{y}{3} \times \frac{d}{dx} \left(\frac{y}{3} \right)$$

$$\Rightarrow 1 = \sec^2 \frac{y}{3} \times \frac{1}{3} \times \frac{dy}{dx}$$

$$\Rightarrow \frac{dy}{dx} = \frac{3}{\sec^2 \frac{y}{3}} = \frac{3}{1 + \tan^2 \frac{y}{3}}$$

Therefore, $\frac{dx}{dy} = \frac{3}{1 + x^2}$.


11. Determine
$$\frac{dy}{dx}$$
 from the equation $y = \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right)$, $0 < x < 1$.
Ans: The given equation is $y = \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right)$

$$\Rightarrow \cos y = \frac{1 - x^2}{1 + x^2}$$
$$\Rightarrow \frac{1 - \tan^2 \frac{y}{2}}{1 + \tan^2 \frac{y}{2}} = \frac{1 - x^2}{1 + x^2}.$$
 (1)

By comparing both sides of the equation (1) give

$$\tan\frac{y}{2} = x \qquad \dots \dots (2)$$

Differentiating both sides of the equation (2) with respect to x, gives



12. Determine $\frac{dy}{dx}$ from the equation $y=\sin^{-1}\left(\frac{1-x^2}{1+x^2}\right)$, 0<x<1

Ans: The given equation is $y=\sin^{-1}\left(\frac{1-x^2}{1+x^2}\right)$.

Now,
$$y = \sin^{-1}\left(\frac{1 - x^2}{1 + x^2}\right)$$

$$\Rightarrow \sin y = \frac{1 - x^2}{1 + x^2}.$$
(1)

Differentiating both sides of the equation with respect to x, gives

$$\frac{d}{dx}(\sin y) = \frac{d}{dx} \left(\frac{1 - x^2}{1 + x^2} \right) \qquad \dots \dots (2)$$

Using chain rule, we get

$$\frac{d}{dx}(\sin y) = \cos y \times \frac{dy}{dx} \qquad \dots (3)$$

$$\cos y = \sqrt{1 - \sin^2 y} = \sqrt{1 - \left(\frac{1 - x^2}{1 + x^2}\right)^2} = \sqrt{\frac{(1 + x^2)^2 - (1 - x^2)^2}{1 + x^2}} = \sqrt{\frac{4x^2}{(1 + x^2)^2}}, \text{ using the}$$

equation (1).

$$\Rightarrow \cos y = \frac{2x}{1+x^2} \qquad \dots \dots (4)$$

Therefore, from the equation (3) and (4) gives

$$\frac{d}{dx}(\sin y) = \frac{2x}{1+x^2} \frac{dy}{dx} \qquad \dots (5)$$

Now,

$$\frac{d}{dx} \left(\frac{1 - x^2}{1 + x^2} \right) = \frac{(1 + x^2)(1 - x^2) - (1 - x^2)(1 + x^2)}{(1 + x^2)^2} , \text{ applying the quotient rule.}$$

$$= \frac{(1 + x^2)(-2x) - (1 - x^2)(2x)}{(1 + x^2)^2}$$

$$= \frac{-2x - 2x^3 - 2x + 2x^3}{(1 - x^2)^2}$$

$$\Rightarrow \frac{d}{dx} \left(\frac{1 - x^2}{1 + x^2} \right) = -\frac{4}{(1 + x^2)^2} \qquad \dots (6)$$

Using the equations (2), (5), and (6), gives

$$\frac{2x}{1+x^2}\frac{dy}{dx} = \frac{-4x}{(1+x^2)^2}$$

Therefore, $\frac{dy}{dx} = \frac{-2}{1+x^2}$.

An alternate method:

$$y=\sin^{-1}\left(\frac{1-x^{2}}{1+x^{2}}\right)$$

$$\Rightarrow \sin y=\frac{1-x^{2}}{1+x^{2}}$$

$$\Rightarrow (1+x^{2})\sin y=1-x^{2}$$

$$\Rightarrow (1+\sin y)x^{2}=1-\sin y$$

$$\Rightarrow x^{2}=\frac{1-\sin y}{1+\sin y}$$

$$\Rightarrow x^{2}=\frac{\left(\cos \frac{y}{2}-\sin \frac{y}{2}\right)^{2}}{\left(\cos \frac{y}{2}+\sin \frac{y}{x}\right)^{2}}$$

$$\Rightarrow x=\frac{\cos \frac{y}{2}-\sin \frac{y}{2}}{\cos \frac{y}{2}+\sin \frac{y}{2}}$$

$$\Rightarrow x=\tan\left(\frac{\pi}{4}-\frac{\pi}{2}\right)$$

$$\frac{d}{dx}(x) = \frac{d}{dx} \left[\tan\left(\frac{\pi}{4} - \frac{y}{2}\right) \right]$$
$$\Rightarrow 1 = \sec^2\left(\frac{\pi}{4} - \frac{y}{2}\right) \times \frac{dy}{dx}\left(\frac{\pi}{4} - \frac{y}{2}\right)$$

$$\Rightarrow 1 = \left[1 + \tan^2 \left(\frac{\pi}{4} - \frac{y}{2} \right) \times \left(-\frac{1}{2} \times \frac{dy}{dx} \right) \right]$$
$$\Rightarrow 1 = (1 + x^2) \left(-\frac{1}{2} \frac{dy}{dx} \right)$$
Therefore, $\frac{dx}{dy} = \frac{-2}{1 + x^2}$.

13. Determine
$$\frac{dy}{dx}$$
 from the equation $y = \cos^{-1}\left(\frac{2x}{1+x^2}\right)$, $-1 < x < 1$

Ans: The given equation is $y = \cos^{-1}\left(\frac{2x}{1+x^2}\right)$.

Now,
$$y = \cos^{-1}\left(\frac{2x}{1+x^2}\right)$$

$$\Rightarrow \cos y = \frac{2x}{1+x^2}.$$
.....(1)

$$\frac{d}{dx}(\cos y) = \frac{d}{dx} \times \left(\frac{2x}{1+x^2}\right)$$

$$\Rightarrow -\sin y \times \frac{dy}{dx} = \frac{(1-x^2) \times \frac{d}{dx}(2x) - 2x \times \frac{d}{dx}(1+x^2)}{(1+x^2)^2}, \text{ applying the quotient rule.}$$

$$\Rightarrow -\sqrt{1-\cos^2 y} \frac{dy}{dx} = \frac{(1+x^2) \times 2 - 2x \times 2x}{(1+x^2)^2}$$

$$\Rightarrow \left[\sqrt{1-\left(\frac{2x}{1+x^2}\right)^2}\right] \frac{dx}{dy} = -\left[\frac{2(1-x)^2}{(1+x^2)^2}\right], \text{ using the equation (1).}$$

$$\Rightarrow \sqrt{\frac{(1-x^2)^2 - 4x^2}{(1+x^2)^2}} = \frac{dy}{dx} = \frac{-2(1-x)^2}{(1+x^2)}$$
$$\Rightarrow \sqrt{\frac{(1-x^2)^2}{(1+x^2)^2}} \frac{dy}{dx} = \frac{-2(1-x)^2}{(1+x^2)}$$
$$\Rightarrow \frac{1-x^2}{1+x^2} \times \frac{dy}{dx} = \frac{-2(1-x)^2}{(1+x^2)}$$
Therefore, $\frac{dy}{dx} = \frac{-2}{(1+x^2)}$.

14. Determine $\frac{dy}{dx}$ from the equation $y=\sin^{-1}\left(2x\sqrt{1-x^2}\right)$, $-\frac{1}{\sqrt{2}} < x < \frac{1}{\sqrt{2}}$.

Ans: The given equation is
$$y=\sin^{-1}(2x\sqrt{1-x^2})$$
.

Now,
$$y=\sin^{-1}(2x\sqrt{1-x^2})$$

 $\Rightarrow \sin y=2x\sqrt{1-x^2}$(1)

$$\cos y \frac{dy}{dx} = 2 \left[x \frac{d}{dx} \left(\sqrt{1 - x^2} \right) + \sqrt{1 - x^2} \frac{dx}{dx} \right]$$

$$\Rightarrow \sqrt{1 - \sin^2 y} \frac{dy}{dx} = 2 \left[\frac{x}{2} \times \frac{-2}{\sqrt{1 - x^2}} + \sqrt{1 - x^2} \right]$$

$$\Rightarrow \sqrt{1 - (2x\sqrt{1 - x^2})^2} = \frac{dy}{dx} = 2 \left[\frac{-x^2 + 1 - x^2}{\sqrt{1 - x^2}} \right], \text{ using the equation (1).}$$

$$\Rightarrow \sqrt{1 - 4x^2(1 - x^2)^2} \frac{dy}{dx} = 2 \left[\frac{1 - 2x^2}{\sqrt{1 - x^2}} \right]$$

$$\Rightarrow \sqrt{(1-2x)^2} \frac{dy}{dx} = 2 \left[\frac{1-2x^2}{\sqrt{1-x^2}} \right]$$
$$\Rightarrow (1-2x^2) \frac{dy}{dx} = 2 \left[\frac{1-2x^2}{\sqrt{1-x^2}} \right]$$
Therefore, $\frac{dy}{dx} = \frac{2}{\sqrt{1-x^2}}$.

15. Determine
$$\frac{dy}{dx}$$
 from the equation $y = \sec^{-1}\left(\frac{1}{2x^2-1}\right)$, $0 < x < \frac{1}{\sqrt{2}}$.
Ans: The given equation is $y = \sec^{-1}\left(\frac{1}{2x^2-1}\right)$.

Now,

$$y = \sec^{-1} \left(\frac{1}{2x^2 - 1} \right)$$

$$\Rightarrow \sec y = \frac{1}{2x^2 - 1}$$

$$\Rightarrow \cos y = 2x^2 - 1$$

$$\Rightarrow 2x^2 = 1 + \cos y$$

$$\Rightarrow 2x^2 = 2\cos^2 \frac{y}{2}$$

$$\Rightarrow x = \cos \frac{y}{2}$$
(1)

$$\frac{d}{dx}(x) = \frac{d}{dx} \left(\cos \frac{y}{2} \right)$$
$$\Rightarrow 1 = \sin \frac{y}{2} \times \frac{d}{dx} \left(\frac{y}{2} \right)$$

$$\Rightarrow \frac{-1}{\sin\frac{y}{2}} = \frac{1}{2} \frac{dy}{dx}$$
$$\Rightarrow \frac{dy}{dx} = \frac{-2}{\sin\frac{y}{2}} = \frac{-2}{\sqrt{1 - \cos^2\frac{y}{2}}} = \frac{-2}{\sqrt{1 - x^2}}, \text{ using the equation (1).}$$
Therefore, $\frac{dy}{dx} = \frac{-2}{\sqrt{1 - x}}.$

Exercise 5.4

1. Find the derivative of the function $y = \frac{e^x}{\sin x}$ with respect to x.

Ans: The given function is $y = \frac{e^x}{\sin x}$.

Then, we have

$$\frac{dy}{dx} = \frac{\sin x \frac{d}{dx} (e^x) \cdot e^x \frac{d}{dx} (\sin x)}{\sin^2 x}, \text{ by applying the quotient rule of derivatives.}$$
$$= \frac{\sin x \times (e^x) \cdot e^x \times (\cos x)}{\sin^2 x}$$

Therefore, the derivative of the function y is

$$\frac{dy}{dx} = \frac{e^{x}(\sin x - \cos x)}{\sin^{2} x}, \ x \neq n\pi, \ n \in \mathbb{Z}.$$

2. Find the derivative of the function $y=e^{\sin^{-1}x}$.

Ans: The given function is $y=e^{\sin^{-1}x}$.

Then, we have

$$\frac{dy}{dx} = \frac{d}{dx} (e^{\sin^{-1}x})$$
$$= e^{\sin^{-1}x} \times \frac{d}{dx} (\sin^{-1}x)$$
$$= e^{\sin^{-1}x} \times \frac{1}{\sqrt{1 - x^2}}$$
$$= \frac{e^{\sin^{-1}x}}{\sqrt{1 - x^2}}$$

Therefore, the derivative of the function y is

$$\frac{dy}{dx} = \frac{e^{\sin^{-1}x}}{\sqrt{1-x^2}}, x \in (-1,1).$$

3. Find the derivative of the function $y = e^{x^3}$ with respect to x.

Ans: The given function is $y=e^{x^3}$.

Given: $y = e^{x^3}$ Diff w.r.tx $\frac{dy}{dx} = e^{x^3} \cdot \frac{d}{dx}(x^3)$ $\frac{dy}{dx} = e^{x^3} \cdot 3x^2.$

4. Find the derivative of the function is $y = sin(tan^{-1}e^{-x})$ with respect to x.

Ans: The given function is $y=sin(tan^{-1}e^{-x})$.

Now, applying the chain rule of derivatives, give

$$\frac{dy}{dx} = \frac{d}{dx} \left[\sin(\tan^{-1}e^{-x}) \right]$$
$$= \cos(\tan^{-1}e^{-x}) \times \frac{d}{dx} (\tan^{-1}e^{-x})$$
$$= \cos(\tan^{-1}e^{-x}) \times \frac{1}{1 + (e^{-x})} \times \frac{d}{dx} (e^{-x})$$

$$= \frac{\cos(\tan^{-1}e^{-x})}{1+(e^{-x})} \times e^{-x} \times \frac{d}{dx}(-x)$$
$$= \frac{e^{-x}\cos(\tan^{-1}e^{-x})}{1+e^{-2x}} \times (-1)$$

Therefore, the derivative of the function y is

$$\frac{dy}{dx} = \frac{-e^{-x}\cos(\tan^{-1}e^{-x})}{1+e^{-2x}}.$$

5. Find the derivative of the function $y = log(cos(e^x))$

Ans: Let $y=log(cos(e^x))$

Now, by applying the chain rule of derivatives give

$$\frac{dy}{dx} = \frac{d}{dx} \left[\log(\cos(e^x)) \right]$$
$$= \frac{1}{\cos e^x} \times \frac{d}{dx} (\cos(e^x))$$
$$= \frac{1}{\cos e^x} \times (-\sin(e^x)) \times \frac{d}{dx} (e^x)$$
$$= \frac{-\sin e^x}{\cos e^x} \times e^x$$

Therefore, the derivative of the function y is

$$\frac{\mathrm{dy}}{\mathrm{dx}} = -\mathrm{e}^{\mathrm{x}} \mathrm{tan}\left(\mathrm{e}^{\mathrm{x}}\right), \ \mathrm{x} \neq (2\mathrm{n}+1)\frac{\pi}{2}, \mathrm{n} \in \mathrm{N}.$$

6. Find the derivative of the function $y=e^{x}+e^{x^{2}}+...+e^{x^{5}}$ with respect to x.

Ans: The given function is $y=e^{x}+e^{x^{2}}+...+e^{x^{5}}$.

Then, differentiating with respect to x both sides, give

$$\frac{dy}{dx} = \frac{d}{dx}(e^{x}+e^{x^{2}}+...+e^{x^{5}})$$

$$= \frac{d}{dx}(e^{x}) + \frac{d}{dx}(e^{x^{2}}) + \frac{d}{dx}(e^{x^{4}}) + \frac{d}{dx}(e^{x^{5}}), \text{ applying the sum rule of derivatives.}$$
$$= e^{x} + \left[e^{x^{2}} \times \frac{d}{dx}(x^{2})\right] + \left[e^{x^{3}} \times \frac{d}{dx}(x^{3})\right] + \left[e^{x^{4}} \times \frac{d}{dx}(x^{4})\right] + \left[e^{x^{5}} \times \frac{d}{dx}(x^{5})\right]$$
$$= e^{x} + (e^{x^{2}} \times 2x) + (e^{x^{3}} \times 3x^{2}) + (e^{x^{4}} \times 4x^{3}) + (e^{x^{5}} \times 5x^{4})$$

Therefore, the derivative of the function y is

$$\frac{dy}{dx} = e^{x} + 2xe^{x^{2}} + 3x^{2}e^{x^{3}} + 4x^{3}e^{x^{4}} + 5x^{4}e^{x^{5}}.$$

- 7. Find the derivative of the function $y=\sqrt{e^{\sqrt{x}}}$, x>0 with respect to x.
- **Ans:** The given function is $y=\sqrt{e^{\sqrt{x}}}$.

Then squaring both sides both sides of the equation give

$$y^2 = e^{\sqrt{x}}$$

Now, differentiating both sides with respect to x gives

$$\begin{aligned} \frac{d}{dx}(y^2) &= \frac{d}{dx}(e^{\sqrt{x}}) \\ \Rightarrow 2y\frac{dy}{dx} &= e^{\sqrt{x}}\frac{d}{dx}(\sqrt{x}) \\ \Rightarrow 2y\frac{dy}{dx} &= e^{\sqrt{x}}\frac{1}{2} \times \frac{1}{\sqrt{x}} \\ \Rightarrow \frac{dy}{dx} &= \frac{e^{\sqrt{x}}}{4y\sqrt{x}} \\ \Rightarrow \frac{dy}{dx} &= \frac{e^{\sqrt{x}}}{4y\sqrt{x}}, \text{ substituting the value of } y. \end{aligned}$$

Therefore,

$$\frac{\mathrm{dy}}{\mathrm{dx}} = \frac{\mathrm{e}^{\sqrt{x}}}{4\sqrt{\mathrm{x}\mathrm{e}^{\sqrt{x}}}}, \, \mathrm{x} > 0.$$

8. Find the derivative of the function y=log(logx), x>1.

Ans: The given function is y=log(logx).

Now, differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{d}{dx} [log(logx)]$$

$$= \frac{1}{logx} \times \frac{d}{dx} (logx), \text{ by applying the chain rule of derivatives.}$$

$$= \frac{1}{logx} \times \frac{1}{x}$$
Therefore, $\frac{dy}{dx} = \frac{1}{xlog}, x > 1.$

9. Find the derivative of the function $y = \frac{\cos x}{\log x}$, x>0 with respect to x.

Ans: The given function is
$$y = \frac{\cos x}{\log x}$$
.

Differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{\frac{d}{dx}(\cos x) \times \log x \cdot \cos x \times \frac{d}{dx}(\log x)}{(\log x)^2}, \text{ by applying the quotient rule.}$$
$$= \frac{\frac{-\sin x \log x \cdot \cos x \times \frac{1}{x}}{(\log x)^2}}{(\log x)^2}$$

Therefore,

$$\frac{dy}{dx} = \frac{-[x \log x \sin x + \cos x]}{x (\log x)^2}, x > 0.$$

10. Find the derivative of the function $y=cos(logx+e^x)$, x>0 with resepct to x

Ans: The given function is $y=\cos(\log x+e^x)$.

Then differentiating both sides with respect to x gives

 $\frac{dy}{dx} = \frac{d}{dx} \Big[\cos \Big(\log x + e^x \Big) \Big].$

$$\Rightarrow \frac{dy}{dx} = -\sin[\log x + e^x] \times \frac{d}{dx} (\log x + e^x), \text{ by applying the chain rule of derivatives.}$$
$$= \sin(\log x + e^x) \times \left[\frac{d}{dx} (\log x) + \frac{d}{dx} (e^x) \right]$$
$$= \sin(\log x + e^x) \times \left(\frac{1}{x} + e^x \right)$$
Therefore, $\frac{dy}{dx} = \left(\frac{1}{x} + e^x \right) \sin(\log x + e^x), x > 0.$

Exercise 5.5

1. Find the derivative of the function $y=\cos 2x \times \cos 3x$ with respect to x

Ans: The given function is $y=\cos 2x \times \cos 3x$.

First, taking logarithm both sides of the equation give,

logy=log(cosx×cos2x×cos3x)

 \Rightarrow logy=log(cosx)+log(cos2x)+log(cos3x), by the property of logarithm.

Now, differentiating both sides of the equation with respect to x gives

$$\frac{1}{y}\frac{dy}{dx} = \frac{1}{\cos x} \times \frac{d}{dx}(\cos x) + \frac{1}{\cos 2x} \times \frac{d}{dx}(\cos 2x) + \frac{1}{\cos 3x} \times \frac{d}{dx}(\cos 3x)$$
$$\Rightarrow \frac{dy}{dx} = y \left[-\frac{\sin x}{\cos x} - \frac{\sin 2x}{\cos 2x} \times \frac{d}{dx}(2x) - \frac{\sin 3x}{\cos 3x} \times \frac{d}{dx}(3x) \right]$$

Therefore,

$$\frac{dy}{dx} = -\cos \times \cos 2x \times \cos 3x [\tan x + 2\tan 2x + 3\tan 3x].$$

2. Find the derivative of the function $y = \sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}}$ with respect to

х.

Ans: The given function is
$$y = \sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}}$$

First taking logarithm both sides of the equation give

$$logy=log \sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}}$$

$$\Rightarrow logy=\frac{1}{2}log \left[\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}\right]$$

$$\Rightarrow logy=\frac{1}{2}[log\{(x-1)(x-2)\}-log\{(x-3)(x-4)(x-5)\}]$$

$$\Rightarrow logy=\frac{1}{2}[log(x-1)+log(x-2)-log(x-3)-log(x-4)-log(x-5)]$$

Now, differentiating both sides of the equation with respect to x gives

$$\frac{dy}{dx} = \frac{1}{2} \frac{d}{dx} [\log(x-1) + \log(x-2) - \log(x-3) - \log(x-4) - \log(x-5)].$$

$$\frac{1}{y} \frac{dy}{dx} = \frac{1}{2} \left[\frac{1}{x-1} \times \frac{d}{dx} (x-1) + \frac{1}{x-2} \times \frac{d}{dx} (x-2) - \frac{1}{x-3} \times \frac{d}{dx} (x-3) - \frac{1}{x-4} \times \frac{d}{dx} (x-4) - \frac{1}{x-5} \times \frac{d}{dx} (x-5) \right]$$

$$\Rightarrow \frac{dy}{dx} = \frac{y}{2} \left(\frac{1}{x-1} + \frac{1}{x-2} + \frac{1}{x-3} + \frac{1}{x-4} + \frac{1}{x-5} \right)$$

Therefore,

$$\frac{\mathrm{dy}}{\mathrm{dx}} = \frac{1}{2} \sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}} \left[\frac{1}{x-1} + \frac{1}{x-2} + \frac{1}{x-3} + \frac{1}{x-4} + \frac{1}{x-5} \right].$$

3. Find the derivative of the function $y=(logx)^{cosx}$ with respect to x.

Ans: The given function is $y=(\log x)^{\cos x}$.

First, taking logarithm both sides of the equation give

logy=cosx.log(logx).

$$\frac{1}{y} \times \frac{dy}{dx} = \frac{d}{dx}(\cos x) \times \log(\log x) + \cos x \times \frac{d}{dx}[\log(\log x)]$$

$$\Rightarrow \frac{1}{y} \times \frac{dy}{dx} = -\sin x \log(\log x) + \cos x \times \frac{1}{\log x} \times \frac{d}{dx}(\log x), \text{ by applying the chain rule.}$$

$$\Rightarrow \frac{dy}{dx} = y \left[-\sin x \log(\log x) + \frac{\cos x}{\log x} \times \frac{1}{x} \right]$$

Therefore,

$$\frac{dy}{dx} = (\log x)^{\cos x} \left[\frac{\cos x}{x \log x} - \sin \times \log(\log x) \right].$$

4. Determine the derivative of the function $y=x^x-2^{sinx}$ with respect to x.

Ans: The given function is $y=x^x-2^{sinx}$.

Now, let
$$x^x = u$$
 (1)

and
$$2^{\sin x} = v$$
.(2)

Then differentiating the equation (3) with respect to x gives

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}u}{\mathrm{d}x} - \frac{\mathrm{d}v}{\mathrm{d}x} \qquad \dots \dots (4)$$

Now, taking logarithm both sides of the equation (1) give

$$log(u) = log(x^{x})$$
$$\Rightarrow log u = xlog x$$

$$\frac{1}{u}\frac{du}{dx} = \left[\frac{d}{dx}(x) \times \log x + x \times \frac{d}{dx}(\log x)\right]$$

$$\Rightarrow \frac{du}{dx} = u \left[1 \times \log x + x \times \frac{1}{x} \right]$$

$$\Rightarrow \frac{du}{dx} = x^{x} (\log x + 1)$$

$$\Rightarrow \frac{du}{dx} = x^{x} (1 + \log x) \qquad \dots \dots (5)$$

Now, taking logarithm both sides of the equation (2) give

$$log(2^{sinx}) = logv$$

 $\Rightarrow logv = sinx \times log2.$

Differentiating both sides of the equation with respect to x, give

$$\frac{1}{v} \times \frac{dv}{dx} = \log 2 \times \frac{d}{dx} (\sin x)$$

$$\Rightarrow \frac{dv}{dx} = v \log 2 \cos x$$

$$\Rightarrow \frac{dv}{dx} = 2^{\sin x} \cos x \log 2 \qquad \dots \dots (6)$$

Therefore, from the equation (4), (5) and (6) give

$$\frac{\mathrm{dy}}{\mathrm{dx}} = x^{x} (1 + \log x) - 2^{\sin x} \cos x \log 2.$$

5. Find the derivative of the function $y=(x+3)^2(x+4)^3(x+5)^4$ with respect to x

Ans: The given function is $y=(x+3)^2(x+4)^3(x+5)^4$.

First, taking logarithm both sides of the equation give

$$\log = \log \left[(x+3)^2 (x+4)^3 (x+5)^4 \right]$$

 $\Rightarrow \log y = 2\log(x+3) + 3\log(x+4) + \log 4(x+5)$

$$\frac{1}{y} \times \frac{dy}{dy} = 2 \times \frac{1}{x-3} \times \frac{d}{dz} (x+3) + 3 \times \frac{1}{x+4} \times \frac{d}{dx} (x+4) + 4 \times \frac{1}{x+5} \times \frac{d}{dx} (x+5)$$

$$\Rightarrow \frac{dy}{dx} = y \left[\frac{2}{x+3} + \frac{3}{x+4} + \frac{4}{x+5} \right]$$

$$\Rightarrow \frac{dy}{dx} = (x+3)^2 (x+4)^3 (x+5)^4 \times \left[\frac{2}{x+3} + \frac{3}{x+4} + \frac{4}{x+5} \right]$$

$$\Rightarrow \frac{dy}{dx} = (x+3)^2 (x+4)^3 (x+5)^4 \times \left[\frac{2(x+4)(x+5) + 3(x+3)(x+5) + 4(x+3)(x+4)}{(x+3)(x+4)(x+5)} \right]$$

$$\Rightarrow \frac{dy}{dx} = (x+3)^2 (x+4)^2 (x+5)^2 - \left[2(x^2 + 9x + 20) + 3(x^2 + 9x + 15) + 4(x^2 + 7x + 12) \right]$$

Therefore,

$$\frac{dy}{dx} = (x+3)(x+4)^2(x+5)^3(9x^2+70x+133)$$

6. Find the derivative of the function $y = \left(x + \frac{1}{x}\right)^x + x^{\left(1 + \frac{1}{x}\right)}$ with respect to x.

Ans: The given function is $y = \left(x + \frac{1}{x}\right)^x + x^{\left(1 + \frac{1}{x}\right)}$.

First, let
$$u = \left(x + \frac{1}{x}\right)^x$$
 and $v = x^{\left(1 + \frac{1}{x}\right)}$

Therefore,
$$y=u+v$$
.(1)

Differentiating the equation (1) both sides with respect to x give

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} = \frac{dv}{dx} \qquad \dots \dots (2)$$

Now, $u = \left(x + \frac{1}{x}\right)^{x}$

$$\Rightarrow \log u = \log \left(x + \frac{1}{x} \right)^{x}$$
$$\Rightarrow \log u = x \log \left(x + \frac{1}{x} \right)$$

$$\frac{1}{u}\frac{du}{dx} = \frac{d}{dx}(x) \times \log\left(x + \frac{1}{x}\right) + x \times \frac{d}{dx}\left[\log\left(x + \frac{1}{x}\right)\right]$$

$$\Rightarrow \frac{1}{u}\frac{du}{dx} = 1 \times \log\left(x + \frac{1}{x}\right) + x \times \frac{1}{\left(x + \frac{1}{x}\right)} \times \frac{d}{dx}\left(x + \frac{1}{x}\right)$$

$$\Rightarrow \frac{du}{dx} = u\left[\log\left(x + \frac{1}{x}\right) + \frac{x}{\left(x + \frac{1}{x}\right)} \times \left(x + \frac{1}{x^{2}}\right)\right]$$

$$\Rightarrow \frac{du}{dx} = \left(x + \frac{1}{x}\right)^{x}\left[\log\left(x + \frac{1}{x}\right) + \frac{\left(x - \frac{1}{x}\right)}{\left(x + \frac{1}{x}\right)}\right]$$

$$\Rightarrow \frac{du}{dx} = \left(x + \frac{1}{x}\right)^{x}\left[\log\left(x + \frac{1}{x}\right) + \frac{x^{2} + 1}{x^{2} - 1}\right]$$

$$\Rightarrow \frac{du}{dx} = \left(x + \frac{1}{x}\right)^{2}\left[\frac{x^{2} + 1}{x^{2} - 1} + \log\left(x + \frac{1}{x}\right)\right] \qquad \dots (3)$$
Also, $v = x^{\left(x + \frac{1}{x}\right)}$

$$\Rightarrow \log v = \log\left[x^{x^{\left(x + \frac{1}{x}\right)}}\right]$$

Hence, from the equations (2), (3) and (4), give

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \left(x + \frac{1}{x}\right)^{x} \left[\frac{x^{2} - 1}{x^{2} + 1} + \log\left(x + \frac{1}{x}\right)\right] + x^{\left(x + \frac{1}{x}\right)} \left(\frac{x + 1 - \log x}{x^{2}}\right).$$

7. Determine derivative of the function $y=(logx)^{x}+x^{logx}$ with respect to x.

Ans: The given function is $y=(\log x)^x + x^{\log x}$.

Then, let $u=(\log x)^x$ and $v=x^{\log x}$.

Therefore, y=u+v.

Differentiating both sides of the equation with respect to x gives

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} \qquad \dots \dots (1)$$

Now, $u=(logx)^x$

$$\Rightarrow \log u = \log \left[(\log x)^{x} \right]$$
$$\Rightarrow \log u = x \log (\log x)$$

$$\frac{1}{u}\frac{du}{dx} = \frac{d}{dx}(x) \times \log(\log x) + x \times \frac{d}{dx} \left[\log(\log x)\right]$$

$$\Rightarrow \frac{du}{dx} = u \left[1 \times \log(\log x) + x \times \frac{1}{\log x} \times \frac{d}{dx} (\log x) \right]$$

$$\Rightarrow \frac{du}{dx} = (\log x)^{x} \left[\log(\log x) + \frac{x}{\log x} \times \frac{1}{x} \right]$$

$$\Rightarrow \frac{du}{dx} = (\log x)^{x} \left[\log(\log x) + \frac{1}{\log x} \right]$$

$$\Rightarrow \frac{du}{dx} = (\log x)^{x} = \left[\frac{\log(\log x) \times \log x + 1}{\log x} \right]$$

$$\frac{du}{dx} = (\log x)^{x-1} \left[1 + \log x \times \log(\log x) \right] \qquad \dots \dots (2)$$
Again, $v = x^{\log x}$

$$\Rightarrow \log v = \log \left(x^{\log x} \right)$$

$$\Rightarrow \log v = \log x \log x = (\log x)^{2}$$

$$\frac{1}{v} \times \frac{dx}{dx} = \frac{d}{dx} \left[(\log x)^{2} \right]$$

$$\Rightarrow \frac{1}{v} \times \frac{dx}{dx} = 2(\log x) \times \frac{d}{dx} (\log x)$$

$$\Rightarrow \frac{dv}{dx} = 2x^{\log x} \frac{\log x}{x}$$

$$\Rightarrow \frac{dv}{dx} = 2x^{\log x} \times \log x \qquad \dots \dots (3)$$

Hence, from the equations (1), (2), and (3), gives

$$\frac{dy}{dx} = (\log x)^{x+1} [1 + \log x \times \log(\log x)] + 2x^{\log x-1} \times \log x.$$

8. Find the derivative of the function $y = (sinx)^2 + sin^{-1}\sqrt{x}$ with respect to x

Ans: The given function is $y=(\sin x)^{x}+\sin^{-1}\sqrt{x}$.

```
Now, let u=(\sin x)^x and v=\sin^{-1}\sqrt{x}.
```

Therefore, y=u+v.

Then, differentiating both sides of the equation with respect to x gives

 $\frac{dy}{dx} = \frac{du}{dx} - \frac{dv}{dx} \qquad \dots \dots (1)$ Now, $u = (\sin x)^x$ $\Rightarrow \log u = x \log(\sin x)^x$ $\Rightarrow \log u = x \log(\sin x)$ Differentiating both sides of the equation with respect to x gives

$$\frac{1}{u}\frac{du}{dx} = \frac{d}{dx}(x) \times \log(\sin x) + x \times \frac{d}{dx}[\log(\sin x)]$$

$$\Rightarrow \frac{du}{dx} = u\left[1 \times \log(\sin x) + x \times \frac{1}{\sin x} \times \frac{d}{dx}(\sin x)\right]$$

$$\Rightarrow \frac{du}{dx} = (\sin x)^{x} \left[\log(\sin x) + \frac{x}{\sin x} \times \cos x\right]$$

$$\Rightarrow \frac{du}{dx} = (\sin x)^{x} (x \cot x + \log \sin x) \qquad \dots \dots (2)$$
Again, $v = \sin^{-1}\sqrt{x}$.

$$\frac{dv}{dx} = \frac{1}{\sqrt{1 - (\sqrt{x})^2}} \times \frac{d}{dx} (\sqrt{x})$$
$$\Rightarrow \frac{dv}{dx} = \frac{1}{\sqrt{1 - x}} \times \frac{1}{2\sqrt{x}}$$

$$\Rightarrow \frac{\mathrm{d}v}{\mathrm{d}x} = \frac{1}{2\sqrt{x-x^2}}$$

Hence, from the equations (1), (2) and (3), gives

 $\frac{dv}{dx} = (\sin x)^2 (x \cot x + \log \sin x) + \frac{1}{2\sqrt{x - x^2}}.$

9. Find the derivative of the function $y = x^{sinx} + (sinx)^{cosx}$ with respect to x.

Ans: The given function is $y=x^{sinx}+(sinx)^{cosx}$.

Then, let $u=x^{sinx}$ and $v=(sinx)^{cosx}$.

Therefore, y=u+v.

Differentiating both sides of the equation with respect to x gives

 $\frac{dy}{dx} = \frac{du}{dx} - \frac{dv}{dx} \qquad \dots \dots (1)$ Now, $u = x^{sinx}$ $\Rightarrow logu = xlog(x^{sinx})$ $\Rightarrow logu = sinxlogx$

Differentiating both sides of the equation with respect to x gives

$$\frac{1}{u}\frac{du}{dx} = \frac{d}{dx}(\sin x) \times \log x + \sin x \times \frac{d}{dx}(\log x)$$

$$\Rightarrow \frac{du}{dx} = u = \left[\cos x \log x + \sin x \times \frac{1}{x}\right]$$

$$\Rightarrow \frac{du}{dx} = x^{\sin x} \left[\cos x \log x + \frac{\sin x}{x}\right] \qquad \dots \dots (2)$$

Again, v=(sinx)^{cosx}

 $\Rightarrow \log v = \log(\sin x)^{\cos x}$ $\Rightarrow \log v = \cos \log(\sin x)$

Then, differentiating both sides of the equation with respect to x gives

$$\frac{1}{v}\frac{dv}{dx} = \frac{d}{dx}(\cos x) \times \log(\sin x) + \cos x \times \frac{d}{dx}[\log(\sin x)]$$

$$\Rightarrow \frac{dv}{dx} = v \left[-\sin x \times \log(\sin x) + \cos x \times \frac{1}{\sin x} \times \frac{d}{dx}(\sin x) \right]$$

$$\Rightarrow \frac{du}{dx} = (\sin x)^{\cos x} \left[-\sin x \log \sin x + \cot x \cos x \right]$$

$$\Rightarrow \frac{dv}{dx} = (\sin x)^{\cos x} \left[\cos x \cot x + \sin x \log \sin x \right] \qquad \dots (3)$$

Hence, from the equations (1), (2) and (3), gives

$$\frac{du}{dx} = x^{sinx} \left(cosxlogx + \frac{sinx}{x} \right) + (sinx)^{cosx} [cosxcotx + sinxlogsinx].$$

10. Find the derivative function $y = x^{xcosx} + \frac{x^2 + 1}{x^2 - 1}$ with respect to x.

Ans: The given function is $y=x^{x\cos x}+\frac{x^2+1}{x^2-1}$.

First, let
$$u=x^{xcosx}$$
 and $v=\frac{x^2+1}{x^2-1}$.

Therefore, y=u+v.

Differentiating both sides of the equation with respect to x gives

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} - \frac{dv}{dx} \qquad \dots \dots (1)$$

Now, $u = x^{x \cos x}$.

$$\frac{1}{u}\frac{du}{dx} = \frac{d}{dx}(x) \times \cos x \log x + x \times \frac{d}{dx}(\cos x) \times \log x + x \cos x \times \frac{d}{dx}(\log x)$$

$$\Rightarrow \frac{du}{dx} = u \left[1 \times \cos x \times \log x + x \times (-\sin x) \log x + x \cos x \times \frac{1}{x} \right]$$

$$\Rightarrow \frac{du}{dx} = x^{x \cos x} (\cos x \log x - x \sin x \log x + \cos x) \qquad \dots \dots (2)$$
Again, $v = \frac{x^2 + 1}{x^2 - 1}$

$$\Rightarrow \log v = \log(x^2 + 1) - \log(x^2 - 1)$$

$$\frac{1}{v} = \frac{dv}{dx} = \frac{2x}{x^{2}+1} - \frac{2x}{x^{2}-1}$$

$$\Rightarrow \frac{dv}{dx} = v \left[\frac{2x(x^{2}-1) - 2x(x^{2}+1)}{(x^{2}+1)(x^{2}-1)} \right]$$

$$\Rightarrow \frac{du}{dx} = \frac{x^{2}+1}{x^{2}-1} \times \left[\frac{-4x}{(x^{2}+1)(x^{2}-1)} \right]$$

$$\Rightarrow \frac{dv}{dx} = \frac{-4x}{(x^{2}-1)^{2}} \qquad \dots \dots (3)$$

Hence, from the equations (1), (2) and (3), give

$$\frac{dv}{dx} = x^{x\cos x} \left[\cos x(1 + \log x) - x\sin x \log x \right] - \frac{4x}{(x^2 - 1)^2}.$$

11. Find the derivative of the function $y=(x\cos x)^2+(x\sin x)^{\frac{1}{2}}$ with respect to x

Ans: The given function is $y=(x\cos x)^2+(x\sin x)^{\frac{1}{2}}$.

Then, let $u=(x\cos x)^2$ and $v=(x\sin x)^{\frac{1}{2}}$.

Therefore, y=u+v.

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} \qquad \dots \dots (1)$$

Again,
$$u = (x \cos x)^2$$

 $\Rightarrow \log u = \log(x \cos x)^{2}$ $\Rightarrow \log u = u \log(x \cos x)$ $\Rightarrow \log u = x[\log x + \log \cos x]$ $\Rightarrow \log u = x \log x + x \log \cos x$

$$\frac{1}{u}\frac{du}{dx} = \frac{d}{dx}(x\log \cos x)$$

$$\Rightarrow \frac{du}{dx} = u \left[\left\{ \log x \times \frac{d}{dx}(x) + x \times \frac{d}{dx}(\log x) \right\} + \left\{ \log \cos x \times \frac{d}{dx}(x) + x \times \frac{d}{dx}(\log \cos x) \right\} \right]$$

$$\Rightarrow \frac{du}{dx} = (x\cos x)^{x} \left[\left\{ \log x \times 1 + x \times \frac{1}{2} \right\} + \left\{ \log \cos x - 1 + x \times \frac{1}{\cos x} \times \frac{d}{dx}(\cos x) \right\} \right]$$

$$\Rightarrow \frac{du}{dx} = (x\cos x)^{x} \left[\left\{ \log x + 1 \right\} + \left\{ \log \cos x - 1 + \frac{x}{\cos x} \times (-\sin x) \right\} \right]$$

$$\Rightarrow \frac{du}{dx} = (x\cos x)^{x} \left[(\log x + 1) + (\log \cos x - x\tan x) \right]$$

$$\Rightarrow \frac{du}{dx} = (x\cos x)^{x} \left[1 - x\tan x + (\log x + \log \cos x) \right]$$
Therefore,

$$\frac{du}{dx} = (x\cos x)^{x} [1 - x\tan x + (\log x(x\cos x))] \qquad \dots \dots (2)$$

Again, $v = (x\sin x)^{\frac{1}{x}}$
$$\Rightarrow \log v = \log (x\sin x)^{\frac{1}{x}}$$

$$\Rightarrow \log v = \frac{1}{x} \log(x \sin x)$$
$$\Rightarrow \log v = \frac{1}{x} (\log x + \log \sin x)$$
$$\Rightarrow \log v = \frac{1}{x} \log x + \frac{1}{x} \log \sin x$$

$$\frac{1}{v}\frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{x}\log x\right) + \frac{d}{dx}\left[\frac{1}{x}\log(\sin x)\right]$$

$$\Rightarrow \frac{1}{v}\frac{dv}{dx} = \begin{bmatrix}\frac{1}{x}\log x \times \frac{d}{dx}\left(\frac{1}{x}\right) + \frac{1}{x} \times \frac{d}{dx}\left(\log x\right)\end{bmatrix} + \begin{bmatrix}\log(\sin x) \times \frac{d}{dx}\left(\frac{1}{x}\right) + \frac{1}{x} \times \frac{d}{dx}\left((\log \sin x)\right)\end{bmatrix}$$

$$\Rightarrow \frac{1}{v}\frac{dv}{dx} = \begin{bmatrix}\frac{1}{x}\log x \times \left(-\frac{1}{x^2}\right) + \frac{1}{x} \times \frac{1}{x}\end{bmatrix} + \begin{bmatrix}\log(\sin x) \times \left(-\frac{1}{x^2}\right) + \frac{1}{x} \times \frac{1}{\sin x} \times \frac{d}{dx}(\sin x)\end{bmatrix}$$

$$\Rightarrow \frac{1}{v}\frac{dv}{dx} = \frac{1}{x^2}(1 - \log x) + \begin{bmatrix}\frac{1 - \log x}{x^2} + \frac{1}{x \sin x} \times \cos x\end{bmatrix}$$

$$\Rightarrow \frac{1}{v}\frac{dv}{dx} = \frac{1}{x^2}(x \sin x)^{\frac{1}{x}} + \begin{bmatrix}\frac{1 - \log x - \log(\sin x) + x \cot x}{x^2}\end{bmatrix}$$

Therefore,

$$\frac{dv}{dx} = (x \sin x)^{\frac{1}{x}} \left[\frac{1 - \log(x \sin x) + x \cot x}{x^2} \right] \qquad \dots \dots (3)$$

Hence, from the equations (1), (2) and (3), gives

$$\Rightarrow \frac{dy}{dx} = (x\cos x)^2 \left[1 - x\tan x + \log(x\cos x)\right] + (x\sin x)^{\frac{1}{x}} \left[\frac{1 - \log(x\sin x) + x\cot x}{x^2}\right].$$

12. Determine $\frac{dy}{dx}$ from the equation $x^y + y^x = 1$.

Ans: The given function is $x^y + y^x = 1$.

Then, let $x^y = u$ and $y^x = v$.

Therefore, u+v=1.

Differentiating both sides of the equation with respect to x gives

$$\frac{du}{dx} + \frac{dv}{dy} = 0$$
Now, $u = x^{y}$ (1)
$$\Rightarrow \log u = \log(x^{y})$$

$$\Rightarrow \log u = y \log x$$

Differentiating both sides of the equation with respect to x gives

$$\frac{1}{u}\frac{du}{dx} = \log x \frac{dy}{dx} + y \times \frac{d}{dx}(\log x)$$

$$\Rightarrow \frac{du}{dx} = u \left[\log x \frac{dy}{dx} + y \times \frac{1}{x} \right]$$

Therefore, $\frac{du}{dx} = x^{y} \left[\log x \frac{dy}{dx} + \frac{y}{x} \right]$ (2)

Also, v=y^x

Taking logarithm both sides of the equation give

$$\Rightarrow \log v = \log(y^3)$$
$$\Rightarrow \log v = x \log y$$

$$\frac{1}{v} \times \frac{dv}{dx} = \log y \times \frac{d}{dx}(x) + x \times \frac{d}{dx}(\log y)$$

$$\Rightarrow \frac{dv}{dx} = v \left(logy \times 1 + x \times \frac{1}{y} \times \frac{dy}{dx} \right)$$

Therefore, $\frac{dv}{dx} = y^{x} \left(logy + \frac{x}{y} \frac{dy}{dx} \right)$ (3)

So, from the equation (1), (2) and (3), gives

$$x^{y}\left(\log x \frac{dy}{dx} + \frac{y}{x}\right) + y^{x}\left(\log y + \frac{x}{y} \frac{dy}{dx}\right) = 0$$
$$\Rightarrow \left(x^{2} + \log x + xy^{y-1}\right) \frac{dy}{dx} = -\left(yx^{y-1} + y^{x}\log y\right)$$
Hence,
$$\frac{dy}{dx} = \frac{yx^{y-1} + y^{x}\log y}{x^{y}\log x + xy^{x-1}}.$$

13. Determine $\frac{dy}{dx}$ from the equation $y^x = x^y$.

Ans: The given equation is $y^x = x^y$.

Then, taking logarithm both sides of the equation give xlogy=ylogx.

$$logy \times \frac{d}{dx}(x) + x \times \frac{d}{dx}(logy) = logx \times \frac{d}{dx}(y) + y \times \frac{d}{dx}(logx)$$
$$\Rightarrow logy \times 1 + x \times \frac{1}{y} \times \frac{dy}{dx} = logx \times \frac{dy}{dx} + y \times \frac{1}{x}$$
$$\Rightarrow logy + \frac{x}{y} \frac{dy}{dx} = logx \frac{dy}{dx} + \frac{y}{x}$$
$$\Rightarrow \left(\frac{x}{y} - logx\right) \frac{dy}{dx} = \frac{y}{x} - logy$$

$$\Rightarrow \left(\frac{x \cdot y \log x}{y}\right) \frac{dy}{dx} = \frac{y \cdot x \log y}{x}$$
$$\Rightarrow \left(\frac{x \cdot y \log x}{y}\right) \frac{dy}{dx} = \frac{y \cdot x \log y}{x}$$
Therefore, $\frac{dy}{dx} = \frac{y}{x} \left(\frac{y \cdot x \log y}{x \cdot y \log x}\right)$.

14. Determine $\frac{dy}{dx}$ from the equation $(\cos x)^{y} = (\cos y)^{x}$.

Ans: The given equation is $(\cos x)^y = (\cos y)^x$.

Then, taking logarithm both sides of the equation give

ylogcosx=xlogcosy.

Now, differentiating both sides of the equation with respect to x gives

$$logcosx \times \frac{dy}{dx} + y \times \frac{d}{dx} (logcosx) = logcosy \times \frac{d}{dx} (x) + x \times \frac{d}{dx} (logcosy)$$

$$\Rightarrow logcosx \frac{dy}{dx} + \frac{y}{cosx} \times (-sinx) = logcosy + \frac{x}{cosy} (-siny) \times \frac{dy}{dx}$$

$$\Rightarrow logcosx \frac{dy}{dx} - ytanx = logcosy - xtany \frac{dy}{dx}$$

$$\Rightarrow (logcosx + xtany) \frac{dy}{dx} = ytanx + logcosy$$

Therefore, $\frac{dy}{dx} = \frac{ytanx + logcosy}{xtany + logcosx}$.

15. Determine
$$\frac{dy}{dx}$$
 from the equation $xy=e^{(x-y)}$.

Ans: The given equation is $xy=e^{(x-y)}$.

Then, taking logarithm both sides of the equation give

$$log(xy)=log(e^{x-y})$$

$$\Rightarrow logx+logy=(x-y)loge$$

$$\Rightarrow logx+logy=(x-y)\times 1$$

$$\Rightarrow logx+logy=x-y$$

Now, differentiating both sides of the equation with respect to x gives

$$\frac{d}{dx}(\log x) + \frac{d}{dx}(\log y) = \frac{d}{dx}(x) - \frac{dy}{dx}$$
$$\Rightarrow \frac{1}{x} + \frac{1}{y}\frac{dy}{dx} = 1 - \frac{1}{x}$$
$$\Rightarrow \left(1 + \frac{1}{y}\right)\frac{dy}{dx} = \frac{x - 1}{x}$$

- Therefore, $\frac{dy}{dx} = \frac{y(x-1)}{x(x+1)}$.
- 16. Determine the derivative of the following function f and hence evaluate f'(1).

 $f(x)=(1+x)(1+x^2)(1+x^4)(1+x^8)$.

Ans: The given function is $f(x)=(1+x)(1+x^2)(1+x^4)(1+x^8)$.

By taking logarithm both sides of the equation give

 $logf(x) = log(1+x) + log(1+x^{2}) + log(1+x^{4}) + log(1+x^{8})$

$$\frac{1}{f(x)} \times \frac{d}{dx} [f(x)] = \frac{d}{dx} \log(1+x) + \frac{d}{dx} \log(1+x^2) + \frac{d}{dx} \log(1+x^4) + \frac{d}{dx} \log(1+x^8)$$

$$\Rightarrow \frac{1}{f(x)} \times f'(x) = \frac{1}{1+x} \times \frac{1}{dx} (1+x) + \frac{1}{1+x^2} \times \frac{d}{dx} \log(1+x^2) + \frac{1}{1+x^4} \times \frac{d}{dx} \log(1+x^4) + \frac{1}{1+x^8} \times \frac{d}{dx} \log(1+x^8)$$
$$\Rightarrow f'(x) = f(x) \left[\frac{1}{1+x} + \frac{1}{1+x^2} \times 2x + \frac{1}{1+x^4} \times 4x^3 + \frac{1}{1+x^8} \times 8x^7 \right]$$

Therefore,

$$f'(x) = (1+x)(1+x^{2})(1+x^{4})(1+x^{8}) \left[\frac{1}{1+x} + \frac{2x}{1+x^{2}} + \frac{4x^{3}}{1+x^{4}} + \frac{8x^{7}}{1+x^{8}}\right]$$

So,

$$f'(1) = (1+1)(1+1^{2})(1+1^{4})(1+1^{8}) \left[\frac{1}{1+1} + \frac{2 \times 1}{1+1^{2}} + \frac{4 \times 1^{3}}{1+1^{4}} + \frac{8 \times 1^{7}}{1+1^{8}} \right]$$
$$= 2 \times 2 \times 2 \times 2 \left[\frac{1}{2} + \frac{2}{2} + \frac{4}{2} + \frac{8}{2} \right]$$
$$= 16 \times \left(\frac{1+2+4+8}{2} \right)$$
$$= 16 \times \frac{15}{2} = 120$$

Hence, f'(1) = 120.

- 17. Differentiate the function y=(x²-5x+8)(x³+7x+9) in three ways as described below. Also, verify whether all the answers are the same.
 (a) By using product rules.
- Ans: The given function is $y=(x^2-5x+8)(x^3+7x+9)$.

Now, let consider $u=(x^2-5x+8)$ and $v=(x^3+7x+9)$

Therefore, y=uv.

$$\Rightarrow \frac{dy}{dx} = \frac{du}{dv} \cdot v + u \cdot \frac{du}{dx}$$

$$\Rightarrow \frac{dy}{dx} = \frac{d}{dx} (x^2 - 5x + 8) \cdot (x^3 + 7x + 9) + (x^2 - 5x + 8) \cdot \frac{d}{dx} (x^3 + 7x + 9)$$

$$\Rightarrow \frac{dy}{dx} = (2x - 5)(x^3 + 7x + 9) \cdot (x^2 - 5x + 8)(3x^2 + 7)$$

$$\Rightarrow \frac{dy}{dx} = 2x(x^3 + 7x + 9) - 5(x^2 - 5x + 8) + x^2(3x^2 + 7) - 5x(3x^2 + 7) - 8(3x^2 + 7)$$

$$\Rightarrow \frac{dy}{dx} = (2x^4 + 14x^2 + 18x) - 5x^3 - 35x - 45 + (3x^4 + 7x^2) - 15x^3 - 35x + 24x^2 + 56$$

Hence, $\frac{dy}{dx} = 5x^4 - 20x^3 + 45x^2 + 52x + 11$.

(b) By expanding the factors as a polynomial.

Ans: The given function is

$$y=(x^2-5x+8)(x^3+7x+9)$$
.

Then, calculating the product, gives

$$y=x^{2}(x^{3}+7x+9)-5x^{4}(x^{3}+7x+9)+8(x^{3}+7x+9)$$

$$\Rightarrow y=x^{5}+7x^{3}+9x^{2}-5x^{3}-26x^{2}+11x+72$$

$$\frac{dy}{dx} = \frac{d}{dx} (x^5 + 7x^3 + 9x^2 - 5x^3 - 26x^2 + 11x + 72)$$

= $\frac{d}{dx} (x^5) - 5\frac{d}{dx} (x^4) + 15\frac{d}{dx} (x^3) - 26\frac{d}{dx} (x^3) + 11\frac{d}{dx} (x) + \frac{d}{dx} (72)$
= $5x^4 - 5 \times 4x^3 + 15 \times 3x^2 - 26 \times 2x + 11 \times 1 + 0$
Hence, $\frac{dy}{dx} = 5x^4 - 20x^3 + 45x^2 - 52x + 11$.

(c) By using a logarithmic function.

Ans: The given function is

 $y=(x^2-5x+8)(x^3+7x+9)$.

Now, taking logarithm both sides of the function give

 $\log = \log(x^2 - 5x + 8) + \log(x^3 + 7x + 9)$

Differentiating both sides of the equation with respect to x gives

$$\frac{1}{y}\frac{dy}{dx} = \frac{d}{dx}\log(x^2-5x+8) + \frac{d}{dx}\log(x^3+7x+9)$$

$$\Rightarrow \frac{1}{y}\frac{dy}{dx} = \frac{1}{x^2-5x+8} \cdot \frac{d}{dx}(x^2-5x+8) + \frac{1}{x^3+7x+9} \cdot \frac{d}{dx}(x^3+7x+9)$$

$$\Rightarrow \frac{dy}{dx} = y\left[\frac{1}{x^2-5x+8} \times (2x-5) + \frac{1}{x^3+7x+9} \times (3x^2+7)\right]$$

$$\Rightarrow \frac{dy}{dx} = (x^2-5x+8)(x^3+7x+9)\left[\frac{2x-5}{x^3-5x+8} + \frac{3x^2+7}{x^3+7x+9}\right]$$

$$\Rightarrow \frac{dy}{dx} = (x^2-5x+8)(x^3+7x+9)\left[\frac{(2x-5)(x^3+7x+9)+(3x^2+7)(x^2-5x+8)}{(x^3-5x+8)+(x^3+7x+9)}\right]$$

$$\Rightarrow \frac{dy}{dx} = 2x(x^3+7x+9x^2)-5(x^3+7x+9)+3x^2(x^2-5x+8)+7(x^3+7x+9)$$

$$\Rightarrow \frac{dy}{dx} = (2x^4+14x^2+18x)+(5x^3-35x+45)+(3x^4-15x^3+24x^2)+(7x^2+35x+56)$$
Therefore, $\frac{dy}{dx} = 5x^2-20x^3+45x^2-52x+11$.

Hence, comparing the above three results, it is concluded that the derivative $\frac{dy}{dx}$ are the same for all methods.

18. Let u, v, and w are functions of x , then prove that

 $\frac{d}{dx}(u.v.w) = \frac{du}{dx}v.w+u\frac{du}{dx}.w+u.v\frac{dw}{dx}$ in two ways. First by using repeated application of product rule and second by applying logarithmic differentiation.

Ans: Let the function y=u.v.w=u.(v.w).

Then applying the product rule of derivatives, give

$$\frac{dy}{dx} = \frac{du}{dx} \cdot (v.w) + u \cdot \frac{d}{dx} (v.w)$$
$$\Rightarrow \frac{dy}{dx} = \frac{du}{dx} \cdot v.w + u \left[\frac{dv}{dx} \cdot w + v \cdot \frac{dv}{dx} \right] \qquad \text{(Using the product rule again)}$$

Thus,

 $\frac{\mathrm{d} y}{\mathrm{d} x} = \frac{\mathrm{d} u}{\mathrm{d} x} v.w + u.\frac{\mathrm{d} v}{\mathrm{d} x}.w + u.v\frac{\mathrm{d} w}{\mathrm{d} x}.$

Now, take logarithm both sides of the function y=u.v.w.

Then, we have logy=logu+logv+logw.

$$\frac{1}{y} \cdot \frac{dy}{dx} = \frac{d}{dx} (\log u) + \frac{d}{dx} (\log v) + \frac{d}{dx} (\log w)$$
$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = \frac{1}{u} \frac{du}{dx} + \frac{1}{v} \frac{dv}{dx} + \frac{1}{w} \frac{dw}{dx}$$
$$\Rightarrow \frac{dy}{dx} = y \left(\frac{1}{u} \frac{du}{dx} + \frac{1}{v} \frac{dv}{dx} + \frac{1}{w} \frac{dw}{dx} \right)$$
$$\Rightarrow \frac{dy}{dx} = u \cdot v \cdot w \left(\frac{1}{u} \frac{du}{dx} + \frac{1}{v} \frac{dv}{dx} + \frac{1}{w} \frac{dw}{dx} \right)$$
Hence, $\frac{dy}{dx} = \frac{du}{dx} v \cdot w + u \frac{du}{dx} \cdot w + u \cdot v \frac{dw}{dx}$.

Exercise 5.6

1. Determine $\frac{dy}{dx}$ from the equations x=2at², y=at⁴, without eliminating the parameter t, where a,b are constants.

Ans: The given equations are

$$x=2at^{2}$$
 (1)

and
$$y=at^4$$
(2)

Then, differentiating both sides of the equation (1) with respect to t gives

$$\frac{\mathrm{d}x}{\mathrm{d}t} = \frac{\mathrm{d}}{\mathrm{d}t}(2\mathrm{a}t^2) = 2\mathrm{a} \times \frac{\mathrm{d}}{\mathrm{d}t}(t^2) = 2\mathrm{a} \times 2\mathrm{t} = 4\mathrm{a}\mathrm{t}. \qquad \dots (3)$$

Also, differentiating both sides of the equation (2) with respect to t gives

$$\frac{dy}{dt} = \frac{d}{dt}(at^4) = a \times \frac{d}{dt}(t^4) = a \times 4 \times t^3 = 4at^3 \qquad \dots \dots (4)$$

Now, dividing the equations (4) by (3) gives

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{4at^3}{4at} = t^2.$$

Hence, $\frac{dy}{dx} = t^2.$

- 2. Determine $\frac{dy}{dx}$ from the equations x=acos θ , y=bcos θ , without eliminating the parameter θ , where a,b are constants.
- **Ans:** The given equations are

 $x = a \cos \theta$ (1)

and
$$y=b\cos\theta$$
(2)

Then, differentiating both sides of the equation (1) with respect to θ gives

$$\frac{\mathrm{dx}}{\mathrm{d\theta}} = \frac{\mathrm{d}}{\mathrm{d\theta}}(\mathrm{acos}\theta) = \mathrm{a}(-\mathrm{sin}\theta) = -\mathrm{asin}\theta \,. \tag{3}$$

Also, differentiating both sides of the equation (1) with respect to θ gives

$$\frac{dy}{d\theta} = \frac{d}{d\theta} (b\cos\theta) = b(-\sin\theta) = -b\sin\theta \qquad \dots \dots (4)$$

Therefore, dividing the equation (4) by (3) gives

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} = \frac{-b\sin\theta}{-a\sin\theta} = \frac{b}{a}.$$

Hence, $\frac{dy}{dx} = \frac{b}{a}.$

3. Determine $\frac{dy}{dx}$ from the equations x=sint, y=cos2t without eliminating the parameter t.

Ans: The given equations are

x=sint	(1)
and y=cos2t	(2)

Then, differentiating both sides of the equation (1) with respect to t gives

$$\frac{\mathrm{dx}}{\mathrm{dt}} = \frac{\mathrm{d}}{\mathrm{dt}}(\mathrm{sint}) = \mathrm{cost} \;. \tag{3}$$

Also, differentiating both sides of the equation (2) with respect to t gives

$$\frac{dy}{dt} = \frac{d}{dt}(\cos 2t) = \sin 2t \times \frac{d}{dt}(2t) = -2\sin 2t \qquad \dots \dots (4)$$

Therefore, by dividing the equation (4) by (3) gives

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{-2\sin 2t}{\cos t} = \frac{-2 \times 2\sin t \cos t}{\cos t} = -4\sin t$$

Hence, $\frac{dy}{dx} = -4\sin t$.

- 4. Determine $\frac{dy}{dx}$ from the equations x=4t, y= $\frac{4}{t}$ without eliminating the parameter t.
- Ans: The given equations are

x=4t (1)
and
$$y=\frac{4}{t}$$
 (2)

$$\frac{\mathrm{dx}}{\mathrm{dt}} = \frac{\mathrm{d}}{\mathrm{dt}}(4t) = 4. \qquad \dots \dots (3)$$

Also, differentiating both sides of the equation (2) with respect to t gives

$$\frac{dy}{dt} = \frac{d}{dt} \left(\frac{4}{t}\right) = 4 \times \frac{d}{dt} \left(\frac{1}{t}\right) = 4 \times \left(\frac{-1}{t^2}\right) = \frac{-4}{t^2} \qquad \dots \dots (4)$$

Therefore, dividing the equation (4) by (3) gives

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{\left(\frac{-4}{t^2}\right)}{4} = \frac{-1}{t^2}.$$

Hence, $\frac{dy}{dx} = -\frac{1}{t^2}.$

5. Determine $\frac{dy}{dx}$ from the equations x=cos θ -cos 2θ , y=sin θ -sin 2θ , without eliminating the parameter θ .

Ans: The given equations are

$$x = \cos\theta - \cos 2\theta$$
(1)

and
$$y=\sin\theta-\sin2\theta$$
(2)

Then, differentiating both sides of the equation (1) with respect to θ gives

$$\frac{dx}{d\theta} = \frac{d}{d\theta}(\cos\theta - \cos2\theta) = \frac{d}{d\theta}(\cos\theta) - \frac{d}{d\theta}(\cos2\theta) = -\sin\theta(-2\sin2\theta) = 2\sin2\theta - \sin\theta \dots (3)$$

Also, differentiating both sides of the equation (2) with respect to θ gives

$$\frac{dy}{d\theta} = \frac{d}{d\theta} (\sin\theta - \sin2\theta) = \frac{d}{d\theta} (\sin\theta) - \frac{d}{d\theta} (\sin2\theta) = \cos\theta - 2\cos\theta \qquad \dots (4)$$

Therefore, dividing the equation (4) by (3) gives

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} = \frac{\cos\theta - 2\cos\theta}{2\sin2\theta - \sin\theta}.$$

Hence, $\frac{dy}{dx} = \frac{\cos\theta - 2\cos\theta}{2\sin2\theta - \sin\theta}.$

6. Determine $\frac{dy}{dx}$ from the equations $x=a(\theta-\sin\theta)$, $y=a(1+\cos\theta)$, without eliminating the parameter θ , where a, b are constants.

Ans: The given equations are

$$x=a(\theta-\sin\theta)$$
(1)

and
$$y=a(1+\cos\theta)$$
(2)

Then, differentiating both sides of the equation (1) with respect to θ gives

$$\frac{\mathrm{d}x}{\mathrm{d}\theta} = a \left[\frac{\mathrm{d}}{\mathrm{d}\theta}(\theta) - \frac{\mathrm{d}}{\mathrm{d}\theta}(\sin\theta) \right] = a(1 - \cos\theta) \qquad \dots \dots (3)$$

Also, differentiating both sides of the equation (2) with respect to θ gives

$$\frac{dy}{d\theta} = a \left[\frac{d}{d\theta} (1) + \frac{d}{d\theta} (\cos\theta) \right] = a \left[0 + (-\sin\theta) \right] = -a\sin\theta \qquad \dots \dots (4)$$

Therefore, by dividing the equation (4) by (3) gives

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} = \frac{-a\sin\theta}{a(1-\cos\theta)} = \frac{-2\sin\frac{\theta}{2}\cos\frac{\theta}{2}}{2\sin^2\frac{\theta}{2}} = \frac{-\cos\frac{\theta}{2}}{\sin\frac{\theta}{2}} = -\cos\frac{\theta}{2}.$$

Hence, $\frac{dy}{dx} = -\cos\frac{\theta}{2}.$

7. Determine $\frac{dy}{dx}$ from the equations $x=-\frac{\sin^3 t}{\sqrt{\cos 2t}}$, $y=\frac{\cos^3 t}{\sqrt{\cos 2t}}$, without eliminating the parameter t.

Ans: The given equations are,

$$x = -\frac{\sin^{3}t}{\sqrt{\cos 2t}} \qquad \dots \dots (1)$$

and
$$y = \frac{\cos^{3}t}{\sqrt{\cos 2t}} \qquad \dots \dots (2)$$

$$\frac{dx}{dt} = \frac{d}{dt} \left[\frac{\sin^3 t}{\sqrt{\cos 2t}} \right]$$
$$= \frac{\sqrt{\cos 2t} - \frac{d}{dt} (\sin^3 t) - \sin^3 t \times \frac{d}{dt} \sqrt{\cos 2t}}{\cos 2t}$$

$$=\frac{\sqrt{\cos 2t} \times 3\sin^2 t \times \frac{d}{dt}(\sin t) - \sin^3 t \times \frac{1}{2\sqrt{\cos 2t}} \times \frac{d}{dt}(\cos 2t)}{\cos 2t}$$
$$=\frac{3\sqrt{\cos 2t} \times \sin^2 t \cos t - \frac{\sin^3 t}{2\sqrt{\cos 2t}} \times (-2\sin 2t)}{\cos 2t\sqrt{\cos 2t}}$$

$$\frac{dx}{dt} = \frac{3\cos 2t\sin^2 t\cos t + \sin^2 t\sin 2t}{\cos 2t \sqrt{\cos 2t}}.$$
(3)

$$\frac{dy}{dt} = \frac{d}{dt} \left[\frac{\cos^3 t}{\sqrt{\cos 2t}} \right]$$

$$= \frac{\sqrt{\cos 2t} \times \frac{d}{dt} (\cos^3 t) - \cos^3 t \times \frac{d}{dt} (\sqrt{\cos 2t})}{\cos 2t}$$

$$= \frac{3\sqrt{\cos 2t} \cos^2 t (-\sin t) - \cos^3 t \times \frac{1}{2(\sqrt{\cos 2t})} \times \frac{d}{dt} (\cos 2t)}{\cos 2t}$$

$$\frac{dy}{dt} = \frac{-3\cos 2t \times \cos^2 t \times \sin t + \cos^3 t \sin 2t}{\cos 2t} \qquad \dots (4)$$

Thus, dividing the equation (4) by the equation (3) gives

$$\frac{dy}{dx} = \frac{\left(\frac{dx}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{-3\cos 2t \times \cos^2 t \times \sin t + \cos^3 t \sin 2t}{3\cos 2t \cosh^2 t + \sin^3 t \sin 2t}$$
$$= \frac{\sin t \cot \left[-3\cos 2t \times \cos t + 2\cos^3 t\right]}{\sin t \cot \left[3\cos 2t \sin t + 2\sin^3 t\right]}$$
$$= \frac{\left[-3(2\cos^2 t - 1)\cos t + 2\cos^3 t\right]}{\left[3(1 - 2\sin^3 t)\sin t + 2\sin^3 t\right]} \qquad \left[\cos 2t = (2\cos^2 t - 1)\right]$$

$$=\frac{-4\cos^{3}t+3\cos t}{3\sin t-4\sin^{3}t}$$

$$=\frac{-\cos 3t}{\sin 3t}$$

$$\begin{bmatrix}\cos 3t=4\cos^{3}t-3\cos t\\\sin 3t=3\sin t-4\sin^{2}t\end{bmatrix}$$

Hence,
$$\frac{dy}{dx} = -\cot 3t$$
.

8. Determine $\frac{dy}{dx}$ from the parametric equations

$$x=a\left(cost+logtanrac{t}{2}\right)$$
, y=asint , without eliminating the parameter t .

Ans: The given equations are

$$x=a\left(\cosh + \log \tan \frac{t}{2}\right) \qquad \dots \dots (1)$$

and y=asint
$$\dots \dots (2)$$

$$\frac{dx}{dt} = a \times \left[\frac{d}{d\theta} (\cos t) + \frac{d}{d\theta} (\log tan \frac{t}{2}) \right]$$
$$= a \left[-\sin t + \frac{1}{\tan \frac{t}{2}} \times \frac{d}{dt} \left(\tan \frac{t}{2} \right) \right]$$
$$= a \left[-\sin t + \cot \frac{t}{2} \times \sec^2 \frac{t}{2} \times \frac{d}{dt} \left(\frac{t}{2} \right) \right]$$
$$= a \left(-\sin t + \frac{\cos \frac{t}{2}}{\sin \frac{t}{2}} \times \frac{1}{\cos^2 \frac{t}{2}} \times \frac{1}{2} \right)$$

$$=a\left(-\sin t + \frac{1}{2\sin \frac{t}{2}\cos \frac{t}{2}}\right)$$
$$=a\left(-\sin t + \frac{1}{\sin t}\right)$$
$$=a\left(\frac{-\sin^{2}t + 1}{\sin t}\right)$$
Therefore, $\frac{dx}{dt} = a\frac{\cos^{2}t}{\sin t}$ (3)

$$\frac{dy}{dt} = a \frac{d}{dt} (sint) = acost \qquad \dots \dots (4)$$

Thus, dividing the equation (4) by the equation (3) gives

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{acost}{\left(a\frac{cos^2t}{sint}\right)} = \frac{sint}{cost} = tant$$

Hence, $\frac{dy}{dx} = tant$.

9. Determine $\frac{dy}{dx}$ from the parametric equations x=asec θ , y=btan θ , without eliminating the parameter θ , where a,b are constants.

Ans: The given equations are

$$x=asec$$
...... (1)and $y=btan\theta$ (2)

Then, differentiating both sides of the equation (1) with respect to θ gives

$$\frac{dx}{d\theta} = a \times \frac{d}{d\theta} (\sec\theta) = \operatorname{asec}\theta \tan\theta \qquad \dots \dots (3)$$

Also, differentiating both sides of the equation (2) with respect to θ gives

$$\frac{dy}{d\theta} = b \times \frac{d}{d\theta} (\tan \theta) = b \sec^2 \theta \qquad \dots \dots (4)$$

Thus, dividing the equation (4) by the equation (3) gives

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} = \frac{b\sec^2\theta}{a\sec\theta\tan\theta} = \frac{b}{a}\sec\theta\tan\theta = -\frac{b\cos\theta}{a\cos\theta\sin\theta} = \frac{b}{a} \times \frac{1}{\sin\theta} = \frac{b}{a}\csc\theta$$
Hence, $\frac{dy}{dx} = \frac{b}{a}\csc\theta$.

10. Determine $\frac{dy}{dx}$ from the parametric equations

x=a(cos θ + θ sin θ), y=a(sin θ - θ cos θ), without eliminating the parameter θ , where a,b are constants.

Ans: The given equations are

$$x=a(\cos\theta+\theta\sin\theta) \qquad \qquad \dots \dots (1)$$

and $y=a(\sin\theta-\theta\cos\theta)$ (2)

Then, differentiating both sides of the equation (1) with respect to θ gives

$$\frac{dx}{d\theta} = a \left[\frac{d}{d\theta} \cos\theta + \frac{d}{d\theta} (\theta \sin\theta) \right] = a \left[-\sin\theta + \theta \frac{d}{d\theta} (\sin\theta) + \sin\theta \frac{d}{d\theta} (\theta) \right]$$
$$= a \left[-\sin\theta + \theta \cos\theta + \sin\theta \right].$$
Therefore, $\frac{dx}{d\theta} = a\theta \cos\theta$. (2)

Therefore,
$$\frac{dx}{d\theta} = a\theta\cos\theta$$
(3)

Also, differentiating both sides of the equation (2) with respect to θ gives

$$\frac{dy}{d\theta} = a \left[\frac{d}{d\theta} (\sin\theta) - \frac{d}{d\theta} (\theta \cos\theta) \right] = a \left[\cos\theta - \left\{ \theta \frac{d}{d\theta} (\cos\theta) + \cos\theta \times \frac{d}{d\theta} (\theta) \right\} \right]$$
$$\Rightarrow \frac{dy}{d\theta} = a \left[\cos\theta + \theta \sin\theta - \cos\theta \right]$$
Therefore, $\frac{dy}{d\theta} = a\theta \sin\theta$

Thus, dividing the equation (4) by the equation (3) gives

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} = \frac{a\theta\sin\theta}{a\theta\sin\theta} = \tan\theta$$

Hence,
$$\frac{dy}{dx} = tan\theta$$
.

11. Prove that $\frac{dy}{dx} = \frac{y}{x}$, where it is provided that $x = \sqrt{a^{\sin^{-1}t}}$, $y = \sqrt{a^{\cos^{-1}t}}$.

Ans: The given parametric equations are $x=\sqrt{a^{\sin^{-1}t}}$ and $y=\sqrt{a^{\cos^{-1}t}}$.

Now,
$$x = \sqrt{a^{\sin^{-1}t}}$$
 and $y = \sqrt{a^{\cos^{-1}t}}$
 $\Rightarrow x = (a^{\sin^{-1}t})$ and $y = (a^{\cos^{-1}t})^{\frac{1}{2}}$
 $\Rightarrow x = a^{\frac{1}{2}\sin^{-1t}}$ and $y = a^{\frac{1}{2}\cos^{-1}t}$

Therefore, first consider $x=a^{\frac{1}{2}sin^{-1}t}$.

Take logarithms on both sides of the equation.

Then, we have

$$\log x = \frac{1}{2} \sin^{-1} t \log a$$
.

$$\frac{1}{x} \times \frac{dx}{dt} = \frac{1}{2} \log a \times \frac{d}{dt} (\sin^{-1}t)$$

$$\Rightarrow \frac{dx}{dt} = \frac{x}{2} \log a \times \frac{1}{\sqrt{1-t^2}}$$

Therefore, $\frac{dx}{dt} = \frac{x \log a}{2\sqrt{1-t^2}}$(1)

$$\frac{1}{\cos^{-1}t}$$

Again, consider the equation $y=a^{\frac{1}{2}\cos^{4}t}$.

Take logarithm both sides of the equation.

Then, we have

$$\log y = \frac{1}{2} \cos^{-1} t \log a$$

Differentiating both sides of the equation with respect to t gives

$$\frac{1}{y} \times \frac{dx}{dt} = \frac{1}{2} \log a \times \frac{d}{dt} (\cos^{-1}t)$$

$$\Rightarrow \frac{dx}{dt} = \frac{y \log a}{2} \times \left(\frac{1}{\sqrt{1 - t^2}}\right)$$

Therefore, $\frac{dx}{dt} = \frac{-y \log a}{2\sqrt{1 - t^2}}$(2)

Thus, dividing the equation (2) by the equation (1) gives

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\left(\frac{\mathrm{d}y}{\mathrm{d}t}\right)}{\left(\frac{\mathrm{d}x}{\mathrm{d}t}\right)} = \frac{\left(\frac{-\mathrm{yloga}}{2\sqrt{1-t^2}}\right)}{\left(\frac{\mathrm{xloga}}{2\sqrt{1-t^2}}\right)} = \frac{\mathrm{y}}{\mathrm{x}}.$$

Hence,
$$\frac{dy}{dx} = \frac{y}{x}$$
.

Exercise 5.7

- 1. Determine the second order derivative for the following function $y=x^2+3x+2$.
- **Ans:** The given function is $y=x^2+3x+2$.

Then, differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{d}{dx}(x^2) + \frac{d}{dx}(3x) + \frac{d}{dx}(2) = 2x + 3 + 0 = 2x + 3$$

That is,

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 2x + 3.$$

Again, differentiating both sides with respect to x gives

$$\frac{d^2y}{dx^2} = \frac{d}{dx}(2x+3) = \frac{d}{dx}(2x) + \frac{d}{dx}(3) = 2 + 0 = 2$$

Hence, $\frac{d^2y}{dx^2} = 2$.

2. Determine the second order derivative for the following function $y = x^{20}$.

Ans: The given function is $y=x^{20}$.

Then, differentiating both sides with respect to x gives

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}}{\mathrm{d}x}(x^{20}) = 20x^{19}$$

Again, differentiating both sides with respect to x gives

$$\frac{d^2y}{dx^2} = \frac{d}{dx} (20x^{19}) = 20 \frac{d}{dx} (x^{19}) = 20(19)x^{18} = 380x^{18}.$$

Hence, $\frac{d^2y}{dx^2} = 380x^{18}.$

- 3. Determine the second order derivative for the following function $y = x \cdot \cos x$.
- **Ans:** The given function is y=x.cosx.

Then, differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{d}{dx}(x.\cos x) = \cos x \cdot \frac{d}{dx}(x) + x \frac{d}{dx}(\cos x) = \cos x \cdot 1 + x(-\sin x) = \cos x \cdot x \sin x$$

That is, $\frac{dy}{dx} = \cos x \cdot x \sin x$.

Again, differentiating both sides with respect to x gives

$$\frac{d^2y}{dx^2} = \frac{d}{dx}(\cos x - x\sin x) = \frac{d}{dx}(\cos x) - \frac{d}{dx}(x\sin x)$$
$$= -\sin x - [\sin x \cdot \frac{d}{dx}(x) + x \cdot \frac{d}{dx}(\sin x)$$
$$= -\sin x - (\sin x + \cos x)$$
Hence,
$$\frac{d^2y}{dx^2} = -(x\cos x + 2\sin x)$$
.

4. Determine the second order derivative for the following function $y = \log x$

Ans: The given function is y=logx.

$$\frac{dy}{dx} = \frac{d}{dx}(\log x) = \frac{1}{x}$$

Again, differentiating both sides with respect to x gives

$$\frac{d^2y}{dx^2} = \frac{d}{dx} \left(\frac{1}{x}\right) = \frac{-1}{x^2}$$

Hence, $\frac{d^2y}{dx^2} = -\frac{1}{x^2}$.

5. Determine the second order deriva tive for the following function $y=x^{3}\log x$

Ans: The given function is $y=x^{3}\log x$.

Then, differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{d}{dx} \left[x^3 \log x \right] = \log x \cdot \frac{d}{dx} (x^3) + x^3 \frac{d}{dx} (\log x) = \log x \cdot 3x^2 + x^3 \cdot \frac{1}{x} = \log x \cdot 3x^2 + x^2$$

That is, $\frac{dy}{dx} = x^2 (1 + 3\log x)$.

Again, differentiating both sides with respect to x gives

$$\frac{d^{2}y}{dx^{2}} = \frac{d}{dx}(x^{2}(1+3\log x))$$

=(1+3logx). $\frac{d}{dx}(x^{2})+x^{2}\frac{d}{dx}(1+3\log x)$
=(1+3logx).2x+x³. $\frac{3}{x}$
=2x+6logx+3x
=5x+6xlogx
 $d^{2}x$

Hence,
$$\frac{d^2y}{dx^2} = x(5+6\log x)$$
.

6. Determine the second order derivative for the following function. $y = e^x \sin 5x$

Ans: The given function is $y=e^x \sin 5x$.

Then, differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{d}{dx} \left[e^x \sin 5x \right] = \sin x \frac{d}{dx} (e^x) + e^x \frac{d}{dx} (\sin 5x)$$
$$\Rightarrow \frac{dy}{dx} = \sin 5x \cdot e^x + e^x \cdot \cos 5x \cdot \frac{d}{dx} (5x)$$

That is, $\frac{dy}{dx} = e^x(\sin 5x + 5\cos 5x)$.

Again, differentiating both sides with respect to x gives

$$\frac{d^{2}y}{dx^{2}} = \frac{d}{dx} \Big[e^{x} (\sin 5x + 5\cos 5x) \Big]$$

=(\sin5x+5\cos5x).\frac{d}{dx} (e^{x}) + e^{x}.\frac{d}{dx} (\sin5x+5\cos5x) = (\sin5x+5\cos5x)(e^{x}) + e^{x} \Big[\cos 5x.\frac{d}{dx} (5x) + 5(-\sin5x).\frac{d}{dx} (5x) \Big]
=e^{x} (\sin5x+5\cos5x)(e^{x}) + e^{x} (5\cos5x-25\sin5x) = e^{x} (10\cos5x-24\sin5x).
Hence, \frac{d^{2}y}{dx^{2}} = 2e^{x} (5\cos5x-12\sin5x).

7. Determine the second order derivative for the following function. $y=e^{6x}\cos 3x$.

Ans: The given function is $y=e^{6x}\cos 3x$.

$$\frac{dy}{dx} = \frac{d}{dx} (e^{6x} \cos 3x) = \cos 3x \times \frac{d}{dx} (e^{6x}) + e^{6x} \times \frac{d}{dx} (\cos 3x)$$
$$\Rightarrow \frac{dy}{dx} = \cos 3x \times e^{6x} \times \frac{d}{dx} (6x) + e^{6x} \times (-\sin 3x) \times \frac{d}{dx} (3x)$$

Therefore,

$$\frac{\mathrm{dy}}{\mathrm{dx}} = 6\mathrm{e}^{6\mathrm{x}}\cos 3\mathrm{x} \cdot 3\mathrm{e}^{6\mathrm{x}}\sin 3\mathrm{x} \qquad \dots \dots (1)$$

Again, differentiating both sides with respect to x gives

$$\frac{d^{2}y}{dx^{2}} = \frac{d}{dx} (6e^{6x}\cos 3x - 3e^{6x}\sin 3x) = 6 \times \frac{d}{dx} (e^{6x}\cos 3x) - 3 \times \frac{d}{dx} (e^{6x}\sin 3x)$$

$$= 6 \times \left[6e^{6x}\cos 3x - 3e^{6x}\sin 3x \right] - 3 \times \left[\sin 3x \times \frac{d}{dx} (e^{6x}) + e^{6x} \times \frac{d}{dx} (\sin 3x) \right] [using (1)]$$

$$= 36e^{6x}\cos 3x - 18e^{6x}\sin 3x - 3\left[\sin 3x \times e^{6x} \times 6 + e^{6x} \times \cos 3x - 3 \right]$$

$$= 36e^{6x}\cos 3x - 18e^{6x}\sin 3x - 18e^{6x}\sin 3x - 9e^{6x}\cos 3x$$
Hence, $\frac{d^{2}y}{dx^{2}} = 9e^{6x} (3\cos 3x - 4\sin 3x).$

8. Determine the second order derivative for the following function. $y = tan^{-1}x$.

Ans: The given function is $y=\tan^{-1}x$.

Then, differentiating both sides with respect to x gives

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}}{\mathrm{d}x} \tan^{-1}x = \frac{1}{1 - x^2}$$

Again, differentiating both sides with respect to x gives

$$\frac{d^{2}y}{dx^{2}} = \frac{d}{dx} \left(\frac{1}{1+x^{2}} \right) = \frac{d}{dx} (1+x^{2}) = (-1) \times (1+x^{2}) \times \frac{d}{dx} (1+x^{2})$$
$$= \frac{1}{(1+x^{2})} \times 2x$$

Hence, $\frac{d^2 y}{dx^2} = \frac{-2x}{(1+x^2)}$.

9. Determine the second order derivative for the following function. y=log(logx).

Ans: The given function is y=log(logx).

Now, differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{d}{dx} [\log(\log x)]$$
$$\Rightarrow \frac{1}{\log x} \times \frac{d}{dx} (\log x)$$
$$\Rightarrow \frac{1}{\log x} = (x \log x)^{-1}$$

Again, differentiating both sides with respect to x gives

$$\frac{d^2 y}{dx^2} = \frac{d}{dx} \Big[(x \log x)^{-1} \Big] = (-1) \times (x \log x)^{-2} \frac{d}{dx} (x \log x)$$
$$= \frac{-1}{(x \log x)^2} \times \Big[\log x \times \frac{d}{dx} (x) + x \times \frac{d}{dx} (\log x) \Big]$$
$$= \frac{-1}{(x \log x)^2} \times \Big[\log x \times 1x \times \frac{1}{x} \Big]$$
Hence,
$$\frac{d^2 y}{dx^2} = \frac{-(1 + \log x)}{(x \log x)^2}.$$

10. Determine the second order derivative for the follo wing function. y=sin(logx).

Ans: The given function is y=sin(logx).

Now, differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{d}{dx} [\sin(\log x)] = \cos(\log x) \times \frac{d}{dx} (\log x) = \frac{\cos(\log x)}{x}$$

$$\frac{d^{2}y}{dx^{2}} = \frac{d}{dx} \left[\frac{\cos(\log x)}{x} \right]$$
$$= \frac{x \left[\cos(\log x) \right] - \cos(\log x) \times \frac{d}{dx}(x) \right]}{x^{2}}$$
$$= \frac{x \left[-\sin(\log x) \times \frac{d}{dx}(\log x) \right] - \cos(\log x) \times 1}{x^{2}}$$
$$= \frac{-x \sin(\log x) \times \frac{1}{x} - \cos(\log x)}{x^{2}}$$
Hence,
$$\frac{d^{2}y}{dx^{2}} = \frac{\left[-\sin(\log x) + (\log x) \right]}{x^{2}}.$$

11. Prove that
$$\frac{d^2y}{dx^2} + y = 0$$
 when y=5cosx-3sinx.

Ans: The given equation is $y=5\cos 3\sin x$.

Then, differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{d}{dx}(5\cos x) - \frac{d}{dx}(3\sin x) = 5\frac{d}{dx}(\cos x) - 3\frac{d}{dx}(\sin x) = 5(-\sin x) - 3\cos x$$

Therefore, $\frac{dy}{dx} = -(5\sin x + 3\cos x)$.

$$\frac{d^2y}{dx^2} = \frac{d}{dx} [-(5\sin x + 3\cos x)]$$
$$= -\left[5 \times \frac{d}{dx} (\sin x) + 3 \times \frac{d}{dx} (\cos x)\right]$$
$$= [5\cos x + 3(-\sin x)]$$
$$= -y$$

That is,
$$\frac{d^2y}{dx^2} = -y$$
.
Hence, $\frac{d^2y}{dx^2} + y = 0$.

12. Determine $\frac{d^2y}{dx^2}$ containing the terms of y only when y=cos⁻¹x.

Ans: The given function is $y=\cos^{-1}x$.

Now, differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{d}{dx}(\cos^{-1}x) = \frac{-1}{\sqrt{1-x^2}} = -(1-x^2)^{\frac{-1}{2}}$$

Again, differentiating both sides with respect to x gives

$$\frac{d^{2}y}{dx^{2}} = \frac{d}{dx} \left[-(1-x^{2})^{\frac{-1}{2}} \right]$$

$$= \left(\frac{-1}{2} \right) \times (1-x^{2})^{\frac{-3}{2}} \times \frac{d}{dx} (1-x^{2})$$

$$= \frac{1}{\sqrt[2]{(1-x^{2})^{3}}} \times (-2x)$$

$$\Rightarrow \frac{d^{2}y}{dx^{2}} = \frac{-x}{\sqrt{(1-x^{2})^{3}}} \qquad \dots \dots (1)$$

Now, $y = \cos^{-1}x \implies x = \cos y$.

Therefore, substituting x=cosy into equation (1), gives

$$\frac{\mathrm{d}^2 \mathrm{x}}{\mathrm{d} \mathrm{y}^2} = \frac{-\mathrm{cosy}}{\sqrt{(1-\mathrm{cos}^2 \mathrm{y})^3}}$$

$$= \frac{-\cos y}{\sin^3 y}$$
$$= \frac{-\cos y}{\sin y} \times \frac{1}{\sin^2 y}$$
Hence, $\frac{d^2 y}{dx^2} = \cot y \times \csc^2 y$.

13. Prove that $x^2y_2+xy_1+y=0$ when $y=3\cos(\log x)+4\sin(\log x)$.

Ans: The given equations are
$$y=3\cos(\log x)+4\sin(\log x)$$
(1)

and
$$x^2y_2 + xy_1 + y = 0$$
 (2)

Then, differentiating both sides of the equation (1) with respect to x gives

$$y_{1} = 3 \times \frac{d}{dx} [\cos(\log x)] + 4 \times \frac{d}{dx} [\sin(\log x)]$$
$$= 3 \times \left[-\sin(\log x) \times \frac{d}{dx} (\log x) \right] + 4 \times \left[\cos(\log x) \times \frac{d}{dx} (\log x) \right]$$
$$y_{1} = \frac{-3\sin(\log x)}{x} + \frac{4\cos(\log x)}{x} = \frac{4\cos(\log x) - 3\sin(\log x)}{x}$$

$$y_{2} = \frac{d}{dx} \left(\frac{4\cos(\log x) - 3\sin(\log x)}{x} \right)$$

= $\frac{x\{4\cos(\log x) - 3\sin(\log x)\} - \{4\cos(\log x) - 3\sin(\log x)\}\}}{x^{2}}$
= $\frac{x[4\{\cos(\log x)\} - \{3\sin(\log x)\}] - \{4\cos(\log x) - 3\sin(\log x)\} \times 1]}{x^{2}}$
= $\frac{x[-4\sin(\log x)(\log x)' - 3\cos(\log x)(\log x)] - 4\cos(\log x) + 3\sin(\log x)]}{x^{2}}$

$$=\frac{x\left[-4\sin(\log x)\frac{1}{x}-3\cos(\log x)\frac{1}{x}\right]-4\cos(\log x)+3\sin(\log x)}{x^{2}}$$
$$=\frac{-4\sin(\log x)-3\cos(\log x)-4\cos(\log x)+3\sin(\log x)}{x^{2}}$$
Therefore, $y_{2}=\frac{-\sin(\log x)-7\cos(\log x)}{x^{2}}$.

Now, substituting the derivatives y_1 , y_2 and y into the LHS of the equation (2) gives

$$x^{2}y_{2}+xy_{1}+y$$

$$=x^{2}\left(\frac{-\sin(\log x)-7\cos(\log x)}{x^{2}}\right)+x\left(\frac{4\cos(\log x)-3\sin(\log x)}{x^{2}}\right)+3\cos(\log x)+4\sin(\log x)$$

$$=-\sin(\log x)-7\cos(\log x)+4\cos(\log x)-3\sin(\log x)+4\sin(\log x)$$

$$=0$$

Hence, it has been proved that $x^2y_2 + xy_1 + y = 0$.

14. Prove that
$$\frac{d^2y}{dx^2}$$
-(m+n) $\frac{dy}{dx}$ +mny=0 when y=Ae^{mx}+Be^{nx}.

Ans: The given equations are $y=Ae^{mx}+Be^{nx}$

and
$$\frac{d^2 y}{dx^2} - (m+n)\frac{dy}{dx} + mny = 0$$
(2)

Then, differentiating both sides of the equation (1) with respect to x gives

..... (1)

$$\frac{dy}{dx} = A \cdot \frac{d}{dx}(e^{mx}) + B \cdot \frac{d}{dx}(e^{mx}) = A \cdot e^{mx} \cdot \frac{d}{dx}(mx) + B \cdot e^{nx} \cdot \frac{d}{dx}(nx) = Ame^{mx} + Bne^{nx}$$

$$\frac{d^2y}{dx^2} = \frac{d}{dx} (Ame^{mx} + Bne^{nx}) = Am \cdot \frac{d}{dx} (e^{mx}) + Bn \cdot \frac{d}{dx} (e^{nx})$$

=Am.e^{mx}.
$$\frac{d}{dx}(mx)$$
+Bn.e^{nx}. $\frac{d}{dx}(nx)$
Therefore, $\frac{d^2y}{dx^2}$ =Am²e^{mx}+Bn²e^{nx}.

Thus, substituting the derivatives y_1 , y_2 and y into the LHS of the equation (2) gives

.....(1)

$$\frac{d^{2}y}{dx^{2}} - (m+n)\frac{dy}{dx} + mny$$

=Am²ex^{mx}+Bn²e^{nx}-(m+n).(Ame^{mx}+Bne^{nx})+mn(Ae^{mx}+Be^{nx})
=Am²ex^{mx}+Bn²e^{nx}-Amex^{mx}+Bmne^{nx}+Amne^{mx}+Bn²e^{nx}+Amne^{mx}+Bmne^{nx}=0

Thus, it has been proved that $\frac{d^2y}{dx^2}$ -(m+n) $\frac{dy}{dx}$ +mny=0.

15. Prove that
$$\frac{d^2y}{dx^2} = 49y$$
 when $y = 500e^{7x} + 600e^{-7x}$.

Ans: The given equation is $y=500e^{7x}+600e^{-7x}$

Then, differentiating both sides with respect to x gives

$$\frac{dy}{dx} = 500 \times (e^{7x}) + 600 \times \frac{d}{dx} (-7x)$$
$$= 500 \times e^{7x} \times \frac{d}{dx} (7x) + 600 \times e^{-7x} \times \frac{d}{dx} (-7x)$$
$$= 3500e^{7x} - 4200e^{-7x}$$

$$\frac{d^{2}y}{dx^{2}} = 3500 \times \frac{d}{dx} (e^{7x}) - 4200 \times \frac{d}{dx} (e^{-7x})$$
$$= 3500 \times e^{7x} \times \frac{d}{dx} (7x) - 4200 \times e^{-7x} \times \frac{d}{dx} (-7x)$$

$$=7 \times 3500 \times e^{7x} + 7 \times 4200 \times e^{-7x}$$
$$=49 \times 500e^{7x} + 49 \times 600e^{-7x}$$
$$=49(500e^{7x} + 600e^{-7x})$$

=49y, using the equation (1).

Thus, it has been proved that $\frac{d^2y}{dx^2} = 49y$.

16. Prove that
$$\frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$$
 when $e^y(x+1)=1$.

Ans: The given equation is $e^{y}(x+1)=1$.

Now,
$$e^{y}(x+1)=1 \Longrightarrow e^{y}=\frac{1}{x+1}$$
.

So, taking logarithm bth sides of the equation gives

$$y = \log \frac{1}{(x+1)}$$

Therefore, differentiating both sides with respect to x gives

$$\frac{dy}{dx} = (x+1)\frac{d}{dx}\left(\frac{1}{x+1}\right) = (x+1) \times \frac{-1}{(x+1)^2} = \frac{-1}{x+1}$$

That is,

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{-1}{x+1} \qquad \dots \dots (1)$$

Again, differentiating both sides with respect to x gives

$$\frac{d^2 y}{dx^2} = \frac{d}{dx} = \left(\frac{1}{x+1}\right) = -\left(\frac{-1}{(x+1)^2}\right) = \frac{1}{(x+1)^2}$$
$$\Rightarrow \frac{d^2 y}{dx^2} = \left(\frac{-1}{x+1}\right)^2$$
$$\Rightarrow \frac{d^2 y}{dx^2} = \left(\frac{dy}{dx}\right)^2, \text{ using the equation (1).}$$

Thus, it is proved that $\frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$.

17. Prove that
$$(x^2+1)^2y_2+2x(x^2+1)y_1=2$$
 when $y=(\tan^{-1}x)^2$.

Ans: The given equations are $y=(\tan^{-1}x)^2$.

Then, differentiating both sides with respect to x gives

$$y_1 = 2\tan^{-1}x \frac{d}{dx}(\tan^{-1}x)$$
$$\Rightarrow y_1 = 2\tan^{-1}x \times \frac{1}{1+x^2}$$
$$\Rightarrow (1+x^2)y_1 = 2\tan^{-1}x$$

Again, differentiating both sides with respect to x gives

$$(1+x^{2})y_{2}+2xy_{1}=2\left(\frac{1}{1+x^{2}}\right)$$

 $\Rightarrow (1+x^{2})y_{2}+2x(1+x^{2})y_{1}=2$

Thus, it has been proved that $(1+x^2)y_2+2x(1+x^2)y_1=2$.

Miscellaneous Exercise

1. Differentiate the function $y=(3x^2-9x+5)^9$ with respect to x.

Ans: The given function is $y=(3x^2-9x+5)^9$.

Differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{d}{dx} = (3x^2 - 9x + 5)^9$$

=9(3x^2 - 9x + 5)⁸× $\frac{d}{dx}$ (3x² - 9x + 5)
=9(3x^2 - 9x + 5)⁸×(6x - 9x)
=9(3x^2 - 9x + 5)⁸×3(2x - 3)
=27(3x^2 - 9x + 5)⁸(2x - 3)

2. Differentiate the function $y=\sin^3x+\cos^6x$ with respect to x.

Ans: The given function is $y=\sin^3 x + \cos^6 x$.

Differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{d}{dx} = (\sin^3 x) + \frac{d}{dx} (\cos^6 x)$$
$$= 3\sin^2 x \times \frac{d}{dx} (\sin x) + 6\cos^5 x \frac{d}{dx} (\cos x)$$
$$= 3\sin^2 x \times \cos x + 6\cos^5 x (-\sin x)$$
$$= 3\sin x \cos x (\sin x - 2\cos^4 x)$$

3. Differentiate the function $y=(5x)^{3\cos 2x}$ with respect to x.

Ans: The given function is $y=(5x)^{3\cos 2x}$.

First, take the logarithm of both sides of the function.

 $\log y = 3\cos 2x \log 5x.$

$$\frac{1}{y}\frac{dy}{dx} = 3\left[\log 5 \cdot \frac{d}{dx}(\cos 2x) + \cos 2x \cdot \frac{d}{dx}(\log 5x)\right]$$

$$\Rightarrow \frac{dy}{dx} = 3y\left[\log 5x(-\sin 2x) \cdot \frac{d}{dx}(2x) + \cos 2x \cdot \frac{1}{5x} \cdot \frac{d}{dx}(5x)\right]$$

$$\Rightarrow \frac{dy}{dx} = 3y\left[-2\sin 2x\log 5x + \frac{\cos 2x}{x}\right]$$

$$\Rightarrow \frac{dy}{dx} = 3y\left[\frac{3\cos 2x}{x} - 6\sin 2x\log 5x\right]$$

Hence, $\frac{dy}{dx} = (5x)^{3\cos 2x}\left[\frac{3\cos 2x}{x} - 6\sin 2x\log 5x\right]$.

- 4. Differentiate the functiony=sin⁻¹ $(x\sqrt{x})$, $0 \le x \le 1$ with respect to x.
- **Ans:** The given function is $y=\sin^{-1}(x\sqrt{x})$.

Then, differentiating both sides with respect to x by using the chain rule gives

$$\frac{dy}{dx} = \frac{d}{dx} \sin^{-1} \left(x \sqrt{x} \right)$$
$$= \frac{1}{\sqrt{1 - \left(x \sqrt{x} \right)^3}} \times \frac{d}{dx} \left(x \sqrt{x} \right)$$
$$= \frac{1}{\sqrt{1 - x^3}} \cdot \frac{d}{dx} \left(x^{\frac{1}{3}} \right)$$
$$= \frac{1}{\sqrt{1 - x^3}} \times \frac{3}{2} \cdot x^{\frac{1}{2}}$$
$$= \frac{3\sqrt{x}}{2\sqrt{1 - x^3}}$$

Hence, $\frac{dy}{dx} = \frac{3}{2}\sqrt{\frac{x}{1-x^3}}$.

5. Differentiate the function $y = \frac{\cos^{-1} \frac{x}{2}}{\sqrt{2+7}}$, -2<x<2 with respect to x.

Ans: The given function is
$$y = \frac{\cos^{-1} \frac{x}{2}}{\sqrt{2+7}}$$
.

Then, differentiating both sides with respect to x using the quotient rule gives

$$\frac{dy}{dx} = \frac{\sqrt{2x+7} \frac{d}{dx} \left(\cos^{-1} \frac{x}{2}\right) - \left(\cos^{-1} \frac{x}{2}\right) \frac{d}{dx} \left(\sqrt{2x+7}\right)}{\left(\sqrt{2x+7}\right)^2}$$

$$= \frac{\sqrt{2x+7} \left[\frac{-1}{\sqrt{1-\left(\frac{x}{2}\right)^2}} \cdot \frac{d}{dx} \left(\frac{x}{2}\right)\right] - \left(\cos^{-1} \frac{x}{2}\right) \frac{1}{2\sqrt{2x+7}} \cdot \frac{d}{dx} (2x+7)}{2x+7}\right]$$

$$= \frac{\sqrt{2x+7} \frac{-1}{\sqrt{4-x^2}} - \left(\cos^{-1} \frac{x}{2}\right) \frac{2}{2\sqrt{2x+7}}}{2x+7}$$

$$= \frac{-\sqrt{2x+7}}{\sqrt{4-x^2} \times (2x+7)} - \frac{\cos^{-1} \frac{x}{2}}{\left(\sqrt{2x+7}\right)(2x+7)}$$
Hence, $\frac{dy}{dx} = -\left[\frac{1}{\sqrt{4-x^2}\sqrt{2x+7}} + \frac{\cos^{-1} \frac{x}{2}}{\left(2x+7\right)^2}\right]$.

6. Differentiate the function $y = \cot^{-1} \left[\frac{\sqrt{1 + \sin x} + \sqrt{1 - \sin x}}{\sqrt{1 + \sin x} - \sqrt{1 - \sin x}} \right]$, 0<x<2 with respect to x.

Ans: The given function is
$$y = \cot^{-1} \left[\frac{\sqrt{1 + \sin x} + \sqrt{1 - \sin x}}{\sqrt{1 + \sin x} - \sqrt{1 - \sin x}} \right]$$
(1)

Now,
$$\frac{\sqrt{1+\sin x} + \sqrt{1-\sin x}}{\sqrt{1+\sin x} - \sqrt{1-\sin x}}$$
$$= \frac{\left(\sqrt{1+\sin x} + \sqrt{1-\sin x}\right)}{\left(\sqrt{1+\sin x} - \sqrt{1-\sin x}\right)\sqrt{1+\sin x} + \sqrt{1-\sin x}}$$
$$= \frac{(1+\sin x) + (1-\sin x) + 2\sqrt{(1+\sin x) - (1-\sin x)}}{(1+\sin x) - (1-\sin x)}$$
$$= \frac{2+2\sqrt{1-\sin^2 x}}{2\sin x}$$
$$= \frac{1+\cos x}{\sin x}$$
$$= \frac{2\cos^2 \frac{x}{2}}{2\sin x \frac{x}{2}\cos \frac{x}{2}}$$

Therefore,

$$\frac{\sqrt{1+\sin x} + \sqrt{1-\sin x}}{\sqrt{1+\sin x} - \sqrt{1-\sin x}} = \cot \frac{x}{2}.$$
 (2)

So, from the equations (1) and (2) we obtain,

$$y = \cot^{-1} \left(\cot^{\frac{x}{2}} \right)$$
$$\Rightarrow y = \frac{x}{2}$$

Now, differentiating both sides with respect to x gives

 $\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}x}(x)$

Hence,
$$\frac{dy}{dx} = \frac{1}{2}$$
.

7. Differentiate the function $y = (\log x)^{\log x}$, x > 1 with respect to x.

Ans: The given function is $y=(\log x)^{\log x}$.

First take logarithm both sides of the function.

logy=logx×log(logx).

Now, differentiating both sides with respect to x gives

$$\frac{1}{y}\frac{dy}{dx} = \frac{d}{dx} \left[\log x \times \log(\log x) \right]$$

$$\Rightarrow \frac{1}{y}\frac{dy}{dx} = \log(\log x) \times \frac{d}{dx}(\log x) + \frac{d}{dx} \left[\log(\log x) \right]$$

$$\Rightarrow \frac{dy}{dx} = y \left[\log(\log x) \times \frac{1}{x} + \log x \times \frac{1}{\log x} \times \frac{d}{dx}(\log x) \right]$$

$$\Rightarrow \frac{dy}{dx} = y \left[\frac{1}{x} \log(\log x) + \frac{1}{x} \right]$$
Hence, $\frac{dy}{dx} = (\log x)^{\log x} \left[\frac{1}{x} + \frac{\log(\log x)}{x} \right].$

8. Differentiate the function y=cos(acosx+bsinx), where a and b are any constants.

Ans: The given function is y=cos(acosx+bsinx).

Now, differentiating both sides with respect to x by using the chain rule of derivatives gives

 $\frac{dy}{dx} = \frac{d}{dx}\cos(a\cos x + b\sin x)$

$$\Rightarrow \frac{dy}{dx} = -\sin(a\cos x + b\sin x) \times \frac{d}{dx}(a\cos x + b\sin x)$$
$$= -\sin(a\cos x + b\sin x) \times [a(-\sin x) + b\cos x]$$
Hence, $\frac{dy}{dx} = (a\sin x + b\cos x) \times \sin(a\cos x + b\sin x)$.

9. Differentiate the function y=(sinx-cosx)^(sinx-cosx),
$$\frac{\pi}{4} < x < \frac{3\pi}{4}$$
 with respect to x

Ans: The given function is $y=(sinx-cosx)^{(sinx-cosx)}$.

First take logarithm both sides of the function.

 $logy=log[(sinx-cosx)^{(sinx-cosx)}]$ $\Rightarrow logy=(sinx-cosx) \times log(sinx-cosx)$

Now, differentiating both sides with respect to x gives

$$\frac{1}{y}\frac{dy}{dx} = \frac{d}{dx} [(\sin x - \cos x) \times \log(\sin x - \cos x)]$$

$$\Rightarrow \frac{1}{y}\frac{dy}{dx} = \log(\sin x - \cos x) \times \frac{d}{dx}(\sin x - \cos x) + (\sin x - \cos x) \times \frac{d}{dx}\log(\sin x - \cos x)$$

$$\Rightarrow \frac{1}{y}\frac{dy}{dx} = \log(\sin x - \cos x) \times (\cos x + \sin x) + (\sin x - \cos x) \times \frac{1}{(\sin x - \cos x)} \times \frac{d}{dx}(\sin x - \cos x)$$

$$\Rightarrow \frac{dy}{dx} = (\sin x - \cos x)^{(\sin x - \cos x)} [(\cos x + \sin x) \times \log(\sin x - \cos x) + (\cos x + \sin x)]$$

Hnece, the required derivative is

$$\frac{dy}{dx} = (\sin x - \cos x)^{(\sin x - \cos x)} (\cos x + \sin x) [1 + \log(\sin x - \cos x)].$$

10. Differentiate the function $y=x^x+x^a+a^x+a^a$ with respect to x, where for a>0 and x>0 are any fixed constants.

Ans: The given function is $y=x^{x}+x^{a}+a^{x}+a^{a}$.

Now, assume that $x^x = u$, $x^a = v$, $a^x = w$ and $a^a = s$

Therefore, we have y=u+v+w+s.

So, differentiating both sides with respect to x gives

$$\frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} + \frac{dw}{dx} + \frac{ds}{dx} \qquad \dots \dots (1)$$

Also, $u = x^{x}$
 $\Rightarrow \log u = \log x^{x}$
 $\Rightarrow \log u = x \log x$

Then, differentiating both sides with respect to x gives

$$\frac{1}{u}\frac{du}{dx} = \log x \cdot \frac{d}{dx}(x) + x \cdot \frac{d}{dx}(\log x)$$

$$\Rightarrow \frac{du}{dx} = u \left[\log x \cdot 1 + x \cdot \frac{1}{x}\right]$$

Thus, $\frac{du}{dx} = x^{x} [\log x + 1] = x^{x} (1 + \log x)$ (2)
Again, $v = x^{a}$

Then, differentiating both sides with respect to x gives

 $\frac{du}{dx} = \frac{d}{dx}(x^{a})$ $\Rightarrow \frac{dv}{dx} = ax^{a \cdot 1} \qquad \dots (3)$ Also, w=a^x $\Rightarrow \log w = \log a^{x}$ $\Rightarrow \log w = x \log a$

$$\frac{1}{w} \cdot \frac{dw}{dx} = \log a \cdot \frac{d}{dx}(x)$$

$$\Rightarrow \frac{dw}{dx} = w \log a$$

$$\Rightarrow \frac{dw}{dx} = a^{x} \log a \qquad \dots \dots (4)$$

and

s=a^a

Then differentiating both sides with respect to x gives

$$\frac{\mathrm{ds}}{\mathrm{dx}} = 0, \qquad \dots \dots (5)$$

as a is constant, and so a^a is also a constant.

Now, from the equations (1), (2), (3), (4), and (5) we have

$$\frac{\mathrm{dy}}{\mathrm{dx}} = x^2 (1 + \log x) + ax^{a-1} + a^x \log a + 0$$

Hence,
$$\frac{dy}{dx} = x^2(1 + \log x) + ax^{a-1} + a^x \log a$$

11. Differentiate the function $y=x^{x^2-3}+(x-3)^{x^2}$, for x>3 with respect to x.

Ans: The given function is $y=x^{x^2-3}+(x-3)^{x^2}$.

Now suppose that $u=x^{x^2-3}$ and $v=(x-3)^{x^2}$

Therefore, y=u+v.

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}u}{\mathrm{d}x} + \frac{\mathrm{d}v}{\mathrm{d}x} \qquad \dots \dots (1)$$

Also, $u=x^{x^2-3}$.

Take logarithm both sides of the equation.

$$\Rightarrow \log u = \log(x^{x^2 \cdot 3})$$
$$\Rightarrow \log u = (x^2 \cdot 3) \log x$$

Differentiating both sides with respect to x gives

$$\frac{1}{u}\frac{du}{dx} = \log x \cdot \frac{d}{dx}(x^2 - 3) + (x^2 - 3) \cdot \frac{d}{dx}(\log x)$$

$$\Rightarrow \frac{1}{u}\frac{du}{dx} = \log x \cdot 2x + (x^2 - 3) \cdot \frac{1}{x}$$

Hence, $\frac{du}{dx} = x^{x^2 - 3} \cdot \left[\frac{x^2 - 3}{x} + 2 \times \log x\right]$(2)

Again, $v=(x-3)^{x^2}$.

Take logarithm both sides of the equation.

$$\Rightarrow \log v = \log(x-3)^{x^2}$$
$$\Rightarrow \log v = x^2 \log(x-3)$$

Now, differentiating both sides with respect to x gives

$$\frac{1}{u} \cdot \frac{dv}{dx} = \log(x-3) \cdot \frac{d}{dx} (x^2) + x^2 \cdot \frac{d}{dx} [\log(x-3)]$$

$$\Rightarrow \frac{1}{u} \cdot \frac{dv}{dx} = \log(x-3) \cdot 2x + x^2 \cdot \frac{1}{x-3} \cdot \frac{d}{dx} (x-3)$$

$$\Rightarrow \frac{dv}{dx} = v \cdot \left[2x \log(x-3) + \frac{x^2}{x-3} \cdot 1 \right]$$
Hence, $\frac{dv}{dx} = (x-3)^{x^2} \left[\frac{x^2}{x-3} + 2x \log(x-3) \right]$ (3)

Thus, from the equations (1), (2) and (3) we obtain

$$\frac{dy}{dx} = x^{x^2 - 3} \left[\frac{x^2 - 3}{x} + 2x \log x \right] + (x - 3)^{x^2} \left[\frac{x^2}{x - 3} + 2x \log(x - 3) \right].$$

12. Determine $\frac{dy}{dx}$ from the parametric equations

y=12(1-cost),x=10(t-sint), $\frac{\pi}{2} < t < \frac{\pi}{2}$, without eliminating the parameter t.

Ans: The given equations are $y=12(1-\cos t)$,(1)

and
$$x=10(t-sint)$$
 (2)

Then differentiating the equations (1) and (2) with respect to x gives

$$\frac{dx}{dt} = \frac{d}{dt} [10(t-sint)] = 10 \times \frac{d}{dt} (t-sint) = 10(1-cost)$$
$$\frac{dy}{dt} = \frac{d}{dt} [12(1-cost)] = 12 \times \frac{d}{dt} (1-cost) = 12 \times [0-(-sint)] = 12sint$$

Therefore, by dividing
$$\frac{dy}{dt}$$
 by $\frac{dx}{dt}$ we have,

$$\frac{\mathrm{dy}}{\mathrm{dx}} = \frac{\left(\frac{\mathrm{dy}}{\mathrm{dt}}\right)}{\left(\frac{\mathrm{dx}}{\mathrm{dt}}\right)} = \frac{12\mathrm{sint}}{10(1-\mathrm{cost})} = \frac{12\times2\mathrm{sin}\frac{\mathrm{t}}{2}\times\mathrm{cos}\frac{\mathrm{t}}{2}}{10\times2\mathrm{sin}^2\frac{\mathrm{t}}{2}}$$

Hence,
$$\frac{dy}{dx} = \frac{6}{5}\cot\frac{t}{2}$$
.

13. Determine $\frac{dy}{dx}$ from the equation $y = \sin^{-1}x + \sin^{-1}\sqrt{1-x^2}$, $-1 \le x \le 1$.

Ans: The given equation is $y=\sin^{-1}x+\sin^{-1}\sqrt{1-x^2}$.

Differentiating both sides of the equation with respect to x gives

$$\frac{dy}{dx} = \frac{d}{dx} \left[\sin^{-1}x + \sin^{-1}\sqrt{1 - x^2} \right]$$

$$\Rightarrow \frac{dy}{dx} = \frac{d}{dx} (\sin^{-1}x) + \frac{d}{dx} (\sin^{-1}\sqrt{1 - x^2})$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{\sqrt{1 - x^2}} + \frac{1}{\sqrt{1}(\sqrt{1 - x^2})} \times \frac{d}{dx} (\sqrt{1 - x^2})$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{\sqrt{1 - x^2}} + \frac{1}{2 \times \sqrt{1 - x^2}} (-2)$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{\sqrt{1 - x^2}} - \frac{1}{\sqrt{1 - x^2}}$$
Hence, $\frac{dy}{dx} = 0$.

14. Prove that
$$\frac{dy}{dx} = -\frac{1}{(1+x)^2}$$
 when $x\sqrt{1+y} + y\sqrt{1+x} = 0$, for $-1 < x < 1$.

Ans: The given equation is

$$x\sqrt{1+y}+y\sqrt{1+x}=0$$
$$\Rightarrow x\sqrt{1+y}=y\sqrt{1+x}$$

Now, squaring both sides of the equation, gives

$$x^{2}(1+y)=y^{2}(1+x)$$

$$\Rightarrow x^{2}+x^{2}y=y^{2}+xy^{2}$$

$$\Rightarrow x^{2}-y^{2}=xy^{2}-x^{2}y$$

$$\Rightarrow x^{2}-y^{2}=xy(y-x)$$

$$\Rightarrow (x+y)(x-y)=xy(y-x)$$

$$\therefore x+y=-xy$$

$$\Rightarrow (1+x)y=x$$

$$\Rightarrow y=\frac{-x}{(1+x)}$$

Now, differentiating both sides of the equation with respect to x gives

$$\frac{dy}{dx} = \frac{(1+x)\frac{d}{dx}(x) - x\frac{d}{dx}(1+x)}{(1+x)^2} = \frac{(1+x) - x}{(1+x)^2}$$

Hence,
$$\frac{dy}{dx} = \frac{1}{(1+x)^2}$$
.

15. prove that
$$\frac{\left[1+\left(\frac{dy}{dx}\right)^2\right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}$$
 is a constant independent from a and b,

when $(x-a)^2+(y-b)^2=c^2$, for some constant c>0.

Ans: The given equation is
$$(x-a)^2+(y-b)^2=c^2$$
.

Differentiating both sides of the equation with respect to x gives

$$\frac{d}{dx} = [(x-a)^2] + \frac{d}{dx} [(y-b)^2] = \frac{d}{dx} (c^2)$$

$$\Rightarrow 2(x-a) \cdot \frac{d}{dx} (x-a) + 2(y-b) \cdot \frac{d}{dx} (y-b) = 0$$

$$\Rightarrow 2(x-a) \cdot 1 + 2(y-b) \cdot \frac{dy}{dx} = 0$$

Hence, $\frac{dy}{dx} = \frac{-(x-a)}{y-b}$ (1)

Again, differentiating both sides of the equation with respect to x gives

$$\frac{d^2y}{dx^2} = \frac{d}{dx} \left[\frac{-(x-a)}{y-b} \right]$$

$$= -\frac{\left[(y-b) \cdot \frac{d}{dx} (x-a) - (x-a) \cdot \frac{d}{dx} (y-b) \right]}{(y-b)^2}$$
$$= -\left[\frac{(y-b) - (x-a) \cdot \frac{dy}{dx}}{(y-b)^2} \right]$$
$$= -\left[\frac{(y-b) - (x-a) \cdot \left\{ \frac{-(x-a)}{y-b} \right\}}{(y-b)^2} \right]$$
$$= -\left[\frac{(y-b)^2 + (x+a)^2}{(y-b)^2} \right]$$

Therefore,

$$\left[\frac{1+\left(\frac{dy}{dx}\right)^{2}}{\left(\frac{d^{2}y}{dx^{2}}\right)^{2}}\right]^{\frac{3}{2}} = \frac{\left[\left(1+\frac{(x-a)^{2}}{(y-b)^{2}}\right)\right]^{\frac{3}{2}}}{-\left[\frac{(y-a)^{2}+(x-a)^{2}}{(y-a)^{3}}\right]} = \frac{\left[\frac{(y-b)^{2}+(x-a)^{2}}{(y-b)^{2}}\right]^{\frac{3}{2}}}{-\left[\frac{(y-a)^{2}+(x-a)^{2}}{(y-a)^{3}}\right]} = \frac{\left[\frac{(y-b)^{2}}{(y-b)^{2}}\right]^{\frac{3}{2}}}{-\frac{(y-b)^{3}}{(y-b)^{3}}}$$
$$\Rightarrow \left[\frac{1+\left(\frac{dy}{dx}\right)^{2}}{\frac{d^{2}y}{dx^{2}}}\right]^{\frac{3}{2}} = \frac{c^{2}}{\frac{(y-b)^{3}}{(y-b)^{3}}} = -c, \text{ is a constant, and is independent of a and b.}$$

16. Prove that $\frac{dy}{dx} = \frac{\cos^2(a+y)}{\sin a}$, $\cos a \neq \pm 1$ from the equation $\cos y = x\cos(a+y)$.

Ans: The given equation is cosy=xcos(a+y).

Then, differentiating both sides of the equation with respect to x gives

$$\frac{d}{dx}[\cos y] = \frac{d}{dx}[x\cos(a+y)]$$

$$\Rightarrow -\sin y \frac{dy}{dx} = \cos(a+y) \cdot \frac{d}{dx}(x) + x \cdot \frac{d}{dx}[\cos(a+y)]$$

$$\Rightarrow -\sin y \frac{dy}{dx} = \cos(a+y) + x \cdot [-\sin(a+y)] \frac{dy}{dx}$$

$$\Rightarrow [x\sin(a+y) - \sin y] \frac{dy}{dx} = \cos(a+y) \qquad \dots \dots (1)$$
Since $\cos y = x\cos(a+y) \Rightarrow x = \frac{\cos y}{\cos(a+y)}$, so from the equation (1) gives

$$\left[\frac{\cos y}{\cos(a+y)}.\sin(a+y)-\sin y\right]\frac{dy}{dx}=\cos(a+y)$$
$$\Rightarrow [\cos y.\sin(a+y)-\sin y.\cos(a-y)].\frac{dy}{dx}=\cos^2(a+y)$$
$$\Rightarrow \sin(a+y-y)\frac{dy}{dx}=\cos^2(a+b)$$

Hence, it has been proved that
$$\frac{dy}{dx} = \frac{\cos^2(a+b)}{\sin a}$$
.

- 17. Determine $\frac{d^2y}{dx^2}$ from the parametric equations x=a(cost+tsint) and y=a(sint-tcost), without cancelling the parameter t.
 - $x=a(cost+tsint) \qquad \dots \dots (1)$ and y=a(sint-tcost) \qquad \dots \dots (2)

The given equations are

Ans:

Then, differentiating both sides of the equation (1) with respect to x gives

$$\frac{dx}{dt} = a \left[-\sin t + \sin t \cdot \frac{d}{dx}(t) + t \cdot \frac{d}{dt}(\sin t) \right]$$
$$= a \left[-\sin t + \sin t + \cos t \right] = a \cos t$$

Again, differentiating both sides of the equation (2) with respect to x gives

$$\frac{dy}{dt} = a \cdot \frac{d}{dt} (sint-tcost)$$
$$a \left[cost \cdot \left\{ cost \cdot \frac{d}{dt}(t) + t \cdot \frac{d}{dt}(cost) \right\} \right]$$

a[cost-{cost-tsint}]=atsint

Therefore,

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dx}\right)} = \frac{atsint}{atcost} = tant$$

Now, differentiating both sides with respect to x gives

$$\frac{d^{2}y}{dx^{2}} = \frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{dx} (tant) = \sec^{2} t. \frac{dt}{dx} = \sec^{2} t. \frac{1}{atcost}$$
Hence, $\frac{d^{2}y}{dx^{2}} = \frac{\sec^{3} t}{at}, 0 < t < \frac{\pi}{2}$ $\left[\because \frac{dx}{dt} = atcost \Rightarrow \frac{dt}{dx} = \frac{1}{atcost} \right]$

18. Prove that f''(x) exists for all real values of x when $f(x)=|x|^3$ and hence evaluate it.

Ans: Remember that,
$$|x| = \begin{cases} x, \text{if } x \ge 0 \\ -x, \text{if } x < 0 \end{cases}$$

Therefore, if $x \ge 0$, then $f(x) = |x|^3 = x^3$.

Then, $f'(x)=3x^2$.

Differentiating both sides with respect to x gives

$$f''(x) = 6x$$
.

Now, if x<0, then $f(x)=|x|^3=(-x^3)=x^3$.

So,
$$f'(x)=3x^2$$

Therefore, differentiating both sides with respect to x gives

$$f''(x)=6x.$$

Hence, for $f(x)=|x|^3$, f''(x) exists for all real values of x and is provided as

$$f''(x) = \begin{cases} 6x, \text{if } x \ge 0 \\ -6x, \text{if } x < 0 \end{cases}.$$

19. Derive the sum formula for cosine from the sum formula of sine sin(A+B)=sinAcosB+cosAsinB, by using differentiation.

Ans: The given sum formula is sin(A+B)=sinAcosB+cosAsinB.

Now, differentiating both sides with respect to x gives

$$\frac{d}{dx} [\sin(A+B)] = \frac{d}{dx} (\sin A \cos B) + \frac{d}{dx} (\cos A \sin B)$$

$$\Rightarrow \cos(A+B) \times \frac{d}{dx} (A+B) = \cos B \times \frac{d}{dx} (\sin A) + \sin A \times \frac{d}{dx} (\cos B) + \sin B \times \frac{d}{dx} (\cos A)$$

$$+ \cos A \times \frac{d}{dx} (\sin B)$$

$$\Rightarrow \cos(A+B) \times \frac{d}{dx} (A+B) = \cos B \times \cos A \frac{d}{dx} + \sin A (-\sin B) \frac{dB}{dx} + \sin B (-\sin A) \times \frac{dA}{dx}$$

$$+ \cos A \cos B \frac{dB}{dx}$$

$$\Rightarrow \cos(A+B) \left[\frac{dA}{dx} + \frac{dB}{dx} \right] = (\cos A \cos B - \sin A \sin B) \times \left[\frac{dA}{dx} + \frac{dB}{dx} \right]$$


Hence the required sum formula for cosines is cos(A+B)=cosAcosB-sinAsinB.

20. Does there exist a function which is continuous everywhere but not differentiable at exactly tow points? Justify your answer.

Ans: Yes there exist such a function

f(x) = x - 1 + |x - 2|

As you can see the graph of the function, this function is continuous at every point. But it is differentiable at exactly two points, viz (1,1) and (2,1) because of a sharp turn.



Evaluate the determinant.

y=(mc-nb)f(x)-(lc-na)g(x)+(lb-ma)h(x).

$$\frac{dy}{dx} = \frac{d}{dx} [(mc-nb)f(x)] - \frac{d}{dx} [(lc-na)g(x)] + \frac{d}{dx} [(lb-ma)h(x)]$$

$$= (mc-nb)f(x) - (lc-na)g(x) + (lb-ma)h(x)$$

$$= \begin{vmatrix} f(x)g(x)h(x) \\ 1 & m & n \\ a & b & c \end{vmatrix}$$
Hence,
$$\frac{dy}{dx} = \begin{vmatrix} f(x)g(x)h(x) \\ 1 & m & n \\ a & b & c \end{vmatrix}$$
.

22. Prove that
$$(1-x^2)\frac{d^2y}{dx^2} - x\frac{dy}{dx} - a^2y = 0$$
 when $y = e^{a\cos^{-1}x}, -1 \le x \le 1$.

Ans: The given equation is $y=e^{a\cos^{-1}x}$.

Then take logarithm both sides of the equation.

 $\log y = a \cos^{-1} x \log e$ $\Rightarrow \log y = a \cos^{-1} x$

Now, differentiating both sides with respect to x gives

$$\frac{1}{y}\frac{dy}{dx} = ax\frac{1}{\sqrt{1-x^2}}$$
$$\frac{dy}{dx} = \frac{-ax}{\sqrt{1-x^2}}$$

Therefore, squaring both the sides of the equation, gives

$$\left(\frac{dy}{dx}\right)^2 = \frac{a^2y^2}{1-x^2}$$
$$\Rightarrow (1-x^2)\left(\frac{dy}{dx}\right)^2 = a^2y^2$$
$$\Rightarrow (1-x^2)\left(\frac{dy}{dx}\right)^2 = a^2y^2$$

Again, differentiating both sides with respect to x gives

$$\left(\frac{dy}{dx}\right)^{2} \frac{d}{dx}(1-x^{2}) + (1-x^{2}) \times \frac{d}{dx} \left[\left(\frac{dy}{dx}\right)^{2} \right] = a^{2} \frac{d}{dx}(y^{2})$$
$$\Rightarrow \left(\frac{dy}{dx}\right)^{2} (-2x) + (1-x^{2}) \times 2 \frac{dy}{dx} \times \frac{d^{2}y}{dx^{2}} = a^{2} \times 2y \times \frac{dy}{dx}$$
$$\Rightarrow x \frac{dy}{dx} + (1-x^{2}) \frac{d^{2}y}{dx^{2}} = a^{2} \times y$$

Hence, it is proved that $(1-x^2)\frac{d^2y}{dx^2}-x\frac{dy}{dx}-a^2y=0$.