statistics

12 Chapter

Exercise – 12.1

1. A survey conducted by an organisation for the cause of illness and death among the women between the ages 15-44 (in years) worldwide, found the following figures (in %)

S.No.	Causes	Female fatality rate (%)
1.	Reproductive health conditions	31.8
2.	Neuropsychiatric conditions	25.4
3.	Injuries	12.4
4.	Cardiovascular conditions	4.3
5.	Respiratory conditions	4.1
6.	Other causes	22.0

(i) Represent the information given above graphically.

Ans: The graph of the information presented above can be produced as follows by depicting causes on the x-axis and family fatality rate on the y-axis, and selecting an acceptable scale (1 unit = 5% for the y axis).



All the rectangle bars are of the same width and have equal spacing between them.

(ii) Which condition is the major cause of women's ill health and death worldwide?

Ans: Reproductive health issues are the leading cause of women's illness and mortality globally, affecting 31.8% of women.

(iii) Try to find out, with the help of your teacher, any two factors which play a major role in the cause in (ii) above being the major cause

Ans: The factors are as follows:

- 1. Lack of medical facilities
- 2. Lack of correct knowledge of treatment

Section	Number of girls per thousand boys
Scheduled caste (SC)	940
Scheduled tribe (ST)	970
Non SC/ST	920
Backward districts	950
Non – backward districts	920
Rural	930
Urban	910

2. The following data on the number of girls (to the nearest ten) per thousand boys in different sections of Indian society is given below:

(i) Represent the information above by a bar graph.

Ans: The graph of the information presented above may be built by choosing an appropriate scale (1 unit = 100 girls for the y-axis) and representing section (variable) on the x-axis and number of girls per thousand boys on the y-axis.



Here, all the rectangle bars are of the same length and have equal spacing in between them.

(ii) In the classroom discuss what conclusions can be arrived at from the graph.

Ans: The largest number of females per thousand boys (i.e., 970) is found in ST, while the lowest number of girls per thousand boys (i.e., 910) is found in urban.

In addition, the number of females per thousand boys is higher in rural regions than in cities, in backward districts than in non-backward districts, and in SC and ST districts than in non-SC/ST districts.

3. Given below are the seats won by different political parties in the polling outcome of a state assembly elections:

Political	Α	В	С	D	Ε	F
party						
Seats won	75	55	37	29	10	37

(i) Draw a bar graph to represent the polling results.

Ans:



Here, all the rectangle bars are of the same length and have equal spacing in between them.

(ii) Which political party won the maximum number of seats?

Ans: From the above graph it is clear that Political party 'A' won the maximum number of seats.

4. The length of 40 leaves of a plant are measured correct to one millimeter	er,
and the obtained data is represented in the following table:	

Length mm)	(in	Number of leaves
117.5-126.5		3
126.5-135.5		5
135.5-144.5		9
144.5-153.5		12
135.5-162.5		5
162.5-171.5		4
171.5-180.5		2

(i) Draw a histogram to represent the given data.

Ans: The length of leaves is represented in a discontinuous class interval with a difference of 1 between them, as can be seen. To make the class intervals continuous, $\frac{1}{2} = 0.5$ must be added to each upper class limit and 0.5 must be subtracted from the lower class limits.

Length (in mm)	Number of leaves
117.5-126.5	3
126.5-135.5	5
135.5-144.5	9
144.5-153.5	12
135.5-162.5	5
162.5-171.5	4
171.5-180.5	2



The above histogram may be built using the length of leaves on the x-axis and the number of leaves on the y-axis.

On the y-axis, one unit symbolises two leaves.

(ii) Is there any other suitable graphical representation for the same data?

Ans: Frequency polygon is another good graphical representation of this data.

(iii) Is it correct to conclude that the maximum number of leaves are 153 mm long? Why?

Ans: No, because the maximum number of leaves (i.e.12) has a length of 144.5mm to 153.5mm It is not necessary for all of them to be 153mm long.

Length (in hours)	Number of lamps
300-400	14
400-500	56
500-600	60
600-700	86
700-800	74
800-900	62
900-1000	48

5. The following table gives the life times of neon lamps:

(i) Represent the given information with the help of a histogram.

Ans: The histogram of the given data may be produced by plotting the life duration (in hours) of neon lamps on the x-axis and the number of lamps on the y-axis. Here,1



Here, 1 unit on y-axis represents 10 lamps.

(ii) How many lamps have a lifetime of more than 700 hours?

Ans: It may be deduced that the number of neon lamps with a lifetime more than 700 is equal to the sum of the numbers of neon lamps with lifetimes of 700,800 and 900. As a result, there are 184 neon bulbs with a lifetime of more than 700 hours (74+62+48=184).

according to the	e mark obtained by	tnem:		
Section A		Section B		
Marks	Frequency	Marks	Frequency	
0-10	3	0-10	5	
10-20	9	10-20	19	
20-30	17	20-30	15	
30-40	12	30-40	10	

6. The following table gives the distribution of students of two sections according to the mark obtained by them:

40-50	9	40-50	1

Represent the marks of the students of both the sections on the same graph by two frequency polygons. From the two polygons compare the performance of the two sections.

Ans: We can find the class marks of the given class intervals by using the following formula.

Class mark = $\frac{\text{Upper class limit} + \text{Lower class limit}}{\text{Lower class limit}}$

2

Section A			Section B		
Marks	Class marks	Frequency	Marks	Class marks	Frequency
0-10	5	3	0-10	5	5
10-20	15	9	10-20	15	19
20-30	25	17	20-30	25	15
30-40	35	12	30-40	35	10
40-50	45	9	40-50	45	1

The frequency polygon can be constructed as follows, with class markings on the x-axis and frequency on the y-axis, and an appropriate scale (1 unit = 3 for the y-axis).



It can be observed that the performance of students of section 'A' is better than the students of section 'B' in terms of good marks.

Number of balls	Class mark	Team A	Team B
0.5-6.5	3.5	2	5
6.5-12.5	9.5	1	6
12.5-18.5	15.5	8	2
18.5-24.5	21.5	9	10
24.5-30.5	27.5	4	5
30.5-36.5	33.5	5	6
36.5-42.5	39.5	6	3
42.5-48.5	45.5	10	4
48.5-54.5	51.5	6	8
54.5-60.5	57.5	2	10

7. The runs scored by two teams A and B on the first 60 balls in a cricket match are given below:

Represent the data of both the teams on the same graph by frequency polygons.

[Hint: First make the class intervals continuous.]

Ans: A frequency polygon can be created by plotting class grades on the x-axis and running times on the y-axis.



8. A random survey of the number of children of various age groups playing in park was found as follows:

Age (in years)	Number of children
1-2	5
2-3	3
3-5	6
5-7	12
7-10	9
10-15	10
15-17	4

Draw a histogram to represent the data above. Ans:

Age (in years)	Frequency	Width of class	Length of
	(Number of		rectangle
	children)		_
1-2	5	1	5×1 - 5
			$\frac{1}{1} = 3$

2-3	3	1	$\frac{3\times 1}{1} = 3$
3-5	6	2	$\frac{6\times 1}{2} = 3$
5-7	12	2	$\frac{12\times 1}{2} = 6$
7-10	9	3	$\frac{9\times 1}{3} = 3$
10-15	10	5	$\frac{10\times 1}{5} = 2$
15-17	4	2	$\frac{4 \times 1}{2} = 2$



9. 100 surnames were randomly picked up from a local telephone directory and a frequency distribution of the number of letters in the English alphabet in the surnames was found as follows:

Number of letters	Number of surnames
1-4	6
4-6	30

6-8	44	
8-12	16	
12-20	4	

(i) Draw a histogram to depict the given information.

Ans:

Number of	Frequency	Width of class	Length of
letters	(Number of		rectangle
	surnames)		
1-4	6	3	$\frac{6\times 2}{3} = 4$
4-6	30	2	$\frac{30 \times 2}{2} = 30$
6-8	44	2	$\frac{44 \times 2}{2} = 44$
8-12	16	4	$\frac{16 \times 2}{4} = 8$
12-20	4	8	$\frac{4 \times 2}{8} = 1$

The histogram can be generated using the number of letters on the x-axis and the fraction of the number of surnames per 2 letters interval on the y-axis, as well as an acceptable scale (1 unit = 4 students for the y axis).



(ii) Write the class interval in which the maximum number of surname lie.

Ans: The maximum number of surnames in the class interval is 68% since it contains 44 surnames, which is the maximum for this data.