## **Previous Year Question Paper**

### 2013

#### **General Instructions:**

(i) All questions are compulsory.

(ii) The question paper consists of 29 questions divided into three sections A, B and C. Section A comprises of10 questions of one mark each, Section B comprises of 12 questions of four marks each and Section Comprises of 7 questions of six marks each.

(iii) All questions in Section A are to be answered in one word, one sentence or as per the exact requirement of the questions.

(iv) There is no overall choice, However, internal choice has been provided in 4 questions of four marks each and 2 questions of six marks each. You have to attempt only one of the alternatives in all such questions.

(v) Use of calculators is not permitted. You may ask for logarithmic tables, if required.

1. Write the principal value of  $\tan^{-1}(\sqrt{3}) - \cot^{-1}(-\sqrt{3})$ 

Solution:

$$\tan^{-1}(\sqrt{3}) = \pi / 3$$

$$\cot^{-1}(-\sqrt{3}) = \pi - \pi / 6$$

Hence

$$\pi/3 - (\pi - \pi/6) = -\pi/2$$

2. Write the value of 
$$\tan^{-1} \left[ 2\sin\left(2\cos^{-1}\frac{\sqrt{3}}{2}\right) \right]$$
.

Solution:

$$\therefore \cos^{-1} \frac{\sqrt{3}}{2} = \frac{\pi}{6}$$
  
=  $\tan^{-1}(2\sin(2\pi/6))$   
=  $\tan^{-1}\left(2.\sin\frac{\pi}{3}\right)$   
=  $\tan^{-1}\left(2.\frac{\sqrt{3}}{2}\right) = \tan^{-1}\sqrt{3} = \pi/3$ 

3. For what value of x, is the matrix A =  $\begin{bmatrix} 0 & 1 & -2 \\ -1 & 0 & 3 \\ x & -3 & 0 \end{bmatrix}$  a skew-symmetric matrix ?

Sol. The value of determinant of skew symmetric matrix of odd order is always equal to zero

$$\begin{bmatrix} 0 & 1 & -2 \\ -1 & 0 & 3 \\ x & -3 & 0 \end{bmatrix} = 0$$
  
-1(0-3x)-2(3-0) = 0  
$$\Rightarrow 3x - 6 = 0 \Rightarrow \boxed{x = 2}$$

4. If matrix  $A = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$  and  $A^2 = kA$ , then write the value of k. **Sol.** Given A2 = kA $\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} = k \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$   $\begin{bmatrix} 2 & -2 \\ -2 & 2 \end{bmatrix} \begin{bmatrix} k & -k \\ -k & k \end{bmatrix} \Rightarrow \boxed{k = 2}$ 

5. Write the differential equation representing the family of curves y = mx, where m is an arbitrary constant. Sol. y = mx .....(1)

differentiating with respect to x, we get dy/dx = m

 $\therefore$  differential equation of curve

 $\begin{bmatrix} 2 & -3 & 5 \\ 6 & 0 & 4 \end{bmatrix}$ 

 $y = \frac{xdy}{dx}$ 

6. If  $A_{ij}$  is the cofactor of the element  $a_{ij}$  of the determinant  $\begin{bmatrix} 2 & -3 & 5 \\ 6 & 0 & 4 \\ 1 & 5 & -7 \end{bmatrix}$ , then write the value of  $a_{32} \cdot A_{32}$ .

Sol.

$$\begin{bmatrix} 1 & 5 & -7 \end{bmatrix}$$

$$A_{32} = (-1)^{3+2} M_{32} \text{ where } M_{32} \text{ is the min or of } a_{32}.$$

$$A_{32} = (-1)^{3+2} \begin{vmatrix} 2 & 5 \\ 6 & 4 \end{vmatrix}$$

$$A_{32} = -\begin{vmatrix} 2 & 5 \\ 6 & 4 \end{vmatrix} \Rightarrow A_{32} = -(8-30)$$

$$\boxed{A_{32} = 22}$$

$$\therefore a_{32}A_{32} = 5(22) = 110$$

7. P and Q are two points with position vectors  $3\vec{a} - 2\vec{b}$  and  $\vec{a} + \vec{b}$  respectively. Write the position vector of point R which divides the line segment PQ in the ratio 2 : 1 externally. 1

Sol. P.V. of P is  $3\vec{a} - 2\vec{b}$ P Q 1 P.V. of Q is  $\vec{a} + \vec{b}$ Point R divides segment PQ in ratio 2 : 1 externally. P.V.of R =  $\frac{(P.V.of p)1 - (P.V.of Q)(2)}{1 - 2}$ 

P.V.of R = 
$$\frac{(3\vec{a} - 2\vec{b})(1) - (\vec{a} + \vec{b})(2)}{1 - 2} = \frac{\vec{a} - 4\vec{b}}{-1}$$
  
P.V.of R =  $4\vec{b} - \vec{a}$ 

8. Find  $|\vec{x}|$ , if for a unit vector  $\vec{a}$ ,  $(\vec{x} - \vec{a})$ . $(\vec{x} + \vec{a}) = 15$ .

Sol. Given 
$$|\vec{a}| = 1$$
  
 $(\vec{x} - \vec{a}).(\vec{x} + \vec{a}) = 15$   
 $|\vec{x}|^2 - |\vec{a}|^2 = 15$   
 $|\vec{x}|^2 - 1 = 15$   
 $|\vec{x}|^2 = 15 + 1$   
 $|\vec{x}|^2 = 16$   
 $|\vec{x}| = 4$ 

9. Find the length of the perpendicular drawn from the origin to the plane 2x - 3y + 6z + 21 = 0.



10. The money to be spent for the welfare of the employees of a firm is proportional to the rate of change of its total revenue (marginal revenue). If the total revenue (in rupees) received from the sale of x units of a product is given by  $R(x) = 3x^2 + 36x + 5$ , find the marginal revenue, when x = 5, and write which value does the equations indicate.

Sol.  $R(x) = 3x^2 + 36x + 5$ 

$$MR = \frac{dR}{dx} = 6x^{2} + 36$$
  
when x = 5  
$$MR = 30 + 36 + 66$$

11. Consider  $f : R_+ \to [4, \infty)$  given by  $f(x) = x^2 + 4$ . Show that f is invertible with the inverse  $f^{-1}$  of f given by

 $f^{-1}(y) = \sqrt{y-4}$ , where R<sub>+</sub> is the set of all non-negative real numbers.

Sol. 
$$f: R^+ \rightarrow [4.\infty)$$
  
 $f(x) = x^2 + 4$   
 $f(x) = x^2 > 4$   $\therefore$  (one - one)  
As  $f(x) = x^2 + 4 \ge 4$   
 $\Rightarrow$  Rage =  $[4.\infty) = co - domain$   
 $\therefore$  onto  
Further :  $y = x^2 + 4$   
so f is invertible.  
 $\Rightarrow y - 4 = x^2 \Rightarrow x = \pm \sqrt{y - 4}$   
As  $x > 0$  so  $x = \sqrt{y - 4}$   
 $\therefore y = \sqrt{x - 4} = f^{-1}(x)$   
Or  $f^{-1}(y) - \sqrt{y - 4}$ 

12. Show that :

$$\tan\left(\frac{1}{2}\sin^{-1}\frac{3}{4}\right) = \frac{4-\sqrt{7}}{3}$$
*OR*

Solve the following equation :

$$\cos\left(\tan^{-1} x\right) = \sin\left(\cot^{-1} \frac{3}{4}\right)$$

Let 
$$\frac{1}{2}\sin^{-1}\frac{3}{4} = \theta$$
 then  $\frac{3}{4} = \sin 2\theta$   
Now  $\tan\left(\frac{1}{2}\sin^{-1}\frac{3}{4}\right) = \tan\theta$ 

if 
$$\sin 2\theta = \frac{3}{4}$$
 then  $\frac{2 \tan \theta}{1 + \tan^2 \theta} = \frac{3}{4}$   
 $8 \tan \theta = 3 + 3 \tan^2 \theta$   
 $3 \tan^2 \theta - 8 \tan \theta + 3 = 0$   
 $\tan \theta = \frac{8 \pm \sqrt{64 - 4 \times 3 \times 3}|}{6}$   
 $\tan \theta = \frac{8 \pm \sqrt{28}}{6} = \frac{4 \pm \sqrt{7}}{3}$   
 $\tan \theta = \frac{4 + \sqrt{7}}{3}$  or  $\frac{4 - \sqrt{7}}{3}$   
 $\tan \left(\frac{1}{2}\sin^{-1}\frac{3}{4}\right) = \frac{4 - \sqrt{7}}{3}$  Hence proved.  
**OR**  
 $\cos(\tan^{-1} x)$   
LHS. Irt  $\tan^{-1} x = \theta \Rightarrow x = \tan \theta$   
 $\cos \theta = \frac{1}{\sqrt{1 + \tan^2 \theta}} = \frac{1}{\sqrt{1 + x^2}}$   
Hence  $\cos(\tan^{-1} x) = \frac{1}{\sqrt{1 + x^2}}$   
Hence  $\cos(\tan^{-1} x) = \frac{1}{\sqrt{1 + x^2}}$   
R.H.S Let  $\cot^{-1}\frac{3}{4} = \theta \Rightarrow \frac{3}{4} = \cot \theta$   
then  $\sin \theta = \frac{1}{\sqrt{1 + \cot^2 \theta}} = \frac{1}{\sqrt{1 + \frac{9}{16}}} = \frac{4}{5}$   
Now LHS = RHS  
 $\frac{1}{\sqrt{1 + x^2}} = \frac{4}{5}$   
 $25 = 16 + 16x^2$   
 $x^2 = \frac{9}{16} \Rightarrow x = \frac{3}{4}$ 

13. Using properties of determinants, prove the following :

$$\begin{vmatrix} x & x+y & x+2y \\ x+2y & x & x+y \\ x+y & x+2y & x \end{vmatrix} = 9y^{2}(x+y)$$

$$\begin{vmatrix} x & x+y & x+2y \\ x+2y & x & x+y \\ x+y & x+2y & x \end{vmatrix} = 9y^{2}(x+y)$$
LHS 
$$\begin{vmatrix} x & x+y & x+2y \\ x+2y & x & x+y \\ x+2y & x & x+y \\ x+y & x+2y & x \end{vmatrix}$$
Now, apply C<sub>1</sub>  $\rightarrow$  C<sub>1</sub> + C<sub>2</sub> + C<sub>3</sub>  

$$\begin{vmatrix} 3x+3y & x+y & x+2y \\ 3x+3y & x & x+y \\ 3x+3y & x+2y & x \end{vmatrix}$$

$$3(x+y)\begin{vmatrix} 3x+3y & x+y & x+2y \\ 3x+3y & x+2y & x \end{vmatrix}$$

$$3(x+y)\begin{vmatrix} -y & 2y \\ -2y & y \end{vmatrix}$$

$$3(x+y)\begin{vmatrix} -y & 2y \\ -2y & y \end{vmatrix}$$

$$3y^{2}(x+y)\begin{vmatrix} -1 & 2 \\ -2 & 1 \end{vmatrix}$$

$$3y^{2}(x+y)(-1+4) = 9y^{2}(x+y).$$
 Hence proved.

٠

•

14. If 
$$y^{x} = e^{y-x}$$
, prove that  $\frac{dy}{dx} = \frac{(1+\log y)^{2}}{\log y}$   
Sol.  $y^{x} = e^{y-x}$   
 $\Rightarrow x \log_{e} y = y-x$  ....(1)  
Differentiating w.r.t.x  
 $\Rightarrow \log_{e} y + x \cdot \frac{1}{y} \frac{dy}{dx} = \frac{dy}{dx} - 1$   
 $\Rightarrow \log_{e} y + 1 = \frac{dy}{dx} \left(1 - \frac{x}{y}\right) \{form(1) \frac{x}{y} = \frac{1}{1 + \log_{e} y} \}$   
 $\Rightarrow \log_{e} y + 1 = \frac{dy}{dx} \left(1 - \frac{1}{1 + \log_{e} y}\right)$   
 $\Rightarrow (\log_{e} y + 1) = \frac{dy}{dx} \left(1 - \frac{\log_{e} y}{1 + \log_{e} y}\right)$   
 $\Rightarrow \frac{dy}{dx} = \frac{(1 + \log_{e} y)^{2}}{\log_{e} y}$ 

15. Differentiate the following with respect to x :

$$\sin^{-1}\left(\frac{2^{x+1} \cdot 3^x}{1+(36)^x}\right)$$

Solution:

$$y = \sin^{-1} \left[ \frac{2^{x+1} \cdot 3^x}{1 + (36)^x} \right]$$
$$y = \sin^{-1} \left[ \frac{2^x \cdot 2 \cdot 3^x}{1 + (36)^x} \right]$$
$$y = \sin^{-1} \left[ \frac{2 \cdot (6)^x}{1 + (6)^{2x}} \right]$$
$$y = 2 \tan^{-1} (6)^x$$
$$\frac{dy}{dx} = \frac{2}{1 + (6)^{2x}} \cdot 6^x \log 6$$
$$\frac{dy}{dx} = \frac{2 \cdot 6^x \log 6}{1 + (36)^x}$$

16. Find the value of k, for which 
$$f(x) = \begin{cases} \frac{\sqrt{1+kx} - \sqrt{1-kx}}{x}, & \text{if } -1 \le x < 0\\ \frac{x}{x-1}, & \text{if } 0 \le x < 1 \end{cases}$$
  
*OR*  
 $d^2 x = \pi$ 

.

If  $x = a \cos^3\theta$  and  $y = a \sin^3\theta$ , then find the value of  $\frac{d^2y}{dx^2}$  at  $\theta = \frac{\pi}{6}$ .

Solution:

$$f(x) = \begin{cases} \frac{\sqrt{1+kx} - \sqrt{1-kx}}{x} & \text{if } -1 \le x < 0\\ \frac{x}{x-1}, & \text{if } 0 \le x < 1 \end{cases}$$

function f(x) is continuous at x = 0

$$\therefore f(0) = \lim_{x \to 0} f(x)$$

$$\Rightarrow \frac{0+1}{0-1} = \lim_{x \to 0} \left( \frac{\sqrt{1+kx} - \sqrt{1-kx}}{x} \right)$$

$$\Rightarrow -1 = \lim_{x \to 0} \left( \frac{\sqrt{1+kx} - \sqrt{1-kx}}{x} \right) \left( \frac{\sqrt{1+kx} + \sqrt{1-kx}}{\sqrt{1+kx} + \sqrt{1-kx}} \right)$$

$$\Rightarrow -1 = \lim_{x \to 0} \frac{(1+2k) - (1-kx)}{x[\sqrt{1+kx} + \sqrt{1-kx}]}$$

$$\Rightarrow -1 = \lim_{x \to 0} \frac{2k}{\sqrt{1+kx} + \sqrt{1-kx}}$$

$$\Rightarrow -1 = \frac{2k}{2} \Rightarrow k = -1$$

OR

•

$$x = a \cos^{3} \theta \quad \text{and} \quad y = a \sin^{3} \theta$$
$$\frac{dx}{d\theta} = -3a \cos^{2\theta} \sin \theta \text{ and} \quad \frac{dy}{d\theta} = 3a \sin^{2} \theta \cos \theta$$
$$x = a \cos^{3} \theta \quad \text{and} \quad y = a \sin^{3} \theta$$
$$\frac{dx}{d\theta} = -3a \cos^{2\theta} \sin \theta \text{ and} \quad \frac{dy}{d\theta} = 3a \sin^{2} \theta \frac{\cos \theta}{\cos \theta}$$
$$\Rightarrow \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}}$$
$$\Rightarrow \frac{dy}{dx} = -\tan \theta$$
$$\Rightarrow \frac{d^{2}y}{dx^{2}} = -\sec^{2} \theta \frac{1}{(-3a \cos^{2} \theta . \sin \theta)}$$
$$\Rightarrow \frac{d^{2}y}{dx^{2}} = \frac{1}{3a} \sec^{4} \theta \csc \theta$$
$$\left(\frac{d^{2}y}{dx^{2}}\right)_{\theta\frac{\pi}{6}} = \frac{1}{3a} \left(\frac{2}{\sqrt{3}}\right)^{4} . 2 = \frac{32}{27a}$$

17. Evaluate :

·

$$\int \frac{\cos 2x - \cos 2\alpha}{\cos x - \cos \alpha}$$
*OR*

Evaluate :

$$\int \frac{x+2}{\sqrt{x^2+2x+3}} dx$$

$$\int \frac{\cos 2x - \cos 2\alpha}{\cos x - \cos \alpha} dx$$
  
=  $\int \frac{(2\cos^2 x - 1) - (2\cos^2 \alpha - 1)}{\cos x - \cos \alpha} dx$   
=  $\int \frac{2(\cos^2 x - 1)(2\cos^2 \alpha - 1)}{\cos x - \cos \alpha} dx$   
=  $2\int (\cos x + \cos \alpha) dx$   
=  $2(\sin x + x \cos \alpha) + c$ 

$$I = \int \frac{x+2}{\sqrt{x^2+2x+3}} dx$$

$$I = \int \left\{ \frac{(x+1)+1}{\sqrt{x^2+2x+3}} \right\} dx$$

$$I = \int \left\{ \frac{(x+1)}{\sqrt{x^2+2x+3}} \right\} dx + \int \left\{ \frac{1}{\sqrt{x^2+2x+3}} \right\} dx$$

$$I = I_1 + I_2$$
In I<sub>1</sub>let x<sup>2</sup> + 3 = t<sup>2</sup>  
 $\therefore$  (2x + 2)dx = 2t 2t  
 $\Rightarrow$  (x + 1)dx = tdt  
 $\therefore$  I<sub>1</sub> =  $\int \frac{t.dt}{t} = t$   
I<sub>1</sub> =  $\sqrt{x^2+2x+3}$   
Now in I<sub>1</sub> =  $\int \frac{1}{\sqrt{x^2+2x+3}} dx = \int \frac{dx}{\sqrt{x^2+2x+3}}$   
I<sub>2</sub> = log[(x + 1)<sup>2</sup> +  $\sqrt{(x + 1)^2 + 2}$ ]  
Now I = I<sub>1</sub> + I<sub>2</sub>  
 $\Rightarrow$  I =  $\sqrt{x^2+2x+3} + \log(x+1+\sqrt{x^2+2x+3}) + c$ 

18. Evaluate :

$$\int \frac{\mathrm{d}x}{x(x^5+3)}$$

$$I = \int \frac{dx}{x(x^5 + 3)}$$

$$I = \int \frac{x^4 dx}{x(x^5 + 3)}$$
Let  $x^5 = t \Longrightarrow 5x^4 dx = dt$ 

$$I = \frac{1}{5} \int \frac{dt}{t(t + 3)}$$

$$I = \frac{1}{5} \cdot \frac{1}{3} \int \left(\frac{1}{t} - \frac{1}{t + 3}\right) dt$$

$$I = \frac{1}{15} \{\log t - \log(t + 3)\} + c$$

$$I = \frac{1}{15} \log \int \left(\frac{t}{t + 3}\right) + c$$

$$I = \frac{1}{15} \log \int \left(\frac{x^5}{x^5 + 3}\right) + c$$

19. Evaluate

$$\int_0^{2\pi} \frac{1}{1+e^{\sin x}} dx$$

Solution:

$$I = \int_{0}^{2\pi} \frac{1}{1 + e^{\sin x}} dx \qquad \dots (1)$$
$$I = \int_{0}^{2\pi} \frac{1}{1 + e^{\sin(2\pi - x)}} dx$$
$$I = \int_{0}^{2\pi} \frac{1}{1 + e^{-\sin x}} dx$$
$$I = \int_{0}^{2\pi} \frac{e^{\sin x}}{e^{\sin x} + 1} dx$$
Adding (1) & (2) we get
$$\Rightarrow 21 = \int_{0}^{2\pi} \left(\frac{1 + e^{\sin x}}{1 + e^{\sin x}}\right) dx$$
$$\Rightarrow 21 = [x]_{0}^{2\pi}$$
$$\Rightarrow 21 = 2\pi$$
$$I = \pi$$

20. If  $\vec{a} = \vec{i} - \vec{j} + 7k$  and  $\vec{b} = 5\vec{i} + \vec{j} + \lambda \vec{k}$ , then find the value of  $\lambda$ , so that  $\vec{a} + \vec{b}$  and  $\vec{a} - \vec{b}$  are perpendicular vectors. Solution:

$$\vec{a} = \vec{i} - \vec{j} + 7k$$
  

$$\vec{b} = 5\vec{i} + \vec{j} + \lambda\vec{k}$$
  

$$\vec{a} + \vec{b} = 6\vec{i} - 2\vec{j} + (7 + \lambda)\vec{k}$$
  

$$\vec{a} - \vec{b} = 4\vec{i} - 0\vec{j} + (7 - \lambda)\vec{k}$$
  
given  $(\vec{a} + \vec{b})$  and  $(\vec{a} - \vec{b})$  are perpendicular  
 $\therefore (\vec{a} + \vec{b}).(\vec{a} - \vec{b}) = 0$   
 $\{6\vec{i} - 2\vec{j} + (7 + \lambda)\vec{k}\} \times \{4\vec{i} - 0\vec{j} + (7 - \lambda)\vec{k}\} = 0$   
 $6(-4) + 0(-2) + (7 + \lambda)(7 - \lambda) = 0$   
 $-24 + 49 - \lambda^2 = 0$   
 $\lambda^2 = 25 \Rightarrow [\lambda = \pm 5]$ 

21. Show that the lines

$$\vec{r} = 3\vec{i} + 2\vec{j} - 4\vec{k} + \lambda(\vec{i} + 2\vec{j} + 2\vec{k});$$
  
$$\vec{r} = 3\vec{i} + 2\vec{j} + \mu(3\vec{i} + 2\vec{j} + 6\vec{k})$$

are intersecting. Hence find their point of intersection.

#### **OR**

Find the vector equation of the plane through the points (2, 1, -1) and (-1, 3, 4) and perpendicular to the plane x - 2y + 4z = 10.

Sol. If the given lines are intersecting then the shortest distance between the lines is zero and also they have same

common point 
$$\vec{r} = 3\vec{i} + 2\vec{j} - 4\vec{k} + \lambda(\vec{i} + 2\vec{j} + 2\vec{k})$$
  

$$\Rightarrow \frac{x-3}{1} = \frac{y-2}{2} = \frac{z+4}{2} (=\lambda)(\text{Let})$$
Let P is  $(\lambda + 3, 2\lambda + 2, 2\lambda - 4)$   
Also,  $\vec{r} = 5\vec{i} - 2\vec{j} + \mu(3\vec{i} + 2\vec{j} + 6\vec{k})$   

$$\Rightarrow \frac{x-5}{3} = \frac{y+2}{2} = \frac{z-0}{6} (=\mu)(\text{Let})$$
Let Q is  $(\mu + 5, 2\mu - 2, 6\mu)$ 

If lines are intersecting then P and Q will be same.

$$\lambda + 3 = 3\mu + 5$$
 ....(1)  
 $2\lambda + 2 = 2\mu - 2$  ....(2)  
 $2\lambda - 4 = 6\mu$  ....(3)

Solve(2) & (3)  

$$\lambda + 1 = \mu - 1$$
  
 $2\lambda - 2 = 3\mu$   
 $\frac{- + -}{3 = 2\mu - 1}$   
 $4 = 2\mu$   
 $\mu = -2$   
put  $\mu = -2$  ....(3)  
 $2\lambda - 4 = 6(-2)$   
 $2\lambda = -12 + 4$   
 $2\lambda = -8$   
 $\lambda = -4$   
put  $\mu \& \lambda in (1)$   
 $\lambda + 3 = 3\mu + 5$   
 $-4 + 3 = 3(-2) + 5$   
 $-1 = -1$   
 $\therefore$  from  $\lambda = -4$  then P is (-1, -6, -12)  
from  $\mu = -2$  then Q is (-1, -6, -12)  
as P and Q are same

: lines are intersecting lines and their point of intersection is (-1, -6, -12).

OR



The required plane is perpendicular to given plane.

Therefore n r of required plane will be perpendicular to  $\vec{n}_1$  and AB.

$$\therefore \vec{n} \parallel (\vec{n}_{1} \times \vec{AB}) \vec{n}_{1} = \hat{i} - 2\hat{j} + 4k \vec{AB} = -3\hat{i} + 2\hat{j} - 5k \therefore \vec{n}_{1} \times \vec{AB} = -18\hat{i} - 17\hat{j} - 4k \therefore required plane is \vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n} \vec{r} \cdot (-18\hat{i} - 17\hat{j} - 4k) = (2\hat{i} + \hat{j} - k) \cdot (-18\hat{i} - 17\hat{j} - 4k) \vec{r} \cdot (-18\hat{i} - 17\hat{j} - 4k) = -36 - 17 + 4 \vec{r} \cdot (-18\hat{i} - 17\hat{j} - 4k) = -49 \boxed{18x + 17y + 4z = 49}$$

22. The probabilities of two students A and B coming to the school in time are  $\frac{3}{7}$  and  $\frac{5}{7}$  respectively.

Assuming that the events, 'A coming in time' and 'B coming in time' are independent, find the probability of only one of them coming to the school in time. Write at least one advantage of coming to school in time. Sol. If P(A come in school time) = 3/7

P (B come in school time) = 5/7 P (A not come in school time) = 4/7 P (B not come in school time) = 2/7 P (only one of them coming school in time) = P(A)×P( $\vec{B}$ )+P( $\vec{A}$ ).P(B) 3 2 5 4 26

$$=\frac{3}{7}\times\frac{2}{7}+\frac{5}{7}\times\frac{4}{7}=\frac{26}{49}$$

23. Find the area of the greatest rectangle that can be inscribed in an ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ 

**O**R

Find the equations of tangents to the curve  $3x^2 - y^2 = 8$ , which pass through the point  $\left(\frac{4}{3}, 0\right)$ 



Area of reactangle  $A = 2a = \cos \theta$ .  $2b \sin \theta$  A = 2ab.  $\sin 2\theta$  $\therefore A_{max} = 2ab$ 

**OR** 

Let a point 
$$(x_1, y_1)$$

$$3x^{2} - y^{2} = 8 \Longrightarrow 6x - 2y \cdot y' = 0$$
  

$$\Rightarrow y' = \frac{3x}{y}$$
  

$$\therefore \text{ Tangent } y - y_{1} = \frac{3x_{1}}{y_{1}}(x - x_{1})$$
  
It passing throgh  $\left(\frac{4}{3}, 0\right)$   

$$-y_{1} = \frac{3x_{1}}{y_{1}}\left(\frac{4}{3} - x_{1}\right)$$
  

$$\Rightarrow -y_{1}^{2} = 4x_{1} - 3x_{1}^{2} \implies y_{1}^{2} = 4x_{1} - 3x_{1}^{2}$$
  

$$\Rightarrow 3x_{1}^{2} - 8 = 3x_{1}^{2} - 4x_{1}$$
  

$$\therefore x_{1} = 2$$
  
So  $12 - y^{2} = 8$   

$$\Rightarrow y^{2} = 4 \implies y_{1} = \pm 2$$

24. Find the area of the region bounded by the parabola  $y = x^2$  and y = |x|. Sol.



Required area = 2[area of  $\triangle OAB$  – Area of curve OCBA]

$$A = 2\left[\frac{1}{2}(1)(1) - \int_0^1 x^2 dx\right]$$
$$A = 2\left[\frac{1}{2} - \frac{1}{3}\right] \implies A = 2\left[\frac{1}{6}\right] = \frac{1}{3}$$

25. Find the particular solution of the differential equation  $(tan-1y - x)dy = (1 + y^2)dx$ , given that when x = 0, y=0

Sol. 
$$(\tan^{-1}y - x)dy = (1 + y^2)dx$$
  

$$\Rightarrow \frac{dx}{dy} = \frac{\tan^{-1}y}{1 + y^2} - \frac{x}{1 + y^2}$$

$$\Rightarrow \frac{dx}{dy} + \frac{x}{1 + y^2} = \frac{\tan^{-1}y}{1 + y^2}$$

$$IF = e^{\int \frac{1}{1 + y^2} dy}$$

$$IF = e^{\tan^{-1}y}$$

$$x. IF = \int Q.IF dy + c$$

$$\Rightarrow x. e^{\tan^{-1}y} = \int \frac{\tan^{-1}y}{(1 + y^2)} e^{\tan^{-1}y} dy + c$$
Put  $\tan^{-1}y = t$ 

$$\Rightarrow x. e^{\tan^{-1}y} = (t.e^t) - (e^t) + c$$

$$\Rightarrow x. e^{\tan^{-1}y} = \tan^{-1}y. e^{\tan^{-1}y} - e^{\tan^{-1}y} + c$$

26. Find the equation of the plane passing through the line of intersection of the planes  $\vec{r}.(\hat{i}+3\hat{j})-6=0$  and  $\vec{r}.(3\hat{j}-\hat{j}-4k)=0$ , whose perpendicular distance from origin is unity.

#### OR

Find the vector equation of the line passing through the point (1, 2, 3) and parallel to the planes  $\vec{r}.(3\hat{i}-\hat{j}-2k)=5$  and  $\vec{r}.(3\hat{i}+\hat{j}+k)=6$ 

Sol.

P<sub>1</sub>is x + 3y - 6 = 0 P<sub>2</sub>is  $\vec{r} \cdot (3\hat{j} - \hat{j} - 4k) = 0$ P<sub>2</sub>is 3x - y - 4z = 0

 $P_1$  is  $\vec{r}.(3\hat{i}+\hat{j})-6=0$ 

Equation of plane passing through intersection of  $P_1$  and  $P_2$  is  $P_1 + \lambda P_2 = 0$ 

$$(x+3y-6) + \lambda(3x-y-4z) = 0$$
  
(1+3\lambda)x+(3-\lambda)y+(-4\lambda)z+(-6) = 0

Its distance from (0, 0, 0) is 1.

$$\begin{vmatrix} 0+0+0-6\\ \hline \sqrt{(1+3\lambda)^2+(3-\lambda)^2+(-4\lambda)^2} \end{vmatrix}$$
  
36 = (1 + 3\lambda)^2 + (3 - \lambda)^2 + (-4\lambda)^2  
36 = 1 + 9\lambda^2 + 6\lambda + 9 + \lambda^2 - 6\lambda + 16\lambda^2  
36 = 26\lambda^2 + 10 \Rightarrow 26\lambda^2 = 26 \Rightarrow \lambda^2 = 1 \Rightarrow \lambda = \pm 1

Hence required plane is

For  $\lambda = 1$ , (x + 3y - 6) + 1 (3x - y - 4z) = 04x + 2y - 4z - 6 = 0

For 
$$\lambda = -1$$
,  $(x + 3y - 6) - 1(3x - y - 4z) = 0$   
 $-2x + 4y + 4z - 6 = 0$   
OR  
P<sub>1</sub>is  $\vec{r} \cdot (\hat{i} - \hat{j} + 2k) = 5$   
 $\therefore \vec{n_1} = \hat{i} - \hat{j} + 2k$   
P<sub>2</sub>is  $\vec{r} \cdot (3\hat{i} - \hat{j} + 2k) = 6$   
 $\vec{n_2} = 3\hat{i} - \hat{j} + k$ 

The line parallel to plane P<sub>1</sub> & P<sub>2</sub> will be perpendicular to  $\vec{n}_1$  &  $\vec{n}_2$ 

$$\therefore \quad b \parallel (n_1 \times n_2)$$
  

$$\vec{n}_1 = \hat{i} - \hat{j} + 2k$$
  

$$\vec{n}_2 = 3\hat{i} - \hat{j} + k$$
  

$$\vec{n}_1 \times \vec{n}_2 = -3\hat{i} + 5\hat{j} + 4k$$
  

$$\therefore \quad \vec{b} = -3\hat{i} + 5\hat{j} + 4k$$
  
Point is (1, 2, 3)  

$$\therefore \quad \vec{a} = \hat{i} - \hat{j} + 3k$$
  

$$\therefore \quad \text{required line is } \vec{r} = \vec{a} \mid +\lambda\vec{b}$$

 $\vec{r} = (\hat{i} + 2\hat{j} + 3k) + \lambda(-3\hat{i} + 5\hat{j} + 4|k)$ 

27. In a hockey match, both teams A and B scored same number of goals up to the end of the game, so to decide the winner, the referee asked both the captains to throw a die alternately and decided that the team, whose captain gets a six first, will be declared the winner. If the captain of team A was asked to start, find the irrespective probabilities of winning the match and state whether the decision of the referee was fair or not.

Sol. P(6 get) = 
$$1/6$$
  
P(6 not get) =  $P^{\overline{(6get)}} = 5/6$   
P.(A win) = P(Aget6) + P(\overline{6get}).P(\overline{6get})P(\overline{6get}) + P(\overline{6get}).P(\overline{6get}).P(\overline{6get}).P(\overline{6get})P(\overline{6get}) + .....+\infty  
P.(A win) =  $\frac{1}{6} \times \frac{5}{6} \times \frac{5}{6} \times \frac{5}{6} \times \frac{5}{6} \times \frac{5}{6} \times \frac{5}{6} \times \frac{1}{6} + ... + \infty$   
=  $\frac{1}{6} + \left(\frac{5}{6}\right)^2 \times \frac{1}{6} + \left(\frac{5}{6}\right)^4 \times \frac{1}{6} + ... + \infty$   
 $\boxed{\therefore S_{\infty} = \frac{a}{1-r}}$   
=  $\frac{\left(\frac{1}{6}\right)}{\left(1-\frac{25}{36}\right)} = \frac{36}{11 \times 6} = \frac{6}{11}$ 

Similarly winning for B

$$P(B \text{ win}) = 1 - P(A \text{ win})$$
  
=  $1 - \frac{6}{11} = \frac{5}{11}$ 

28. A manufacturer considers that men and women workers are equally efficient and so he pays them at the same rate. He has 30 and 17 units of workers (male and female) and capital respectively, which he uses to produce two types of goods A and B. To produce one unit of A, 2 workers and 3 units of capital are required while 3workers and 1 unit of capital is required to produce one unit of B. If A and B are priced at j- 100 and j- 120per unit respectively, how should he use his resources to maximise the total revenue ? Form the above as an LPP and solve graphically. Do you agree with this view of the manufacturer that men and women workers are equally efficient and so should be paid at the same rate ?

Sol. if $z_{max} = 100x + 120y$						
		typeA	typeB			
work	ker	2	3	30		
capit	:1	3	1	17		
Subject to,						
2x+3y≤30						
3x+y≤17						
x≥0						
Let object of type $A = x$						
Object of type B = y						
y/1						
(0, 1/)						
(0, 10) C E(3, 8)						
В						
(0, 0) A (15, 0)						
$(2\mathbf{x} + 3\mathbf{y} \le 30)$						
$(3x + y) \le 17$						
pts	c00	rdinate	Z <sup>max</sup> =100x	x+ 120	у	
0	(0,0)		Z=0			
А	A $\begin{pmatrix} 17 \\ 0 \end{pmatrix}$		-1700			
$\left(\overline{3},0\right)$		-,0	$2-\overline{3}$			
Е	(3.8)		Z=300+960=1260			
С	(0.1	0)	Z=1200			
			1.0.10			

maximum revenue = 1260.

29. The management committee of a residential colony decided to award some of its members (say x) for honesty, some (say y) for helping others and some other (say z) for supervising the workers to keep the colony neat and clean. The sum of all the awardees is 12. Three times the sum of awardees for cooperation

and supervision added to two times the number of awardees for honesty is 33. If the sum of the number of awardees for honesty and supervision is twice the number of awardees for helping others, using matrix method, find the number of awardees of each category. Apart from these value, namely, honesty, cooperation and supervision, suggest one more value which the management of the colony must include for awards.

# Sol. Given $x + y + z = 12 \dots(1)$ $3(y+z) + 2x = 33 \dots (2)$ $(x + z) = 2y \dots (3)$ x + y + z = 122x + 3y + 3z = 33x - 2y + z = $\begin{bmatrix} 1 & 1 & 1 \\ 2 & 3 & 3 \\ 1 & -2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 12 \\ 33 \\ 0 \end{bmatrix}$ AX = BA - 1(AX) = A - 1(B) $I \cdot X = A - 1 \cdot B$ X = A - 1.B $\mathbf{X} = \frac{(\mathrm{Adj.A}) \cdot \mathbf{B}}{|\mathbf{A}|}$ $|\mathbf{A}| = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 3 & 3 \\ 1 & -2 & 1 \end{bmatrix}$ |A| = 1(3+6) - 1(2-3) + 1(-4-3)|A| = 9 + 1 - 7 = 3 $|\mathbf{A}| \neq 0$ $(\text{Adj. A}) = \begin{bmatrix} 9 & -3 & 0 \\ 1 & 0 & -1 \\ -7 & 3 & 1 \end{bmatrix}$ $(Adj. A).B = \begin{bmatrix} 9 & -3 & 0 \\ 1 & 0 & -1 \\ -7 & 3 & 1 \end{bmatrix}_{3\times 3} \begin{bmatrix} 12 \\ 33 \\ 0 \end{bmatrix}_{3\times 3}$ $(Adj. A).B = \begin{bmatrix} 9\\12\\15 \end{bmatrix}$ $\therefore X = \frac{(Adj.A) \cdot B}{|A|}$

$$X = \frac{1}{3} \begin{bmatrix} 9\\12\\15 \end{bmatrix} \Rightarrow X = \begin{bmatrix} 3\\4\\5 \end{bmatrix} \Rightarrow \begin{bmatrix} x\\y\\z \end{bmatrix} = \begin{bmatrix} 3\\4\\5 \end{bmatrix}$$
$$x = 3, y = 4, z = 5.$$

•