# **SOLVED EXAMPLES**





**Sol.** Length of perpendicular from origin to the line  $x\sqrt{5} + 2y = 3\sqrt{5}$  is

$$OL = \frac{3\sqrt{5}}{\sqrt{(\sqrt{5})^2 + 2^2}} = \frac{3\sqrt{5}}{\sqrt{9}} = \sqrt{5}$$

Radius of the given circle =  $\sqrt{10}$  = OQ = OP

$$PQ = 2QL = 2\sqrt{OQ^2 - OL^2} = 2\sqrt{10 - 5} = 2\sqrt{5}$$
  
Thus area of  $\triangle OPQ = \frac{1}{2} \times PQ \times OL = \frac{1}{2} \times 2\sqrt{5} \times \sqrt{5} = 5$ 



**Ex. 4** Find the equations of the following curves in Cartesian form. Also, find the centre and radius of the circle  $x = a + c \cos \theta$ ,  $y = b + c \sin \theta$ 

Sol. We have : 
$$x = a + c \cos \theta$$
,  $y = b + c \sin \theta$   $\Rightarrow$   $\cos \theta = \frac{x - a}{c}$ ,  $\sin \theta = \frac{y - b}{c}$   
 $\Rightarrow \qquad \left(\frac{x - a}{c}\right)^2 + \left(\frac{y - b}{c}\right)^2 = \cos^2 \theta + \sin^2 \theta$   $\Rightarrow$   $(x - a)^2 + (y - b)^2 = c^2$   
Clearly, it is a circle with centre at (a, b) and radius c.

Ex. 5 If the straight line ax + by = 2;  $a, b \neq 0$  touches the circle  $x^2 + y^2 - 2x = 3$  and is normal to the circle  $x^2 + y^2 - 4y = 6$ , then find the values of a and b are respectively.

**Sol.** Given 
$$x^2 + y^2 - 2x = 3$$

 $\therefore$  centre is (1, 0) and radius is 2

Given 
$$x^2 + y^2 - 4y = 6$$

:. centre is (0, 2) and radius is  $\sqrt{10}$ . Since line ax + by = 2 touches the first circle

$$\therefore \qquad \frac{|a(1)+b(0)-2|}{\sqrt{a^2+b^2}} = 2 \qquad \text{or} \quad |(a-2)| = [2\sqrt{a^2+b^2}] \qquad \dots \dots \dots (i)$$

Also the given line is normal to the second circle. Hence it will pass through the centre of the second circle. a(0) + b(2) = 2 or 2b = 2 or b = 1

Putting this value in equation (i) we get  $|a-2| = 2\sqrt{a^2 + 1^2}$  or  $(a-2)^2 = 4(a^2+1)$ 

$$a^{2}+4-4a=4a^{2}+4$$
 or  $3a^{2}+4a=0$  or  $a(3a+4)=0$  or  $a=0, -\frac{4}{3}$   $(a \neq 0)$ 

- $\therefore$  values of a and b are  $\left(-\frac{4}{3}, 1\right)$ .
- **Ex.6** Find the equation of a circle having the lines  $x^2 + 2xy + 3x + 6y = 0$  as its normal and having size just sufficient to contain the circle x(x-4) + y(y-3) = 0.

**Sol.** Pair of normals are (x+2y)(x+3)=0

Normals are 
$$x + 2y = 0$$
,  $x + 3 = 0$ .

Point of intersection of normals is the centre of required circle i.e.  $C_1(-3, 3/2)$  and centre of given circle is  $C_2(2, 3/2)$ 



...

and radius  $r_2 = \sqrt{4 + \frac{9}{4}} = \frac{5}{2}$ Let  $r_1$  be the radius of required circle  $r_1 = C_1C_2 + r_2 = \sqrt{(-3-2)^2 + (\frac{3}{2} - \frac{3}{2})^2} + \frac{5}{2} = \frac{15}{2}$ Hence equation of required circle is  $x^2 + y^2 + 6x - 3y - 45 = 0$ **Ex.**7 Let a circle be given by 2x(x-a) + y(2y-b) = 0 $(a \neq 0, b \neq 0)$ Find the condition on a and b if two chords, each bisected by the x-axis, can be drawn to the circle from (a, b/2). The given circle is 2x(x-a) + y(2y-b) = 0Sol. or  $x^2 + y^2 - ax - by/2 = 0$ Let AB be the chord which is bisected by x-axis at a point M. Let its co-ordinates be M(h, 0). and  $S \equiv x^2 + y^2 - ax - by/2 = 0$ Equation of chord AB is  $T = S_1$ ...  $hx + 0 - \frac{a}{2}(x+h) - \frac{b}{4}(y+0) = h^{2} + 0 - ah - 0$ Since its passes through (a, b/2) we have  $ah - \frac{a}{2}(a+h) - \frac{b^2}{8} = h^2 - ah$  $h^2 - \frac{3 a h}{2} + \frac{a^2}{2} + \frac{b^2}{8} = 0$ Now there are two chords bisected by the x-axis, so there must be two distinct real roots of h. ....  $B^2 - 4AC > 0$  $\Rightarrow \qquad \left(\frac{-3a}{2}\right)^2 - 4.1 \cdot \left(\frac{a^2}{2} + \frac{b^2}{8}\right) > 0 \quad \Rightarrow \quad a^2 > 2b^2.$ Prove that the circles  $x^2 + y^2 + 2ax + c^2 = 0$  and  $x^2 + y^2 + 2by + c^2 = 0$  touch each other, if  $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{c^2}$ . **Ex.8** Given circles are  $x^2 + y^2 + 2ax + c^2 = 0$ Sol. ..... (i)  $x^2 + y^2 + 2by + c^2 = 0$ and ..... (ii)

Let  $C_1$  and  $C_2$  be the centres of circles (i) and (ii), respectively and  $r_1$  and  $r_2$  be their radii, then

$$C_1 = (-a, 0), C_2 = (0, -b), r_1 = \sqrt{a^2 - c^2}, r_2 = \sqrt{b^2 - c^2}$$

Here we find the two circles touch each other internally or externally.

For touch,  $|C_1 C_2| = |r_1 \pm r_2|$ 

or

$$\sqrt{(a^2 + b^2)} = \left| \sqrt{(a^2 - c^2)} \pm \sqrt{(b^2 - c^2)} \right|$$

On squaring  $a^2 + b^2 = a^2 - c^2 + b^2 - c^2 \pm 2\sqrt{(a^2 - c^2)}\sqrt{(b^2 - c^2)}$ 



or  $c^2 = \pm \sqrt{a^2 b^2 - c^2 (a^2 + b^2) + c^4}$ Again squaring,  $c^4 = a^2 b^2 - c^2 (a^2 + b^2) + c^4$ or  $c^2 (a^2 + b^2) = a^2 b^2$ or  $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{c^2}$ 

**Ex.9** Find the equation of the circle through the points of intersection of  $x^2 + y^2 - 1 = 0$ ,  $x^2 + y^2 - 2x - 4y + 1 = 0$  and touching the line x + 2y = 0.

Sol. Family of circles is 
$$x^2 + y^2 - 2x - 4y + 1 + \lambda(x^2 + y^2 - 1) = 0$$
  
 $(1 + \lambda) x^2 + (1 + \lambda) y^2 - 2x - 4y + (1 - \lambda) = 0$   
 $x^2 + y^2 - \frac{2}{1 + \lambda} x - \frac{4}{1 + \lambda} y + \frac{1 - \lambda}{1 + \lambda} = 0$   
Centre is  $\left(\frac{1}{1 + \lambda}, \frac{2}{1 + \lambda}\right)$  and radius  $= \sqrt{\left(\frac{1}{1 + \lambda}\right)^2 + \left(\frac{2}{1 + \lambda}\right)^2 - \frac{1 - \lambda}{1 + \lambda}} = \frac{\sqrt{4 + \lambda^2}}{|1 + \lambda|}$ 

Since it touches the line x + 2y = 0, Hence Radius = Perpendicular distance from centre to the line.

i.e., 
$$\left| \frac{\frac{1}{1+\lambda} + 2\frac{2}{1+\lambda}}{\sqrt{1^2 + 2^2}} \right| = \frac{\sqrt{4+\lambda^2}}{|1+\lambda|} \implies \sqrt{5} = \sqrt{4+\lambda^2} \implies \lambda = \pm 1$$

 $\lambda = -1$  cannot be possible in case of circle. So  $\lambda = 1$ .

Thus, we get the equation of circle.

- **Ex. 10** Find the pole of the line 3x + 5y + 17 = 0 with respect to the circle  $x^2 + y^2 + 4x + 6y + 9 = 0$
- **Sol.** Given circle is  $x^2 + y^2 + 4x + 6y + 9 = 0$

and given line is 3x + 5y + 17 = 0

Let  $P(\alpha, \beta)$  be the pole of line (ii) with respect to circle (i)

Now equation of polar of point  $P(\alpha, \beta)$  with respect to circle (i) is

$$x\alpha + y\beta + 2(x + \alpha) + 3(y + \beta) + 9 = 0$$

or  $(\alpha + 2)x + (\beta + 3)y + 2\alpha + 3\beta + 9 = 0$ 

Now lines (ii) and (iii) are same, therefore,

$$\frac{\alpha + 2}{3} = \frac{\beta + 3}{5} = \frac{2\alpha + 3\beta + 9}{17}$$
(i) (ii) (iii)

From (i) and (ii), we get  $5\alpha + 10 = 3\beta + 9$  or  $5\alpha - 3\beta = -1$  ..... (iv) From (i) and (iii), we get  $17\alpha + 34 = 6\alpha + 9\beta + 27$  or  $11\alpha - 9\beta = -7$  ..... (v) Solving (iv) & (v), we get  $\alpha = 1, \beta = 2$ .

Hence required pole is (1, 2).



- Ex. 11 Find the equation of a circle which passes through the point (2, 0) and whose centre is the limit of the point of intersection of the lines 3x + 5y = 1 and  $(2 + c)x + 5c^2y = 1$  as  $c \rightarrow 1$ .
- **Sol.** Solving the equations  $(2 + c)x + 5c^2y = 1$  and 3x + 5y = 1

then  $(2+c)x+5c^2\left(\frac{1-3x}{5}\right)=1$  or  $(2+c)x+c^2(1-3x)=1$ 

...

$$x = \frac{1 - c^2}{2 + c - 3c^2}$$
 or  $x = \frac{(1 + c)(1 - c)}{(3c + 2)(1 - c)} = \frac{1 + c}{3c + 2}$ 

$$\therefore \qquad x = \lim_{c \to 1} \frac{1+c}{3c+2} \qquad \text{or} \quad x = \frac{2}{5}$$

$$\therefore \qquad y = \frac{1-3x}{5} = \frac{1-\frac{6}{5}}{5} = -\frac{1}{25}$$

Therefore the centre of the required circle is  $\left(\frac{2}{5}, \frac{-1}{25}\right)$  but circle passes through (2, 0)

:. Radius of the required circle = 
$$\sqrt{\left(\frac{2}{5}-2\right)^2 + \left(-\frac{1}{25}-0\right)^2} = \sqrt{\frac{64}{25} + \frac{1}{625}} = \sqrt{\frac{1601}{625}}$$

Hence the required equation of the circle is  $\left(x - \frac{2}{5}\right)^2 + \left(y + \frac{1}{25}\right)^2 = \frac{1601}{625}$ 

or 
$$25x^2 + 25y^2 - 20x + 2y - 60 = 0$$

- Ex. 12 The circle  $x^2 + y^2 4x 8y + 16 = 0$  rolls up the tangent to it at  $(2 + \sqrt{3}, 3)$  by 2 units, find the equation of the circle in the new position.
- **Sol.** Given circle is  $x^2 + y^2 4x 8y + 16 = 0$

let P =  $(2 + \sqrt{3}, 3)$ 

 $\sqrt{3} x - y - 2\sqrt{3} = 0$ 

Equation of tangent to the circle at  $P(2 + \sqrt{3}, 3)$  will be

$$(2 + \sqrt{3})x + 3y - 2(x + 2 + \sqrt{3}) - 4(y + 3) + 16 = 0$$

or

slope =  $\sqrt{3}$   $\Rightarrow$   $\tan\theta = \sqrt{3}$ 

 $\theta = 60^{\circ}$ line AB is parallel to the tangent at P

 $\Rightarrow$  coordinates of point B = (2 + 2cos60°, 4 + 2sin60°)

thus B = 
$$(3, 4 + \sqrt{3})$$

radius of circle =  $\sqrt{2^2 + 4^2 - 16} = 2$ 

equation of required circle is  $(x-3)^2 + (y-4-\sqrt{3})^2 = 2^2$ 

(2,4)

 $P(2+\sqrt{3},3)$ 



.



Sol. Let  $A \equiv (-a, 0)$  and  $B \equiv (a, 0)$  be two fixed points.

Let one line which rotates about B an angle  $\theta$  with the x-axis at any time t and at that time the second line which rotates about A make an angle  $2\theta$  with x-axis.

Now equation of line through B and A are respectively

 $y-0 = tan\theta(x-a)$  ...... (i) and  $y-0 = tan2\theta(x+a)$  ...... (ii)



Fro

⇒

$$y = \frac{1}{1 - \tan^2 \theta} (x + a)$$

$$= \left\{ \frac{\frac{2y}{(x - a)}}{1 - \frac{y^2}{(x - a)^2}} \right\} (x + a) \quad \text{(from (i))}$$

$$y = \frac{2y(x - a)(x + a)}{(x - a)^2 - y^2} \quad \Rightarrow \quad (x - a)^2 - y^2 = 2(x^2 - a^2)$$

or  $x^2 + y^2 + 2ax - 3a^2 = 0$  which is the required locus.

 $2 \tan \theta$ 

The circle  $x^2 + y^2 - 6x - 10y + k = 0$  does not touch or intersect the coordinate axes and the point (1, 4) is inside the Ex. 16 circle. Find the range of the value of k.

**O**(0,0)

B(a,0)

Since (1, 4) lies inside the circle Sol.

$$\Rightarrow$$
 S<sub>1</sub> < 0

$$\Rightarrow \qquad (1)^2 + (4)^2 - 6(1) - 10(4) + k < 0 \qquad \Rightarrow \qquad k < 29$$

Also centre of given circle is (3, 5) and circle does not touch or intersect the coordinate axes

$$\Rightarrow r < CA & r < CB$$

$$CA = 5$$

$$CB = 3$$

$$\Rightarrow r < 5 & r < 3$$

$$\Rightarrow r < 3 \text{ or } r^{2} < 9$$

$$r^{2} = 9 + 25 - k$$

$$r^{2} = 34 - k \Rightarrow 34 - k < 9$$

$$k > 25$$

$$\Rightarrow k \in (25, 29)$$

- Find the radical centre of circles  $x^2 + y^2 + 3x + 2y + 1 = 0$ ,  $x^2 + y^2 x + 6y + 5 = 0$  and  $x^2 + y^2 + 5x 8y + 15 = 0$ . Ex.17 Also find the equation of the circle cutting them orthogonally.
- Given circles are  $S_1 = x^2 + y^2 + 3x + 2y + 1 = 0$  $S_2 = x^2 + x^2 + x^2 + 3x + 2y + 1 = 0$ Sol.

$$S_2 \equiv x^2 + y^2 - x + 6y + 5 = 0$$
  
 $S_3 \equiv x^2 + y^2 + 5x - 8y + 15 = 0$ 

Equations of two radical axes are  $S_1 - S_2 \equiv 4x - 4y - 4 = 0$ 

x - y - 1 = 0or

and 
$$S_2 - S_3 \equiv -6x + 14y - 10 = 0$$

3x - 7y + 5 = 0or

Solving them the radical centre is (3, 2). Also, if r is the length of the tangent drawn from the radical centre (3, 2) to any one of the given circles, say S1, we have

$$\mathbf{r} = \sqrt{\mathbf{S}_1} = \sqrt{3^2 + 2^2 + 3.3 + 2.2 + 1} = \sqrt{27}$$

Hence (3, 2) is the centre and  $\sqrt{27}$  is the radius of the circle intersecting them orthogonally.

Its equation is  $(x-3)^2 + (y-2)^2 = r^2 = 27 \implies x^2 + y^2 - 6x - 4y - 14 = 0$ 



....

Find the equation of the circle of minimum radius which contains the three circles. **Ex. 18**  $S_1 \equiv x^2 + y^2 - 4y - 5 = 0$  $S_2 \equiv x^2 + y^2 + 12x + 4y + 31 = 0$  $S_3 \equiv x^2 + y^2 + 6x + 12y + 36 = 0$ Sol. For  $S_1$ , centre = (0, 2) and radius = 3 For  $S_2$ , centre = (-6, -2) and radius = 3 For  $S_3$ , centre = (-3, -6) and radius = 3 let P(a, b) be the centre of the circle passing through the centres of the three given circles, then  $(a-0)^{2}+(b-2)^{2}=(a+6)^{2}+(b+2)^{2}$  $(a+6)^2 - a^2 = (b-2)^2 - (b+2)^2$  $\Rightarrow$ (2a+6)6=2b(-4) $b = \frac{2 \times 6(a+3)}{-8} = -\frac{3}{2}(a+3)$ again  $(a-0)^2 + (b-2)^2 = (a+3)^2 + (b+6)^2$  $(a+3)^2 - a^2 = (b-2)^2 - (b+6)^2$ ⇒ (2a+3)3 = (2b+4)(-8) $(2a+3)3 = -16 \left| -\frac{3}{2}(a+3) + 2 \right|$ 6a + 9 = -8(-3a - 5)6a + 9 = 24a + 4018a = -31 $a = -\frac{31}{18}, b = -\frac{23}{12}$ radius of the required circle =  $3 + \sqrt{\left(-\frac{3}{18}\right)^2 + \left(-\frac{23}{12} - 2\right)^2} = 3 + \frac{5}{36}\sqrt{949}$ equation of the required circle is  $\left(x + \frac{31}{18}\right)^2 + \left(y + \frac{23}{12}\right)^2 = \left(3 + \frac{5}{36}\sqrt{949}\right)^2$ ... A fixed circle is cut by a family of circles all of which, pass through two given points  $A(x_1, y_1)$  and  $B(x_2, y_2)$ . Prove that **Ex. 19** the chord of intersection of the fixed circle with any circle of the family passes through a fixed point.

- Sol. Let S = 0 be the equation of fixed circle
  - let  $S_1 = 0$  be the equation of any circle through A and B

which intersect S = 0 in two points.

- $L = S S_1 = 0$  is the equation of the chord of intersection of S = 0 and  $S_1 = 0$
- let  $L_1 = 0$  be the equation of line AB

let  $S_2$  be the equation of the circle whose diametrical ends are  $A(x_1, y_1) \& B(x_2, y_2)$ 

then 
$$S_1 \equiv S_2 - \lambda L_1 = 0$$

$$L \equiv S - (S_2 - \lambda L_1) = 0$$
 or  $L \equiv (S - S_2) + \lambda L_1 = 0$   
 $L \equiv L' + \lambda L_1 = 0$  ......(i)

(i)

Implies each chord of intersection passes through the fixed point, which is the point of intersection of lines  $L' = 0 \& L_1 = 0$ .

Hence proved.



or

# Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141



- **Ex. 20** Let P be any moving point on the circle  $x^2 + y^2 2x = 1$ , from this point chord of contact is drawn w.r.t. the circle  $x^2 + y^2 2x = 0$ . Find the locus of the circumcentre of the triangle CAB, C being centre of the circle and A, B are the points of contact.
- **Sol.** The two circles are

$$(x-1)^2 + y^2 = 1$$
 ..... (i)  
 $(x-1)^2 + y^2 = 2$  ..... (ii)

So the second circle is the director circle of the first. So  $\angle APB = \pi/2$ 

Also 
$$\angle ACB = \pi/2$$

Now circumcentre of the right angled triangle CAB would lie on the mid point of AB So let the point be  $M \equiv (h, k)$ 

Now, 
$$CM = CBsin45^\circ = \frac{1}{\sqrt{2}}$$

So, 
$$(h-1)^2 + k^2 = \left(\frac{1}{\sqrt{2}}\right)^2$$

So, locus of M is 
$$(x - 1)^2 + y^2 = \frac{1}{2}$$



E	xercise # 1	Si	ngle Correct Choice	Type Questions]
1.	The lines $2x - 3y = 5$ and 3 (A) $x^2 + y^2 - 2x - 2y = 47$ (C) $x^2 + y^2 - 2x + 2y = 47$	x - 4y = 7 are diameters of a	a circle of area 154 sq. units. (B) $x^2 + y^2 - 2x - 2y = 62$ (D) $x^2 + y^2 - 2x + 2y = 62$	The equation of the circle is -
2.	Two lines through $(2, 3)$ (A) $2x + 3y = 13, x + 5y =$ (C) $x = 2, 9x - 11y = 51$	from which the circle x <sup>2</sup> - 17	$y^{2} = 25$ intercepts chord <b>(B)</b> $y = 3$ , $12x + 5y = 39$ <b>(D)</b> none of these	ds of length 8 units have equations
3.	The line $2x - y + 1 = x - 2y = 4$ . The radius of t	0 is tangent to the circle he circle is	e at the point (2, 5) and	the centre of the circles lies on
	<b>(A)</b> 3√5	<b>(B)</b> 5√3	( <b>C</b> ) 2√5	( <b>D</b> ) $5\sqrt{2}$
4.	$y = \sqrt{3}x + c_1 \& y = \sqrt{3}$ (A)8	$\mathbf{x} + \mathbf{c}_2$ are two parallel tar <b>(B)</b> 4	ngents of a circle of radius (C) 2	2 units, then $ c_1 - c_2 $ is equal to - (D) 1
5.	Let $C_1$ and $C_2$ are circles d segment PQ that is tangen (A) 15	defined by $x^2 + y^2 - 20x + 6$ at to C <sub>1</sub> at P and to C <sub>2</sub> at Q (B) 18	$54 = 0$ and $x^2 + y^2 + 30x + 14$ 0 is (C) 20	44 = 0. The length of the shortest line (D) 24
6.	The equation to the circle origin is - (A) $x^2 + y^2 - 6x + 8y - 9 =$ (C) $x^2 + y^2 + 6x \pm 8y + 9 =$	e whose radius is 4 and wh 0 0	hich touches the negative x (B) $x^2 + y^2 \pm 6x - 8y + 9$ (D) $x^2 + y^2 \pm 6x - 8y - 9$	x-axis at a distance 3 units from the = 0 = 0
7.	The angle between the t	wo tangents from the orig	gin to the circle $(x - 7)^2 + (x - 7)^2$	$(y+1)^2 = 25$ equals
	(A) $\frac{\pi}{4}$	(B) $\frac{\pi}{3}$	(C) $\frac{\pi}{2}$	(D) none
8.	The centre of the smalles (A) (3,2)	t circle touching the circles (B) (4,4)	$x^{2} + y^{2} - 2y - 3 = 0$ and $x^{2}$ (C) (2,7)	$y^2 + y^2 - 8x - 18y + 93 = 0$ is (D) (2,5)
9.	B and C are fixed points then the locus of the ce	having co-ordinates (3, 0 ntroid of the $\triangle ABC$ has	D) and $(-3, 0)$ respectively the equation -	. If the vertical angle BAC is 90°,
	(A) $x^2 + y^2 = 1$	<b>(B)</b> $x^2 + y^2 = 2$	(C) $9(x^2 + y^2) = 1$	<b>(D)</b> $9(x^2 + y^2) = 4$
10.	The condition so that the (A) $g^2 + f^2 = c + k^2$	e line $(x + g) \cos\theta + (y + g)$ (B) $g^2 + f^2 = c^2 + k$	f) $\sin \theta = k$ is a tangent to (C) $g^2 + f^2 = c^2 + k^2$	$x^{2} + y^{2} + 2gx + 2fy + c = 0$ is (D) $g^{2} + f^{2} = c + k$
11.	If the circle $C_1: x^2 + y^2 = 1$ maximum length and has	6 intersects another circl a slope equal to 3/4, then t	$e C_2$ of radius 5 in such a the co-ordinates of the cen	manner that the common chord is of tre of $C_2$ are
	(A) $\left(\pm\frac{9}{5},\pm\frac{12}{5}\right)$	<b>(B)</b> $\left(\pm\frac{9}{5}, \frac{12}{5}\right)$	(C) $\left(\pm\frac{12}{5},\pm\frac{9}{5}\right)$	<b>(D)</b> $\left(\pm\frac{12}{5}, \frac{9}{5}\right)$



# Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141

12. Number of different circles that can be drawn touching 3 lines, no two of which are parallel and they are neither coincident nor concurrent, are -

13. The area of an equilateral triangle inscribed in the circle  $x^2 + y^2 - 2x = 0$  is :

(A) 
$$\frac{3\sqrt{3}}{2}$$
 (B)  $\frac{3\sqrt{3}}{4}$  (C)  $\frac{3\sqrt{3}}{8}$  (D) none

14. The length of the tangent drawn from any point on the circle  $x^2 + y^2 + 2gx + 2fy + p = 0$  to the circle  $x^2 + y^2 + 2gx + 2fy + q = 0$  is:

(A) 
$$\sqrt{q-p}$$
 (B)  $\sqrt{p-q}$  (C)  $\sqrt{q+p}$  (D) none

15. The equations of the tangents drawn from the point (0,1) to the circle  $x^2 + y^2 - 2x + 4y = 0$  are -

(A) 
$$2x-y+1=0, x+2y-2=0$$
  
(B)  $2x-y-1=0, x+2y-2=0$   
(C)  $2x-y+1=0, x+2y+2=0$   
(D)  $2x-y-1=0, x+2y+2=0$ 

16. A circle of radius unity is centered at origin. Two particles start moving at the same time from the point (1, 0) and move around the circle in opposite direction. One of the particle moves counterclockwise with constant speed v and the other moves clockwise with constant speed 3v. After leaving (1, 0), the two particles meet first at a point P, and continue until they meet next at point Q. The coordinates of the point Q are (A) (1, 0) (B) (0, 1) (C) (0, -1) (D) (-1, 0)

17.The locus of the centers of the circles which cut the circles  $x^2 + y^2 + 4x - 6y + 9 = 0$  and<br/> $x^2 + y^2 - 5x + 4y - 2 = 0$  orthogonally is -<br/>(A) 9x + 10y - 7 = 0<br/>(C) 9x - 10y + 11 = 0(B) x - y + 2 = 0<br/>(D) 9x + 10y + 7 = 0

18. The parametric coordinates of any point on the circle  $x^2 + y^2 - 4x - 4y = 0$  are-(A)  $(-2 + 2\cos\alpha, -2 + 2\sin\alpha)$  (B)  $(2 + 2\cos\alpha, 2 + 2\sin\alpha)$ (C)  $(2 + 2\sqrt{2}\cos\alpha, 2 + 2\sqrt{2}\sin\alpha)$  (D)  $(-2 + 2\sqrt{2}\cos\alpha, -2 + 2\sqrt{2}\sin\alpha)$ 

- 19. The equation of the diameter of the circle  $(x 2)^2 + (y + 1)^2 = 16$  which bisects the chord cut off by the circle on the line x - 2y - 3 = 0 is (A) x + 2y = 0 (B) 2x + y - 3 = 0 (C) 3x + 2y - 4 = 0 (D) none
- 20. A line meets the co-ordinate axes in A and B. A circle is circumscribed about the triangle OAB. If  $d_1$  and  $d_2$  are the distances of the tangent to the circle at the origin O from the points A and B respectively, the diameter of the circle is

(A) 
$$\frac{2d_1 + d_2}{2}$$
 (B)  $\frac{d_1 + 2d_2}{2}$  (C)  $d_1 + d_2$  (D)  $\frac{d_1d_2}{d_1 + d_2}$ 

21. The gradient of the tangent line at the point (a  $\cos \alpha$ , a  $\sin \alpha$ ) to the circle  $x^2 + y^2 = a^2$ , is -

(A) 
$$\tan (\pi - \alpha)$$
 (B)  $\tan \alpha$  (C)  $\cot \alpha$  (D)  $- \cot \alpha$ 

22. The equation of normal to the circle  $x^2 + y^2 - 4x + 4y - 17 = 0$  which passes through (1, 1) is (A) 3x + y - 4 = 0 (B) x - y = 0 (C) x + y = 0 (D) none



23. The equation of the common tangent to the circle  $x^2 + y^2 - 4x - 6y - 12 = 0$  and

- $x^2 + y^2 + 6x + 18y + 26 = 0$  at their point of contact is (A) 12x + 5y + 19 = 0 (B) 5x + 12y + 19 = 0
- (C) 5x 12y + 19 = 0 (D) 12x 5y + 19 = 0
- 24. A pair of tangents are drawn from the origin to the circle  $x^2 + y^2 + 20(x + y) + 20 = 0$ . The equation of the pair of tangents is -

(A) 
$$x^2 + y^2 + 5xy = 0$$
  
(B)  $x^2 + y^2 + 10xy = 0$   
(C)  $2x^2 + 2y^2 + 5xy = 0$   
(D)  $2x^2 + 2y^2 - 5xy = 0$ 

25. In a right triangle ABC, right angled at A, on the leg AC as diameter, a semicircle is described. The chord joining A with the point of intersection D of the hypotenuse and the semicircle, then the length AC equals to

(A) 
$$\frac{AB \cdot AD}{\sqrt{AB^2 + AD^2}}$$
 (B)  $\frac{AB \cdot AD}{AB + AD}$  (C)  $\sqrt{AB \cdot AD}$  (D)  $\frac{AB \cdot AD}{\sqrt{AB^2 - AD^2}}$ 

26. Tangents are drawn from (4, 4) to the circle  $x^2 + y^2 - 2x - 2y - 7 = 0$  to meet the circle at A and B. The length of the chord AB is -

(A) 
$$2\sqrt{3}$$
 (B)  $3\sqrt{2}$  (C)  $2\sqrt{6}$  (D)  $6\sqrt{2}$ 

27. Pair of tangents are drawn from every point on the line 3x + 4y = 12 on the circle  $x^2 + y^2 = 4$ . Their variable chord of contact always passes through a fixed point whose co-ordinates are -

(A) 
$$\left(\frac{4}{3}, \frac{3}{4}\right)$$
 (B)  $\left(\frac{3}{4}, \frac{3}{4}\right)$  (C) (1, 1) (D)  $\left(1, \frac{4}{3}\right)$ 

- 28. The locus of the mid point of a chord of the circle  $x^2 + y^2 = 4$  which subtends a right angle at the origin is: (A) x+y=2 (B)  $x^2+y^2=1$  (C)  $x^2+y^2=2$  (D) x+y=1
- 29. The radical centre of three circles taken in pairs described on the sides of a triangle ABC as diameters is the :
   (A) centroid of the ΔABC
   (B) incentre of the ΔABC
   (D) orthocentre of the ΔABC

(C)  $\frac{\pi}{4}$ 

30. The angle between the two tangents from the origin to the circle  $(x-7)^2 + (y+1)^2 = 25$  equals -

(A) 
$$\frac{\pi}{2}$$

31. The number of common tangents of the circles  $x^2 + y^2 - 2x - 1 = 0$  and  $x^2 + y^2 - 2y - 7 = 0$ . (A) 1 (B) 3 (C) 2 (D) 4

(B)  $\cos^{-1}\frac{4}{5}$  (C)  $\frac{\pi}{2}$ 

**(B)**  $\frac{\pi}{3}$ 

32. Two circles are drawn through the points (1, 0) and (2, -1) to touch the axis of y. They intersect at an angle

(A) 
$$\cot^{-1}\frac{3}{4}$$

33. Equation of the circle cutting orthogonally the three circles  $x^2 + y^2 - 2x + 3y - 7 = 0$ ,  $x^2 + y^2 + 5x - 5y + 9 = 0$  and  $x^2 + y^2 + 7x - 9y + 29 = 0$  is (A)  $x^2 + y^2 - 16x - 18y - 4 = 0$ (B)  $x^2 + y^2 - 7x + 11y + 6 = 0$ (C)  $x^2 + y^2 + 2x - 8y + 9 = 0$ (D) none of these

# Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141

(D) none

**(D)** tan<sup>-1</sup> 1

# MATHS FOR JEE MAIN & ADVANCED

34.	4. A circle is drawn touching the x-axis and centre at the point which is the reflection of (a, b) in the $y - x = 0$ . The equation of the circle is -				
	(A) $x^2 + y^2 - 2bx - 2ay + a^2 = 0$	(B) $x^2 + y^2 - 2bx - 2ay + b^2 = 0$			
	(C) $x^2 + y^2 - 2ax - 2by + b^2 = 0$	( <b>D</b> ) $x^2 + y^2 - 2ax - 2by + a^2 = 0$			
35.	A rhombus is inscribed in the region cor $x^2 + y^2 + 4x - 12 = 0$ with two of its vertices on the line	mmon to the two circles $x^2 + y^2 - 4x - 12 = 0$ and poining the centres of the circles. The area of the rhombous is			
	(A) $8\sqrt{3}$ sq.units (B) $4\sqrt{3}$ sq.units	(C) $16\sqrt{3}$ sq.units (D) none			
36.	The equation of the circle having the lines $y^2 - 2y + 4x$	x - 2xy = 0 as its normals & passing through the point (2,1) is -			
	(A) $x^2 + y^2 - 2x - 4y + 3 = 0$	<b>(B)</b> $x^2 + y^2 - 2x + 4y - 5 = 0$			
	(C) $x^2 + y^2 + 2x + 4y - 13 = 0$	(D) none			
37.	The circumference of the circle $x^2 + y^2 - 2x + 8y - x^2 + y^2 + 4x + 12y + p = 0$ , then $p + q$ is equal to:	q = 0 is bisected by the circle			
	(A) 25 (B) 100	(C) 10 (D) 48			
38.	If the two circles, $x^2 + y^2 + 2g_1x + 2f_1y = 0$ and $x^2$	$f^2 + y^2 + 2g_2x + 2f_2y = 0$ touches each other, then -			
	(A) $f_1g_1 = f_2g_2$ (B) $\frac{f_1}{g_1} = \frac{f_2}{g_2}$	(C) $f_1 f_2 = g_1 g_2$ (D) none			
39.	The equation of a line inclined at an any	gle $\frac{\pi}{4}$ to the axis X, such that the two circles			
	$x^{2} + y^{2} = 4$ $x^{2} + y^{2} - 10x - 14x + 65 = 0$ interced	at equal lengths on it is			
	(A) $2x-2y-3=0$ (B) $2x-2y+3=0$	(C) $x-y+6=0$ (D) $x-y-6=0$			
40.	The distance between the chords of contact of tange	ents to the circle $x^2 + y^2 + 2gx + 2fy + c = 0$ from the origin			
	and the point (g, f) is				
	$\sqrt{g^2 + f^2 - c}$	$g^2 + f^2 - c$ $\sqrt{g^2 + f^2 + c}$			

(A) 
$$\sqrt{g^2 + f^2}$$
 (B)  $\frac{\sqrt{g^2 + f^2 - c}}{2}$  (C)  $\frac{g^2 + f^2 - c}{2\sqrt{g^2 + f^2}}$  (D)  $\frac{\sqrt{g^2 + f^2 + c}}{2\sqrt{g^2 + f^2}}$ 



# Exercise # 2 Part # I [Multiple Correct Choice Type Questions]

(D) Area of the circle touching these four circles internally is  $4\pi(3+2\sqrt{2})$ 



10. If r represent the distance of a point from origin &  $\theta$  is the angle made by line joining origin to that point from line xaxis, then  $r = |\cos\theta|$  represents -**(B)** two circles centered at  $\left(\frac{1}{2}, 0\right)$  &  $\left(-\frac{1}{2}, 0\right)$ (A) two circles of radii  $\frac{1}{2}$  each. (C) two circles touching each other at the origin. (D) pair of straight line Consider the circles  $S_1: x^2 + y^2 = 4$  and  $S_2: x^2 + y^2 - 2x - 4y + 4 = 0$  which of the following statements are correct? 11. (A) Number of common tangents to these circles is 2. (B) If the power of a variable point P w.r.t. these two circles is same then P moves on the line x + 2y - 4 = 0. (C) Sum of the y-intercepts of both the circles is 6. **(D)** The circles  $S_1$  and  $S_2$  are orthogonal. The equation of circles passing through (3, -6) touching both the axes is 12. **(B)**  $x^2 + y^2 + 6x - 6y + 9 = 0$ (A)  $x^2 + y^2 - 6x + 6y + 9 = 0$ (C)  $x^2 + y^2 + 30x - 30y + 225 = 0$ (D)  $x^2 + y^2 - 30x + 30y + 225 = 0$ Tangents are drawn to the circle  $x^2 + y^2 = 50$  from a point 'P' lying on the x-axis. These tangents meet the y-13. axis at points 'P<sub>1</sub>' and 'P<sub>2</sub>'. Possible co-ordinates of 'P' so that area of triangle  $PP_1P_2$  is minimum is/are -**(B)**  $(10\sqrt{2}, 0)$ (C)(-10,0)(**D**)  $(-10\sqrt{2}, 0)$ (A)(10,0)The equation of the circle which touches both the axes and the line  $\frac{x}{3} + \frac{y}{4} = 1$  and lies in the first quadrant 14. is  $(x - c)^2 + (y - c)^2 = c^2$  where c is **(A)** 1 **(B)**2(C)4**(D)** 6 If y = c is a tangent to the circle  $x^2 + y^2 - 2x + 2y - 2 = 0$ , then the value of c can be -(A) 1 (B) 3 (C) -1 15. **(D)**-3 A family of linear functions is given by f(x) = 1 + c(x+3) where  $c \in \mathbb{R}$ . If a member of this family meets a unit circle 16. centered at origin in two coincident points then 'c' can be equal to (A) -3/4**(B)**0 (C) 3/4 **(D)** 1 Equations of circles which pass through the points (1, -2) and (3, -4) and touch the x-axis is 17. (A)  $x^2 + y^2 + 6x + 2y + 9 = 0$ **(B)**  $x^2 + y^2 + 10x + 20y + 25 = 0$ (C)  $x^2 + y^2 - 6x + 4y + 9 = 0$ (D) none 18. The common chord of two intersecting circles  $C_1$  and  $C_2$  can be seen from their centres at the angles of 90° & 60° respectively. If the distance between their centres is equal to  $\sqrt{3}$  + 1 then the radii of C<sub>1</sub> and C<sub>2</sub> are -(B)  $\sqrt{2}$  and  $2\sqrt{2}$  (C)  $\sqrt{2}$  and 2 (D)  $2\sqrt{2}$  and 4 (A)  $\sqrt{3}$  and 3 If  $al^2 - bm^2 + 2 dl + 1 = 0$ , where a, b, d are fixed real numbers such that  $a + b = d^2$  then the line lx + my + 1 = 019. touches a fixed circle : (A) which cuts the x-axis orthogonally (B) with radius equal to b (C) on which the length of the tangent from the origin is  $\sqrt{d^2 - b}$ (D) none of these. Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035

20. The equation(s) of the tangent at the point (0, 0) to the circle, making intercepts of length 2a and 2b units on the co-ordinate axes, is (are) -**(C)** x = y (**D**) bx + ay = 0

(A) ax + by = 0**(B)** ax - by = 0

21. Consider the circles

$$S_1: x^2 + y^2 + 2x + 4y + 1 = 0$$
  

$$S_2: x^2 + y^2 - 4x + 3 = 0$$
  

$$S_3: x^2 + y^2 + 6y + 5 = 0$$

Which of this following statements are correct?

- (A) Radical centre of  $S_1$ ,  $S_2$  and  $S_3$  lies in  $1^{st}$  quadrant.
- **(B)** Radical centre of  $S_1$ ,  $S_2$  and  $S_3$  lies in 4<sup>th</sup> quadrants.
- (C) Radius of the circle intersecting  $S_1$ ,  $S_2$  and  $S_3$  orthogonally is 1.
- (D) Circle orthogonal to  $S_1$ ,  $S_2$  and  $S_3$  has its x and y intercept equal to zero.
- 22. Locus of the intersection of the two straight lines passing through (1, 0) and (-1, 0) respectively and including an angle of 45° can be a circle with

(A) centre (1, 0) and radius  $\sqrt{2}$ . (B) centre (1, 0) and radius 2. (C) centre (0, 1) and radius  $\sqrt{2}$ . (D) centre (0, -1) and radius  $\sqrt{2}$ .

```
[Assertion & Reason Type Questions]
Part # II
```

These questions contains, Statement-I (assertion) and Statement-II (reason).

- (A) Statement-I is true, Statement-II is true; Statement-II is correct explanation for Statement-I.
- (B) Statement-I is true, Statement-II is true; Statement-II is NOT a correct explanation for statement-I.
- (C) Statement-I is true, Statement-II is false.
- (D) Statement-I is false, Statement-II is true.

Consider two circles  $C_1 \equiv x^2 + y^2 + 2x + 2y - 6 = 0$  &  $C_2 \equiv x^2 + y^2 + 2x + 2y - 2 = 0$ . 1.

Two tangents are drawn from a point on the circle  $C_1$  to the circle  $C_2$ , then tangents always Statement-I: perpendicular.

 $C_1$  is the director circle of  $C_2$ . Statement-II:

Passing through a point A(6, 8) a variable secant line L is drawn to the circle 2.  $S: x^2 + y^2 - 6x - 8y + 5 = 0$ . From the point of intersection of L with S, a pair of tangent lines are drawn which intersect at P.

Statement - I	Locus of the point P has the equation $3x + 4y - 40 = 0$ .	
---------------	--	--

- Statement II Point A lies outside the circle.
- Only one normal can be drawn through the point P(2, -3) to the circle 3. Statement - I  $x^2 + y^2 - 4x + 8y - 16 = 0$
- Statement II Passing through any point lying inside a given circle only one normal can be drawn.
- 4. Statement - I If three circles which are such that their centres are non-collinear, then exactly one circle exists which cuts the three circles orthogonally.
  - Statement II Radical axis for two intersecting circles is the common chord.



# MATHS FOR JEE MAIN & ADVANCED

5.	Let C <sub>1</sub> denotes a family of circles with centre on x-axis and touching the y-axis at the origin.					
	and $C_2$ denotes a	and C <sub>2</sub> denotes a family of circles with centre on y-axis and touching the x-axis at the origin.				
	Statement - I	Every member of $C_1$ intersects any member of $C_2$ at right angles at the point other than origin.				
	Statement - II	If two circles intersect at 90° at one point of their intersection, then they must intersect at 90° on the other point of intersection also.				
6.	Let C be a circle the circle C at P a	with centre 'O' and HK is the chord of contact of pair of the tangents from point A. OA intersects and Q and B is the midpoint of HK, then				
	Statement - I	AB is the harmonic mean of AP and AQ.				
	Statement - II	AK is the Geometric mean of AB and AO and OA is the arithmetic mean of AP and AQ.				
7.	Statement - I	Angle between the tangents drawn from the point $P(13, 6)$ to the circle				
		$S: x^2 + y^2 - 6x + 8y - 75 = 0$ is 90°.				
	Statement - II	Point P lies on the director circle of S.				
8.	Consider the line	s				
	L:(k+'	7)x - (k-1)y - 4(k-5) = 0  where  k  is a parameter				
	and the circle					
	$C: x^2 + y^2$	$y^2 + 4x + 12y - 60 = 0$				
	Statement - I	Every member of L intersects the circle 'C' at an angle of 90°				
	Statement - II	Every member of L is tangent to the circle C.				
9.	Statement - I	The line $(x - 3)\cos\theta + (y - 3)\sin\theta = 1$ touches a circle $(x - 3)^2 + (y - 3)^2 = 1$ for all values of $\theta$ .				
	Statement - II	$x\cos\theta + y\sin\theta = a$ is a tangent of circle $x^2 + y^2 = a^2$ for all values of $\theta$ .				
10.	Statement - I	The circle $C_1: x^2 + y^2 - 6x - 4y + 9 = 0$ bisects the circumference of the circle				
		$C_2: x^2 + y^2 - 8x - 6y + 23 = 0.$				
	Statement - II	Centre of the circle $C_1$ lies on the circumference of $C_2$ .				
11.	Statement - I	The length of intercept made by the circle $x^2 + y^2 - 2x - 2y = 0$ on the x-axis is 2.				
	Statement - II	$x^{2} + y^{2} - \alpha x - \beta y = 0$ is a circle which passes through origin with centre $\left(\frac{\alpha}{2}, \frac{\beta}{2}\right)$ and				
		radius $\sqrt{\frac{\alpha^2+\beta^2}{2}}$ .				
12.	Consider the circ	les, $S_1 : x^2 + y^2 + 2x - 4 = 0$ and $S_2 : x^2 + y^2 - y + 1 = 0$				
	Statement - I	Tangents from the point $P(0, 5)$ on $S_1$ and $S_2$ are equal.				
	Statement - II	Point $P(0, 5)$ lies on the radical axis of the two circles.				
13.	A circle is circum Let $x = PA$ , $y = PI$	scribed about an equilateral triangle ABC and a point P on the minor arc joining A and B, is chosen. B and $z = PC$ . (z is larger than both x and y.)				

- **Statement I** Each of the possibilities (x + y) greater than z, equal to z or less than z, is possible for some P.
  - **Statement II** In a triangle ABC, sum of the two sides of a triangle is greater than the third and the third side is greater than the difference of the two.



# Exercise # 3 Part # I [Matrix Match Type Questions]

Following question contains statements given in two columns, which have to be matched. The statements in **Column-I** are labelled as A, B, C and D while the statements in **Column-II** are labelled as p, q, r and s. Any given statement in **Column-I** can have correct matching with **one or more** statement(s) in **Column-II**.

1.		Column-I			Column	-11
	<b>(A)</b>	If point of intersection and number of common tangents of	two		<b>(p)</b>	$\mu-\lambda=3$
		circles $x^2 + y^2 - 2x - 6y + 9 = 0$ and $x^2 + y^2 + 6x - 2y + 1 = 0$				
		are $\lambda$ and $\mu$ respectively, then				
	<b>(B)</b>	If point of intersection and number of tangents of two circle	es		<b>(q)</b>	$\mu+\lambda=5$
		$x^2+y^2-6x=0$ and $x^2+y^2+2x=0$ are $\lambda$ and $\mu$				
		respectively, then				
	<b>(C)</b>	If the straight line $y = mx \forall m \in I$ touches or lies outside			<b>(r)</b>	$\mu-\lambda=4$
		the circle $x^2 + y^2 - 20y + 90 = 0$ and the maximum and				
		minimum values of $ m $ are $\mu \& \lambda$ respectively then				
	<b>(D</b> )	If two circle $x^2 + y^2 + px + py - 7 = 0$ and			<b>(S)</b>	$\mu+\lambda=4$
		$x^2 + y^2 - 10x + 2py + 1 = 0$ cut orthogonally and				
		the value of p are $\lambda \& \mu$ respectively then				
2.		Column – I			Column	-II
	<b>(A)</b>	Number of values of a for which the common chord			<b>(p)</b>	4
		of the circles $x^2 + y^2 = 8$ and $(x - a)^2 + y^2 = 8$ subtends				
		a right angle at the origin is				
	<b>(B)</b>	A chord of the circle $(x-1)^2 + y^2 = 4$ lies along the			<b>(q)</b>	2
		line y = $22\sqrt{3}$ (x - 1). The length of the chord is equal to				
	<b>(C)</b>	The number of circles touching all the three lines			<b>(r)</b>	0
		3x + 7y = 2, $21x + 49y = 5$ and $9x + 21y = 0$ are				
	<b>(D</b> )	If radii of the smallest and largest circle passing through			<b>(s)</b>	1
		the point $(\sqrt{3}, \sqrt{2})$ and touching the circle				
		$x^2 + y^2 - 2 \sqrt{2} y - 2 = 0$ are r <sub>1</sub> and r <sub>2</sub> respectively, then				
		the mean of $r_1$ and $r_2$ is				
_						
3.		Column-I		Column-	. <b>]]</b> amman t	
	(A)	Two intersecting circles	(P)	nave a c		angent
	<b>(B)</b>	Two circles touching each other	<b>(q)</b>	have a co	ommon n	ormal
	(C)	Two non concentric circles, one strictly inside the other	(r)	do not ha	ave a con	nmon normal
	<b>(D)</b>	Two concentric circles of different radii	(\$)	do not ha	ave a rad	ical axis.



# MATHS FOR JEE MAIN & ADVANCED

4.	Colum	1 – I	Colum	n – 11		
	<b>(A)</b>	Number of common tangents of the circles	<b>(p)</b>	1		
		$x^2 + y^2 - 2x = 0$ and $x^2 + y^2 + 6x - 6y + 2 = 0$ is				
	(B) Number of indirect common tangents of the circles (			2		
		$x^{2} + y^{2} - 4x - 10y + 4 = 0$ & $x^{2} + y^{2} - 6x - 12y - 55 = 0$ is				
	(C)	Number of common tangents of the circles $x^2 + y^2 - 2x - 4y = 0$	(r)	3		
		& $x^2 + y^2 - 8y - 4 = 0$ is				
	<b>(D</b> )	Number of direct common tangents of the circles	<b>(s)</b>	0		
		$x^{2} + y^{2} + 2x - 8y + 13 = 0$ & $x^{2} + y^{2} - 6x - 2y + 6 = 0$ is				
5.	Column-I			ı-II		
	<b>(A)</b>	If the straight line $y = kx \forall K \in I$ touches or passes outside	<b>(p)</b>	1		
		the circle $x^2 + y^2 - 20y + 90 = 0$ then $ k $ can have the value				
	<b>(B)</b>	Two circles $x^2 + y^2 + px + py - 7 = 0$	<b>(q)</b>	2		
		and $x^2 + y^2 - 10x + 2py + 1 = 0$ intersect each other orthogonally				
		then the value of p is				
	(C)	If the equation $x^2 + y^2 + 2\lambda x + 4 = 0$ and $x^2 + y^2 - 4\lambda y + 8 = 0$	<b>(r)</b>	3		
		represent real circles then the value of $\lambda$ can be				
	<b>(D)</b>	Each side of a square is of length 4. The centre of the square is $(3, 7)$ .	<b>(s)</b>	5		
		One diagonal of the square is parallel to $y = x$ . The possible abscissas				
		of the vertices of the square can be				

Part # II

#### [Comprehension Type Questions]

#### **Comprehension # 1**

Let  $A \equiv (-3, 0)$  and  $B \equiv (3, 0)$  be two fixed points and P moves on a plane such that PA = nPB (n > 0). On the basis of above information, answer the following questions :

1.	If $n \neq 1$ , then locus of a point P is -		
	(A) a straight line (B) a circle	(C) a parabola	<b>(D)</b> an ellipse
2.	If $n = 1$ , then the locus of a point P is -		
	(A) a straight line (B) a circle	(C) a parabola	<b>(D)</b> a hyperbola
3.	If $0 < n < 1$ , then -		
	(A) A lies inside the circle and B lies outside the circle		
	(B) A lies outside the circle and B lies inside the circle		
	(C) both A and B lies on the circle	<b>(D)</b> both A and B lies	inside the circle
4.	If $n > 1$ , then -		
	(A) A lies inside the circle and B lies outside the circle	(B) A lies outside the circl	le and B lies inside the circle
	(C) both A and B lies on the circle	<b>(D)</b> both A and B lies ins	side the circle
5.	If locus of P is a circle, then the circle -		
	(A) passes through A and B	(B) never passes throu	igh A and B
	(C) passes through A but does not pass through B	(D) passes through B bu	at does not pass through A



Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141

### **Comprehension # 2**

	Two circles are $S_1 \equiv (x + x)$	$(+3)^2 + y^2 = 9$ $S_2 \equiv (x + 3)^2 + y^2 = 9$	$(-5)^2 + y^2 = 16$ with centres	$C_1 \& C_2$
1.	A direct common tangen of area of $\triangle PQC_1 \& \triangle PRC_1$	t is drawn from a point P w $C_2^2$ .	which touches $S_1 \& S_2$ at Q a	& R, respectively. Find the ratio
	<b>(A)</b> 3 : 4	<b>(B)</b> 9 : 16	<b>(C)</b> 16 : 9	<b>(D)</b> 4 : 3
2.	From point 'A' on $S_2$ whice	ch is nearest to C <sub>1</sub> , a variabl	le chord is drawn to S <sub>1</sub> . The	locus of mid point of the chord.
	(A) circle		<b>(B)</b> Diameter of $s_1$	
	(C) Arc of a circle		<b>(D)</b> chord of $s_1$ but not di	iameter
3.	Locus of 7 cuts the circle	e S <sub>1</sub> at B & C, then line seg	ment BC subtends an angl	e on the major arc of circle S <sub>1</sub> is
	(A) $\cos^{-1}\frac{3}{4}$		<b>(B)</b> $\frac{\pi}{2} - \tan^{-1}\frac{4}{3}$	
	(C) $\frac{\pi}{2} - \frac{1}{2} \tan^{-1} \frac{3}{4}$		<b>(D)</b> $\frac{\pi}{2} \cot^{-1} \frac{4}{3}$	
		Comprehe	nsion # 3	
	Consider a line pair $ax^2 + forming a triangle ABC with the above the triangle according to th$	$3xy - 2y^2 - 5x + 5y + c = 0$ returns the x-axis.	epresenting perpendicular lin	es intersecting each other at C and
1.	If $x_1$ and $x_2$ are intercepts of is equal to	on the x-axis and y <sub>1</sub> and y <sub>2</sub> a	re the intercepts on the y-ax	is then the sum $(x_1 + x_2 + y_1 + y_2)$
	<b>(A)</b> 6	<b>(B)</b> 5	<b>(C)</b> 4	<b>(D)</b> 3
2.	Distance between the orth	ocentre and circumcentre o	f the triangle ABC is	
	<b>(A)</b> 2	<b>(B)</b> 3	<b>(C)</b> 7/4	<b>(D)</b> 9/4
3.	If the circle $x^2 + y^2 - 4y +$	k = 0 is orthogonal with the	circumcircle of the triangle	ABC then 'k' equals
	<b>(A)</b> 1/2	<b>(B)</b> 1	(C) 2	<b>(D)</b> 3/2
		Comprehensi	on # 4	
1.	Let $S_1, S_2, S_3$ be the circles Point from which length of	$x^2 + y^2 + 3x + 2y + 1 = 0, x^2$ of tangents to these three cir	$+y^2 - x + 6y + 5 = 0$ and $x^2 +$ scles is same is	$-y^2 + 5x - 8y + 15 = 0$ , then
	(A) (1.0)	<b>(B)</b> (3, 2)	<b>(C)</b> (10, 5)	(D)(-2,1)
2.	Equation of circle $S_4$ which	h cut orthogonally to all giv	ven circle is	
	(A) $x^2 + y^2 - 6x + 4y - 14 =$	= 0	<b>(B)</b> $x^2 + y_2 + 6x + 4y - 14 =$	0
	(C) $x^2 + y^2 - 6x - 4y + 14 =$	= 0	<b>(D)</b> $x^2 + y^2 - 6x - 4y - 14 =$	0
3.	Radical centre of circles S	$_{1}, S_{2}, \& S_{4}$ is		
	$(\mathbf{A})\left(-\frac{3}{5},-\frac{8}{5}\right)$	<b>(B)</b> (3,2)	<b>(C)</b> (1,0)	<b>(D)</b> $\left(-\frac{4}{5}, -\frac{3}{2}\right)$



#### **Comprehension # 5**

P is a variable point of the line L = 0. Tangents are drawn to the circle  $x^2 + y^2 = 4$  from P to touch it at Q and R. The parallelogram PQSR is completed.

On the basis of above information, answer the following questions :

1. If  $L \equiv 2x + y - 6 = 0$ , then the locus of circumcetre of  $\triangle PQR$  is -

(A) 2x - y = 4 (B) 2x + y = 3 (C) x - 2y = 4 (D) x + 2y = 3

2. If  $P \equiv (6, 8)$ , then the area of  $\triangle QRS$  is -

(A) 
$$\frac{(6)^{3/2}}{25}$$
 sq. units (B)  $\frac{(24)^{3/2}}{25}$  sq. units (C)  $\frac{48\sqrt{6}}{25}$  sq. units (D)  $\frac{192\sqrt{6}}{25}$  sq. units

3. If  $P \equiv (3, 4)$ , then coordinate of S is -

(A) 
$$\left(-\frac{46}{25},-\frac{63}{25}\right)$$
 (B)  $\left(-\frac{51}{25},-\frac{68}{25}\right)$  (C)  $\left(-\frac{46}{25},-\frac{68}{25}\right)$  (D)  $\left(-\frac{68}{25},-\frac{51}{25}\right)$ 

#### **Comprehension # 6**

Let C be a circle of radius *r* with centre at O. Let P be a point outside C and D be a point on C. A line through P intersects C at Q and R, S is the midpoint of QR.

1. For different choices of line through P, the curve on which S lies, is

(A) a straight line
(B) an arc of circle with P as centre
(C) an arc of circle with PS as diameter
(D) an arc of circle with OP as diameter

2. Let P is situated at a distance 'd' from centre O, then which of the following does not equal the product (PQ) (PR)?

(A)  $d^2 - r^2$ (B) PT<sup>2</sup>, where T is a point on C and PT is tangent to C (C) (PS)<sup>2</sup>-(QS)(RS) (D) (PS)<sup>2</sup>

3. Let XYZ be an equilateral triangle inscribed in C. If  $\alpha$ ,  $\beta$ ,  $\gamma$  denote the distances of D from vertices X, Y, Z respectively, the value of product  $(\beta + \gamma - \alpha) (\gamma + \alpha - \beta) (\alpha + \beta - \gamma)$ , is

(A) 0 (B) 
$$\frac{\alpha\beta\gamma}{8}$$
 (C)  $\frac{\alpha^3 + \beta^3 + \gamma^3 - 3\alpha\beta\gamma}{6}$  (D) None of these



	Comprehension # 7						
	Let A, B, C be three sets o	f real numbers (x, y) define	d as				
	$A:\{(x,y):y\ge 1\}$	B: {(x, y): $x^2 + y^2 - 4x - 2$	$y-4=0$ } C: {(x,	y): $x + y = \sqrt{2}$ }			
1.	Number of elements in the	$A \cap B \cap C$ is					
	<b>(A)</b> 0	<b>(B)</b> 1	<b>(C)</b> 2	(D) infinite			
2.	$(x+1)^2 + (y-1)^2 + (x-5)^2$	$(y-1)^2$ has the value eq	jual to				
	<b>(A)</b> 16	<b>(B)</b> 25	<b>(C)</b> 36	<b>(D)</b> 49			
3.	If the locus of the point of intersection of the pair of perpendicular tangents to the circle B is the curve S then the area enclosed between B and S is						
	(Α) 6π	<b>(B)</b> 8π	<b>(C)</b> 9π	<b>(D)</b> 18π			
	Comprehension # 8						
	Consider a circle $x^2 + y^2 = 4$ and a point P(4, 2). $\theta$ denotes the angle enclosed by the tangents from P on the circle and A, B are the points of contact of the tangents from P on the circle.						
1.	The value of $\theta$ lies in the interval						
	<b>(A)</b> (0, 15°)	<b>(B)</b> (15°, 30°)	(C) 30°, 45°)	<b>(D)</b> (45°, 60°)			
2.	The intercept made by a t	angent on the x-axis is					
	<b>(A)</b> 9/4	<b>(B)</b> 10/4	<b>(C)</b> 11/4	<b>(D)</b> 12/4			

3. Locus of the middle points of the portion of the tangent to the circle terminated by the coordinate axes is

**(B)**  $x^{-2} + y^{-2} = 2^{-2}$ 

(A)  $x^{-2} + y^{-2} = 1^{-2}$ 

(C)  $x^{-2} + y^{-2} = 3^{-2}$  (D)  $x^{-2} - y^{-2} = 4^{-2}$ 





- 12. Show that the locus of the point the tangents from which to the circle  $x^2 + y^2 a^2 = 0$  include a constant angle  $\alpha$  is  $(x^2 + y^2 2a^2)^2 \tan^2 \alpha = 4a^2 (x^2 + y^2 a^2).$
- 13. The curves whose equations are

 $S = ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ 

$$S' = a'x^2 + 2h'xy + b'y^2 + 2g'x + 2f'y + c' = 0$$

intersect in four concyclic points then, what is the relation between a,b,h,a'b'h'



- 14. A circle touches the line y = x at a point P such that  $OP = 4\sqrt{2}$ , where O is the origin. The circle contains the point (-10,2) in its interior and the length of its chord on the line x + y = 0 is  $6\sqrt{2}$ . Determine the equation of the circle.
- 15. If the line  $x \sin \alpha y + a \sec \alpha = 0$  touches the circle with radius 'a' and centre at the origin then find the most general values of ' $\alpha$ ' and sum of the values of ' $\alpha$ ' lying in [0, 100 $\pi$ ].
- 16. P is a variable point on the circle with centre at C. CA & CB are perpendiculars from C on x-axis & y-axis respectively. Show that the locus of the centroid of the triangle PAB is a circle with centre at the centroid of the triangle CAB & radius equal to one third of the radius of the given circle.
- 17. Show that the equation  $x^2 + y^2 2x 2\lambda y 8 = 0$  represents, for different values of  $\lambda$ , a system of circles passing through two fixed points A, B on the x axis, and find the equation of that circle of the system the tangents to which at A & B meet on the line x + 2y + 5 = 0.
- 18. A circle touches the line y = x at a point P such that  $OP = 4\sqrt{2}$  where O is the origin. The circle contains the point (-10, 2) in its interior and the length of its chord on the line x + y = 0 is  $6\sqrt{2}$ . Find the equation of the circle.
- 19. Find the intervals of values of 'a' for which the line y + x = 0 bisects two chords drawn from a point  $\left(\frac{1+\sqrt{2}a}{2}, \frac{1-\sqrt{2}a}{2}\right)$  to the circle  $2x^2 + 2y^2 \left(1 + \sqrt{2}a\right)x \left(1 \sqrt{2}a\right)y = 0$ .
- 20. Show that the locus of the centres of a circle which cuts two given circles orthogonally is a straight line & hence deduce the locus of the centre of the circles which cut the circles  $x^2 + y^2 + 4x 6y + 9 = 0$  &  $x^2 + y^2 5x + 4y + 2 = 0$  orthogonally.



# MATHS FOR JEE MAIN & ADVANCED

E	xercise # 5	Part # I  Pre	vious Year Questions]	AIEEE/JEE-MAIN]
1.	The square of the lengt	h of tangent from (3, –4) on	the circle $x^2 + y^2 - 4x - 6y + $	-3 = 0 [AIEEE-2002
	<b>(A)</b> 20	<b>(B)</b> 30	<b>(C)</b> 40	<b>(D)</b> 50
2.	Radical axis of the circ	$\log x^2 + y^2 + 6x - 2y - 9 = 0 \text{ ar}$	$dx^2 + y^2 - 2x + 9y - 11 = 0$	s- [AIEEE-2002
	(A) $8x - 11y + 2 = 0$	<b>(B)</b> $8x + 11y + 2 = 0$	(C) $8x + 11y - 2 = 0$	<b>(D)</b> $8x - 11y - 2 = 0$
3.	If the two circles $(x - 1)$	$^{2} + (y-3)^{2} = r^{2}$ and $x^{2} + y^{2} - y^{2}$	8x + 2y + 8 = 0 intersect in	two distinct points, then- [AIEEE-2003
	(A) $r > 2$	<b>(B)</b> $2 < r < 8$	( <b>C</b> ) r < 2	<b>(D)</b> $r = 2$
4.	The lines $2x - 3y = 5$ as circle is-	nd $3x - 4y = 7$ are diameters	of a circle having area as 15	4 sq. units. Then the equation of th [AIEEE-2003
	(A) $x^2 + y^2 - 2x + 2y = 6$	2	<b>(B)</b> $x^2 + y^2 + 2x - 2y = 62$	2
	(C) $x^2 + y^2 + 2x - 2y = 4$	7	<b>(D)</b> $x^2 + y^2 - 2x + 2y = 4^2$	7
5.	If a circle passes throug	gh the point (a, b) and cuts th	e circle $x^2 + y^2 = 4$ orthogon	ally, then the locus of its centre is- [AIEEE-2004
	(A) $2ax + 2by + (a^2 + b^2)$	(+4) = 0	<b>(B)</b> $2ax + 2by - (a^2 + b^2)$	(+4) = 0
	(C) $2ax - 2by + (a^2 + b^2)$	+4) = 0	<b>(D)</b> $2ax - 2by - (a^2 + b^2)$	(+4) = 0
6.	A variable circle passes through A is-	through the fixed point A(p,	q) and touches x-axis. The l	ocus of the other end of the diamete [AIEEE-2004
	(A) $(x-p)^2 = 4qy$	<b>(B)</b> $(x-q)^2 = 4py$	(C) $(y-p)^2 = 4qx$	<b>(D)</b> $(y-q)^2 = 4px$
7.	If the lines $2x + 3y + 1 =$ the circle is-	= 0  and  3x - y - 4 = 0  lie alon	g diameters of a circle of circ	cumference $10\pi$ , then the equation (AIEEE-2004)
	(A) $x^2 + y^2 - 2x + 2y - 2$	3=0	<b>(B)</b> $x^2 + y^2 - 2x - 2y - 23$	3 = 0
	(C) $x^2 + y^2 + 2x + 2y - 2$	3=0	(D) $x^2 + y^2 + 2x - 2y - 2$	3=0
8.	The intercept on the lin	the $y = x$ by the circle $x^2 + y^2$	-2x = 0 is AB. Equation of	the circle on AB as a diameter is- [AIEEE-2004
	(A) $x^2 + y^2 - x - y = 0$	<b>(B)</b> $x^2 + y^2 - x + y = 0$	(C) $x^2 + y^2 + x + y = 0$	<b>(D)</b> $x^2 + y^2 + x - y = 0$
9.	If the circles $x^2 + y^2 + 2a^2$ 5x + by - a = 0 passes	$ax + cy + a = 0$ and $x^2 + y^2 - 3$ through P and Q for-	ax + dy - 1 = 0 intersect in two	o distinct point P and Q then the lir [AIEEE-2003]
	(A) exactly one value of	fa	(B) no value of a	
	(C) infinitely many val	ues of a	(D) exactly two values of	ofa
10.	A circle touches the x-a circle is-	ixis and also touches the circ	le with centre at $(0, 3)$ and ra	dius 2. The locus of the centre of th [AIEEE-200:

(A) an ellipse	(B) a circle	(C) a hyperbola	(D) a parabola



Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141

[AIEEE-2005]

	(A) $x^2 + y^2 - 3ax - 4by + (a^2 + b^2 - p^2) = 0$ (C) $x^2 + y^2 - 2ax - 3by + (a^2 - b^2 - p^2) = 0$	(B) $2ax + 2by - (a^2 - b^2 + p^2) = 0$ (D) $2ax + 2by - (a^2 + b^2 + p^2) = 0$			
12.	If the pair of lines $ax^2 + 2(a + b)xy + by^2 = 0$ lie along d that the area of one of the sectors is thrice the area of	diameters of a circle and divide the circle into four sectors of another sector then-	s such -2005]		
	(A) $3a^2 - 10ab + 3b^2 = 0$ (C) $3a^2 + 10ab + 3b^2 = 0$	(B) $3a^2 - 2ab + 3b^2 = 0$ (D) $3a^2 + 2ab + 3b^2 = 0$			
13.	If the lines $3x - 4y - 7 = 0$ and $2x - 3y - 5 = 0$ are two the circle is-	diameters of a circle of area $49\pi$ square units, the equat [AIEEE	ion of - <b>2006]</b>		
	(A) $x^2 + y^2 + 2x - 2y - 62 = 0$ (C) $x^2 + y^2 - 2x + 2y - 47 = 0$	(B) $x^2 + y^2 - 2x + 2y - 62 = 0$ (D) $x^2 + y^2 + 2x - 2y - 47 = 0$			
14.	Let C be the circle with centre $(0, 0)$ and radius 3 unit	ts. The equation of the locus of the mid points of the cho	rds of		
	the circle C that subtend an angle of $\frac{2\pi}{3}$ at its centre	e is - [AIEEE-2006, IIT-	1996]		
	(A) $x^2 + y^2 = 1$ (B) $x^2 + y^2 = \frac{27}{4}$	(C) $x^2 + y^2 = \frac{9}{4}$ (D) $x^2 + y^2 = \frac{3}{2}$			
15.	Consider a family of circles which are passing through ordinates of the centre of the circles, then the set of v (A) $0 < k < 1/2$ (B) $k \ge 1/2$	h the point (-1, 1) and are tangent to x-axis. If (h, k) are the values of k is given by the interval- (C) $-1/2 \le k \le 1/2$ (D) $k \le 1/2$	he co- -2007]		
16.	The point diametrically opposite to the point $(1, 0)$ on <b>(A)</b> $(3,-4)$ <b>(B)</b> $(-3,4)$	the circle $x^2 + y^2 + 2x + 4y - 3 = 0$ is- (C) (-3, -4) (D) (3, 4)	-2008]		
17.	Three distinct points A, B and C are given in the 2-di	imensional coordinate plane such that the ratio of the dis	stance		
	of any one of them from the point (1, 0) to the distance from the point (-1, 0) is equal to $\frac{1}{3}$ . Then the circumcentre				
	of the triangle ABC is at the point :-	[AIEEE	-2009]		
	$(\mathbf{A})\left(\frac{5}{2},0\right) \qquad \qquad (\mathbf{B})\left(\frac{5}{3},0\right)$	(C) (0,0) (D) $\left(\frac{5}{4}, 0\right)$			
18.	If P and Q are the points of intersection of $x^2 + y^2 + 2x + 2y - p^2 = 0$ , then there is a circle pass	f the circles $x^2 + y^2 + 3x + 7y + 2p - 5 = 0$ sing through P, Q and (1, 1) for :-	) and - <b>2009]</b>		
	(A) All except two values of p	(B) Exactly one value of p			
	(C) All values of p	(D) All except one value of p			
19.	For a regular polygon, let r and R be the radii of the inst the following is :-	scribed and the circumscribed circles. A false statement a [AIEEE	mong - <b>2010]</b>		
	(A) There is a regular polygon with $\frac{r}{R} = \frac{1}{2}$	(B) There is a regular polygon with $\frac{r}{R} = \frac{1}{\sqrt{2}}$			
	(C) There is a regular polygon with $\frac{r}{R} = \frac{2}{3}$	(D) There is a regular polygon with $\frac{r}{R} = \frac{\sqrt{3}}{2}$			
20.	The circle $x^2 + y^2 = 4x + 8y + 5$ intersects the line 3:	x - 4y = m at two distinct points if :- [AIEEE	-2010]		
	(A) $-85 < m < -35$ (B) $-35 < m < 15$	(C) $15 < m < 65$ (D) $35 < m < 85$			

If a circle passes through the point (a, b) and cuts the circle  $x^2 + y^2 = p^2$  orthogonally, then the equation of the locus



11.

of its centre is-

# Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141

# MATHS FOR JEE MAIN & ADVANCED

21.	The two circles $x^2$ +	[AIEEE-2011]									
	$(\mathbf{A}) \mathbf{a} = 2\mathbf{c}$	<b>(B)</b> $ a  = 2c$	(C) $2 a  = c$	$(\mathbf{D})  \mathbf{a}  = \mathbf{c}$							
22.	The equation of the	circle passing through the po	points $(1, 0)$ and $(0, 1)$ a	est radius is:							
	(A) $x^2 + y^2 + x + y =$ (C) $x^2 + y^2 - x - y =$	-2 = 0 = 0	(B) $x^2 + y^2 - 2x - 2x^2$ (C) $x^2 + y^2 + 2x + 2x^2$	(B) $x^2 + y^2 - 2x - 2y + 1 = 0$ (C) $x^2 + y^2 + 2x + 2y - 7 = 0$							
23.	The length of the di $(2, 3)$ is :	ameter of the circle which tou	uches the x-axis at the p	hes the x-axis at the point (1, 0) and passes through the p [AIEEE-2							
	<b>(A)</b> 5/3	<b>(B)</b> 10/3	<b>(C)</b> 3/5	<b>(D)</b> 6/5							
24.	The circle passing t $(A)$ (-5, 2)	hrough $(1, -2)$ and touching (B) $(2, -5)$	the axis of x at (3, 0) a (C) (5, -2)	also passes through (D) (-2, 5)	the point : [JEE(Main)-2013]						
25.	Let C be the circle v and touching the ci	entered at (0, y), pas	ussing through origin [JEE Main 2014]								
	(A) $\frac{\sqrt{3}}{\sqrt{2}}$	<b>(B)</b> $\frac{\sqrt{3}}{2}$	(C) $\frac{1}{2}$	<b>(D)</b> $\frac{1}{4}$							
26.	Locus of the image o	f the point $(2, 3)$ in the line $(2x)$	-3y+4)+k(x-2y+3)	$=0, k \in \mathbb{R}$ , is a :	[JEE Main 2015]						
	<ul><li>(A) circle of radius</li><li>(C) straight line para</li></ul>	$\sqrt{2}$ llel to x-axis	(B) circle of radius (D) straight line par	(B) circle of radius $\sqrt{3}$ (D) straight line parallel to y-axis							
27.	The number of com	non tangents to the circles $x^2$ +	$-y^2 - 4x - 6y - 12 = 0$ ar	$x^2 + y^2 + 6x + 18y$	+26 = 0, is :						
	<b>(A)</b> 3	<b>(B)</b> 4	(C) 1	<b>(D)</b> 2	[ <b>JEE Main 2015</b> ]						
28.	The centres of those	circle which touch the circle, $x^2$	$y^2 + y^2 - 8x - 8y - 4 = 0$ , ex	xternally and also tou	ch the x-axis, lie on : [JEE Main 2016]						
	(A) an ellipse which (C) a parabola	is not a circle	(B) a hyperbola (D) a circle								
29.	If one of the diameter centre is at $(-3, 2)$ , t	ers of the circle, given by the e hen the radius of S is :	quation, $x^2 + y^2 - 4x + 6$	y-12=0, is a chord	of a circle S, whose [JEE Main 2016]						
	(A) 5√3	<b>(B)</b> 5	<b>(C)</b> 10	<b>(D)</b> $5\sqrt{2}$							
	Part # II	Previous Year Que	estions][IIT-JEE Al	DVANCED]							
1.	Let PQ and RS be tar X on the circumfere	ngents at the extremities of the on the one of the circle then 2r equals	liameter PR of a circle of	fradius r. If PS and R	Q intersect at a point [JEE 2001]						
	(A) $\sqrt{PQ \cdot RS}$	<b>(B)</b> $\frac{PQ + RS}{2}$	(C) $\frac{2PQ \cdot RS}{PQ + RS}$	<b>(D)</b> $\sqrt{\frac{(PQ)^2}{2}}$	$\frac{1}{2} + (RS)^2$						
2	Let $2x^2 + y^2 - 3xy =$	0 be the equation of a pair of ta	ngents drawn from the o	rigin 'O' to a circle of	fradius 3 with centre						

Let  $2x^2 + y^2 - 3xy = 0$  be the equation of a pair of tangents drawn from the origin 'O' to a circle of radius 3 with centre in the first quadrant. If A is one of the points of contact, find the length of OA.

[**JEE 2001**]

3. Find the equation of the circle which passes through the points of intersection of circles  $x^2 + y^2 - 2x - 6y + 6 = 0$  and  $x^2 + y^2 + 2x - 6y + 6 = 0$  and intersects the circle  $x^2 + y^2 + 4x + 6y + 4 = 0$  orthogonally. [JEE 2001]

- 4. Tangents TP and TQ are drawn from a point T to the circle  $x^2 + y^2 = a^2$ . If the point T lies on the line px + qy = r, find the locus of centre of the circumcircle of triangle TPQ.
- 5. If the tangent at the point P on the circle  $x^2 + y^2 + 6x + 6y = 2$  meets the straight line 5x 2y + 6 = 0 at a point Q on the y-axis, then the length of PQ is [JEE 2002]
  - (A) 4 (B)  $2\sqrt{5}$  (C) 5 (D)  $3\sqrt{5}$
- 6. If a > 2b > 0 then the positive value of m for which  $y = mx b\sqrt{1 + m^2}$  is a common tangent to  $x^2 + y^2 = b^2$  and  $(x a)^2 + y^2 = b^2$  is [JEE 2002]

(A) 
$$\frac{2b}{\sqrt{a^2 - 4b^2}}$$
 (B)  $\frac{\sqrt{a^2 - 4b^2}}{2b}$  (C)  $\frac{2b}{a - 2b}$  (D)  $\frac{b}{a - 2b}$ 

- 7. The radius of the circle, having centre at (2, 1), whose one of the chord is a diameter of the circle  $x^2+y^2-2x-6y+6=0$  [JEE 2004]
  - (A) 1 (B) 2 (C) 3 (D)  $\sqrt{3}$
- 8. Line 2x + 3y + 1 = 0 is a tangent to a circle at (1, -1). This circle is orthogonal to a circle which is drawn having diameter as a line segment with end points (0, -1) and (-2, 3). Find equation of circle. [JEE 2004]
- 9. A circle is given by  $x^2 + (y-1)^2 = 1$ , another circle C touches it externally and also the x-axis, then the locus of its centre is [JEE 2005]

(A) 
$$\{(x, y) : x^2 = 4y\} \cup \{(x, y) : y \le 0\}$$
(B)  $\{(x, y) : x^2 + (y - 1)^2 = 4\} \cup \{x, y) : y \le 0\}$ (C)  $\{(x, y) : x^2 = y\} \cup \{(0, y) : y \le 0\}$ (D)  $\{(x, y) : x^2 = 4y\} \cup \{(0, y) : y \le 0\}$ 

- Let ABCD be a quadrilateral with area 18, with side AB parallel to the side CD and AB = 2CD. Let AD be perpendicular to AB and CD. If a circle is drawn inside the quadrilateral ABCD touching all the sides, then its radius is
  - (A) 3 (B) 2 (C) 3/2 (D) 1
- 11. Tangents are drawn from the point (17, 7) to the circle  $x^2 + y^2 = 169$ .

Statement-1 : The tangents are mutually perpendicular.

Statement-2: The locus of the points from which mutually perpendicular tangents can be drawn to the given circle is  $x^2 + y^2 = 338$ .

- (A) Statement-1 is true, statement-2 is true; statement-2 is correct explanation for statement-1.
- (B) Statement-1 is true, statement-2 is true; statement-2 is NOT a correct explanation for statement-1.
- (C) Statement-1 is true, statement-2 is false.
- (D) Statement-1 is false, statement-2 is true.
- 12. Consider the two curves  $C_1 : y^2 = 4x$ ;  $C_2 : x^2 + y^2 6x + 1 = 0$ . Then,
  - (A)  $C_1$  and  $C_2$  touch each other only at one point
  - **(B)**  $C_1$  and  $C_2$  touch each other exactly at two points
  - (C)  $C_1$  and  $C_2$  intersect (but do not touch) at exactly two points
  - **(D)**  $C_1$  and  $C_2$  neither intersect nor touch each other

[**JEE 2007**]



### MATHS FOR JEE MAIN & ADVANCED

**13.** Consider,  $L_1: 2x + 3y + p - 3 = 0; L_2: 2x + 3y + p + 3 = 0,$ 

where p is a real number, and  $C: x^2 + y^2 + 6x - 10y + 30 = 0$ .

**Statement-1**: If line  $L_1$  is a chord of circle C, then line  $L_2$  is not always a diameter of circle C.

**Statement-2**: If line  $L_1$  is a diameter of circle C, then line  $L_2$  is not a chord of circle C.

- (A) Statement-1 is True, Statement-2 is True; statement-2 is a correct explanation for statement-1
- (B) Statement-1 is True, Statement-2 is True; statement-2 is NOT a correct explanation for statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

#### 14. Comprehension

A circle C of radius 1 is inscribed in an equilateral triangle PQR. The points of contact of C with the sides PQ, QR, RP

are D, E, F respectively. The line PQ is given by the equation  $\sqrt{3} x + y - 6 = 0$  and the point D is  $\left(\frac{3\sqrt{3}}{2}, \frac{3}{2}\right)$ .

Further, it is given that the origin and the centre of C are on the same side of the line PQ.

(i) The equation of circle C is

(A) 
$$(x-2\sqrt{3})^2 + (y-1)^2 = 1$$

(C) 
$$(x - \sqrt{3})^2 + (y + 1)^2 = 1$$

(ii) Points E and F are given by

(A) 
$$\left(\frac{\sqrt{3}}{2}, \frac{3}{2}\right), \left(\sqrt{3}, 0\right)$$
  
(C)  $\left(\frac{\sqrt{3}}{2}, \frac{3}{2}\right), \left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ 

(iii) Equations of the sides RP, RQ are

(A) 
$$y = \frac{2}{\sqrt{3}}x + 1$$
,  $y = -\frac{2}{\sqrt{3}}x - 1$   
(C)  $y = \frac{\sqrt{3}}{2}x + 1$ ,  $y = -\frac{\sqrt{3}}{2}x - 1$ 

(B)  $(x-2\sqrt{3})^2 + (y+\frac{1}{2})^2 = 1$ (D)  $(x-\sqrt{3})^2 + (y-1)^2 = 1$ (B)  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right), (\sqrt{3}, 0)$  [**JEE 2008**]

**(D)**
$$\left(\frac{3}{2}, \frac{\sqrt{3}}{2}\right), \left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$$

(B) 
$$y = \frac{1}{\sqrt{3}} x, y = 0$$
  
(D)  $y = \sqrt{3} x, y = 0$  [JEE 2008]

15. Tangents drawn from the point P(1, 8) to the circle  $x^2 + y^2 - 6x - 4y - 11 = 0$  touch the circle at the points A and B. The equation of the circumcircle of the triangle PAB is

(A) 
$$x^2 + y^2 + 4x - 6y + 19 = 0$$
  
(B)  $x^2 + y^2 - 4x - 10y + 19 = 0$   
(C)  $x^2 + y^2 - 2x + 6y - 29 = 0$   
(D)  $x^2 + y^2 - 6x - 4y + 19 = 0$   
[JEE 2009]

16. The centres of two circles  $C_1$  and  $C_2$  each of unit radius are at a distance of 6 units from each other. Let P be the mid point of the line segment joining the centres of  $C_1$  and  $C_2$  and C be a circle touching circles  $C_1$  and  $C_2$  externally. If a common tangent to  $C_1$  and C passing through P is also a common tangent to  $C_2$  and C, then the radius of the circle C is [JEE 2009]

17. Two parallel chords of a circle of radius 2 are at a distance  $\sqrt{3} + 1$  apart. If the chords subtend at the center,

angles of 
$$\frac{\pi}{k}$$
 and  $\frac{2\pi}{k}$ , where  $k > 0$ , then the value of [k] is [JEE 2010]

[Note : [k] denotes the largest integer less than or equal to k]



[**JEE 2011**]

[**JEE 2012**]

= 5

18. The circle passing through the point (-1,0) and touching the y-axis at (0, 2) also passes through the point -

(A) 
$$\left(-\frac{3}{2}, 0\right)$$
 (B)  $\left(-\frac{5}{2}, 2\right)$  (C)  $\left(-\frac{3}{2}, \frac{5}{2}\right)$  (D) (-4,0) [JEE 2011]

The straight line 2x - 3y = 1 divides the circular region  $x^2 + y^2 \le 6$  into two parts. If 19.

$$S = \left\{ \left(2, \frac{3}{4}\right), \left(\frac{5}{2}, \frac{3}{4}\right), \left(\frac{1}{4}, -\frac{1}{4}\right), \left(\frac{1}{8}, \frac{1}{4}\right) \right\},\$$

then the number of point(s) in S lying inside the smaller part is

20. The locus of the mid-point of the chord of contact of tangents drawn from points lying on the straight line 4x - 5y = 20 to the circle  $x^2 + y^2 = 9$  is-[**JEE 2012**] **(B)**  $20(x^2 + y^2) + 36x - 45y = 0$ **(D)**  $36(x^2 + y^2) + 20x - 45y = 0$ (A)  $20(x^2 + y^2) - 36x + 45y = 0$ (C)  $36(x^2 + y^2) - 20x + 45y = 0$ 

#### Paragraph for Question 21 and 22

A tangent PT is drawn to the circle  $x^2 + y^2 = 4$  at the point  $P(\sqrt{3}, 1)$ . A straight line L, perpendicular to PT is a tangent to the circle  $(x-3)^2 + y^2 = 1$ .

21. A common tangent of the two circles is

(A) $x = 4$	<b>(B)</b> $y = 2$	(C) $x + \sqrt{3}y = 4$	<b>(D)</b> $x + 2\sqrt{2}y = 6$	
A possible equation	n of L is		[ <b>JEE 201</b> ]	21

22.

(A) 
$$x - \sqrt{3}y = 1$$
 (B)  $x + \sqrt{3}y = 1$  (C)  $x - \sqrt{3}y = -1$  (D)  $x + \sqrt{3}y$ 

Circle(s) touching x-axis at a distance 3 from the origin and having an intercept of length  $2\sqrt{7}$  or y-axis 23. [JEE Ad. 2013] is (are)

(A) 
$$x^{2} + y^{2} - 6x + 8y + 9 = 0$$
  
(B)  $x^{2} + y^{2} - 6x + 7y + 9 = 0$   
(D)  $x^{2} + y^{2} - 6x - 7y + 9 = 0$ 

- A circle S passes through the point (0, 1) and is orthogonal to the circles  $(x 1)^2 + y^2 = 16$  and  $x^2 + y^2 = 1$ . Then, 24. (A) radius of S is 8 **(B)** radius of S is 7 [JEE Ad. 2014] (C) centre of S is (-7, 1)(D) centre of S is (-8, 1)
- 25. Let RS be the diameter of the circle  $x^2 + y^2 = 1$ , where S is the point (1, 0). Let P be a variable point (other than R and S) on the circle and tangents to the circle at S and P meets at the point Q. The normal to the circle at P intersects a line drawn through Q parallel to RS at point E. Then the locus of E passes through the point(s) [**JEE Ad. 2016**]

**(A)** 
$$\left(\frac{1}{3}, \frac{1}{\sqrt{3}}\right)$$
 **(B)**  $\left(\frac{1}{4}, \frac{1}{2}\right)$  **(C)**  $\left(\frac{1}{3}, -\frac{1}{\sqrt{3}}\right)$  **(D)**  $\left(\frac{1}{4}, -\frac{1}{2}\right)$ 

26.

The circle  $C_1$ :  $x^2 + y^2 = 3$ , with centre at O, intersects the parabola  $x^2 = 2y$  at the point P in the first quadrant. Let the tangent to the circle C<sub>1</sub> at P touches other two circles C<sub>2</sub> and C<sub>3</sub> at R, and R<sub>3</sub>, respectively. Suppose C<sub>2</sub> and C<sub>3</sub> have equal radii  $2\sqrt{3}$  and centres Q<sub>2</sub> and Q<sub>3</sub>, respectively. If Q<sub>2</sub> and Q<sub>3</sub> lie on the y-axis, then [JEE Ad. 2016] **(B)**  $R_2 R_3 = 4\sqrt{6}$ (A)  $Q_2 Q_3 = 12$ 

(C) area of the triangle  $OR_2R_3$  is  $6\sqrt{2}$ 



# Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91 - 9350679141

<sup>(</sup>**D**) area of the triangle PQ<sub>2</sub>Q<sub>3</sub> is  $4\sqrt{2}$ 



#### **SECTION - I : STRAIGHT OBJECTIVE TYPE**

1. The axes are translated so that the new equation of the circle  $x^2 + y^2 - 5x + 2y - 5 = 0$  has no first degree terms. Then the new equation is :

(A)  $x^2 + y^2 = 9$  (B)  $x^2 + y^2 = \frac{49}{4}$  (C)  $x^2 + y^2 = \frac{81}{16}$  (D) none of these

- 2. S(x, y) = 0 represents a circle. The equation S(x, 2) = 0 gives two identical solutions x = 1 and the equation S(1, y) = 0 gives two distinct solutions y = 0, 2. Find the equation of the circle. (A)  $x^2 + y^2 + 2x - 2y + 1 = 0$ (B)  $x^2 + y^2 - 2x + 2y + 1 = 0$ (C)  $x^2 + y^2 - 2x - 2y - 1 = 0$ (D)  $x^2 + y^2 - 2x - 2y + 1 = 0$
- 3. A line meets the co-ordinate axes in A and B. A circle is circumscribed about the triangle OAB. If  $d_1$  and  $d_2$  are the distances of the tangent to the circle at the origin O from the points A and B respectively, then diameter of the circle is:

(A) 
$$\frac{2d_1 + d_2}{2}$$
 (B)  $\frac{d_1 + 2d_2}{2}$  (C)  $d_1 + d_2$  (D)  $\frac{d_1d_2}{d_1 + d_2}$ 

- 4. Consider a family of circles passing through two fixed points A(3,7) & B(6,5). Find the point of concurrency of the chords in which the circle  $x^2 + y^2 4x 6y 3 = 0$  cuts the members of the family :
  - (A)  $\left(\frac{11}{17}, \frac{3}{7}\right)$  (B)  $\left(2, \frac{23}{3}\right)$  (C) (-4, 3) (D) chords are not concurrent
- 5. A circle is inscribed (i.e. touches all four sides ) into a rhombous ABCD with one angle 60°. The distance from the centre of the circle to the nearest vertex is equal to 1. If P is any point of the circle, then

$$|PA|^{2} + |PB|^{2} + |PC|^{2} + |PD|^{2}$$
 is equal to:  
(A) 12 (B) 11 (C) 10 (D) 13

6. If the radius of the circumcircle of the triangle TPQ, where PQ is chord of contact corresponding to point T with respect to circle  $x^2 + y^2 - 2x + 4y - 11 = 0$ , is 6 units, then minimum distance of T from the director circle of the given circle is:

(A) 6 (B) 12 (C) 
$$6\sqrt{2}$$
 (D)  $12-4\sqrt{2}$ 

7. Consider points A ( $\sqrt{13}$ , 0) and B ( $2\sqrt{13}$ , 0) lying on x-axis. These points are rotated in an-anticlockwise direction about the origin through an angle of  $\tan^{-1}\left(\frac{2}{3}\right)$ . Let the new position of A and B be A' and B' respectively. With A' as centre and radius  $\frac{2\sqrt{13}}{3}$  a circle C<sub>1</sub> is drawn and with B' as a centre and radius  $\frac{\sqrt{13}}{3}$  circle C<sub>2</sub> is drawn. The radical axis of C<sub>1</sub> and C<sub>2</sub> is : (A) 9x + 6y = 65 (B) 3x + 3y = 10 (C) 3x + 2y = 20 (D) none of these



8. A circle touches the lines  $y = \frac{x}{\sqrt{3}}$ ,  $y = x\sqrt{3}$  and has unit radius. If the centre of this circle lies in the first quadrant,

then one possible equation of this circle is -

(A) 
$$x^2 + y^2 - 2x(\sqrt{3} + 1) - 2y(\sqrt{3} + 1) + 8 + 4\sqrt{3} = 0$$
  
(B)  $x^2 + y^2 - 2x(\sqrt{3} + 1) - 2y(\sqrt{3} + 1) + 5 + 4\sqrt{3} = 0$   
(C)  $x^2 + y^2 - 2x(\sqrt{3} + 1) - 2y(\sqrt{3} + 1) + 7 + 4\sqrt{3} = 0$   
(D)  $x^2 + y^2 - 2x(\sqrt{3} + 1) - 2y(\sqrt{3} + 1) + 6 + 4\sqrt{3} = 0$ 

9. A circle of constant radius 'r' passes through origin O and cuts the axes of coordinates in points P and Q, then the equation of the locus of the foot of perpendicular from O to PQ is :

(A) 
$$(x^2 + y^2)(x^{-2} + y^{-2}) = 4r^2$$
  
(B)  $(x^2 + y^2)^2(x^{-2} + y^{-2}) = r^2$   
(C)  $(x^2 + y^2)^2(x^{-2} + y^{-2}) = 4r^2$   
(D)  $(x^2 + y^2)(x^{-2} + y^{-2}) = r^2$ 

- 10.  $S_1$ : If the length of tangent drawn from an external point P to the circle of radius r is  $\bullet$ , then area of triangle formed by pair of tangents and its chord of contact is  $\frac{rl^3}{r^2 + 1^2}$ .
  - **S**<sub>2</sub>: If the points where the lines  $a_1x + b_1y + c_1 = 0$  and  $a_2x + b_2y + c_2 = 0$  meet the co-ordinate axes are concyclic, then  $a_1c_1 = a_2c_2$
  - $S_3$ : A circle is inscribed in an equilateral triangle of side a, the area of any square inscribed in the

circle is  $\frac{a^2}{8}$ 

S<sub>4</sub>: The equation of the circle with origin as centre passing the vertices of an equilateral triangle whose median is of length 3a is  $x^2 + y^2 = 4a^2$ 

(A) FFTT (B) TTTF (C) TFFT (D) TTTT

#### **SECTION - II : MULTIPLE CORRECT ANSWER TYPE**

- 11. Consider the circle  $x^2 + y^2 10x 6y + 30 = 0$ . Let O be the centre of the circle and tangent at A(7, 3) and B(5, 1) meet at C. Let S = 0 represents family of circles passing through A and B, then -
  - (A) area of quadrilateral OACB = 4
  - (B) the radical axis for the family of circles S = 0 is x + y = 10
  - (C) the smallest possible circle of the family S = 0 is  $x^2 + y^2 12x 4y + 38 = 0$
  - **(D)** the coordinates of point C are (7, 1)

12. The centre of a circle passing through the points (0, 0), (1, 0) & touching the circle  $x^2 + y^2 = 9$  is :

**(A)**  $\left(\frac{3}{2}, \frac{1}{2}\right)$  **(B)**  $\left(\frac{1}{2}, \sqrt{2}\right)$  **(C)**  $\left(\frac{1}{2}, \frac{1}{2}\right)$  **(D)**  $\left(\frac{1}{2}, -\sqrt{2}\right)$ 



13. Point M moved on the circle  $(x - 4)^2 + (y - 8)^2 = 20$ . Then it broke away from it and moving along a tangent to the circle, cuts the x-axis at the point (-2, 0). The co-ordinates of a point on the circle at which the moving point broke away is

(A) 
$$\left(-\frac{3}{5}, \frac{46}{5}\right)$$
 (B)  $\left(-\frac{2}{5}, \frac{44}{5}\right)$  (C) (6, 4) (D) (3, 5)

- 14. If  $a \oplus 2 bm^2 + 2 d \oplus + 1 = 0$ , where a, b, d are fixed real numbers such that  $a + b = d^2$ , then the line  $\bigoplus x + my + 1 = 0$  touches a fixed circle
  - (A) which cuts the x-axis orthogonally
  - (B) with radius equal to b
  - (C) on which the length of the tangent from the origin is  $\sqrt{d^2 b}$
  - **(D)** none of these.
- 15. If the area of the quadrilateral formed by the tangents from the origin to the circle
  - $x^2 + y^2 + 6x 10y + c = 0$  and the radii corresponding to the points of contact is 15, then values of c is/are

#### SECTION - III : ASSERTION AND REASON TYPE

**16.** Consider the lines

L: (k+7)x - (k-1)y - 4(k-5) = 0 where k is a parameter

and the circle

 $C: x^2 + y^2 + 4x + 12y - 60 = 0$ 

Statement-I: Every member of L intersects the circle 'C' at an angle of 90°

Statement-II: Every member of L is tangent to the circle C.

(A) Statement-1 is true, statement-2 is true; statement-2 is correct explanation for statement-1.

(B) Statement-1 is true, statement-2 is true; statement-2 is NOT the correct explanation for statement-1.

- (C) Statement-1 is true, statement-2 is false.
- (D) Statement-1 is false, statement-2 is true.
- 17. Let C be a circle with centre 'O' and HK is the chord of contact of pair of the tangents from point A. OA intersects the circle C at P and Q and B is the midpoint of HK, then

Statement-I: AB is the harmonic mean of AP and AQ.

Statement-II : AK is the Geometric mean of AB and AO and OA is the arithmetic mean of AP and AQ.

- (A) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.
- (B) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.
- (C) Statement-1 is true, statement-2 is false.
- (D) Statement-1 is false, statement-2 is true.
- **18.** Statement-I: Let  $S_1: x^2 + y^2 10x 12y 39 = 0$ and  $S_3: 2x^2 + 2y^2 - 20x - 24y + 78 = 0$  $S_2: x^2 + y^2 - 2x - 4y + 1 = 0$

The radical centre of these circles taken pairwise is (-2, -3)

Statement-II : Point of intersection of three radical axis of three circles taken in pairs is known as radical centre

(A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.

- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True



- 19. Statement-I : Angle between the tangents drawn from the point P(13, 6) to the circle S :  $x^2 + y^2 6x + 8y 75 = 0$  is 90°. Statement-II : Point P lies on the director circle of S.
  - (A) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.
  - (B) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.
  - (C) Statement-1 is true, statement-2 is false.
  - (D) Statement-1 is false, statement-2 is true.
- 20. Statement-I : Only one normal can be drawn through the point P(2, -3) to the circle  $x^2+y^2-4x+8y-16=0$

Statement-II : Passing through any point lying inside a given circle only one normal can be drawn.

- (A) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.
- (B) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.
- (C) Statement-1 is true, statement-2 is false.
- **(D)** Statement-1 is false, statement-2 is true.

#### **SECTION - IV : MATRIX - MATCH TYPE**

21.		Column-I	Colun	nn-II
	<b>(A)</b>	If the straight line $y = kx \forall K \in I$ touches or passes outside the circle $x^2 + y^2 - 20y + 90 = 0$ then $ k $ can have the value	<b>(p)</b>	1
	<b>(B)</b>	Two circles $x^2 + y^2 + px + py - 7 = 0$ and $x^2 + y^2 - 10x + 2py + 1 = 0$ intersect each other orthogonally then the value of p is	<b>(q)</b>	2
	(C)	If the equation $x^2 + y^2 + 2\lambda x + 4 = 0$ and $x^2 + y^2 - 4\lambda y + 8 = 0$ represent real circles then the value of $\lambda$ can be	<b>(r)</b>	3
	<b>(D)</b>	Each side of a square is of length 4. The centre of the square is $(3, 7)$ . One diagonal of the square is parallel to $y = x$ . The possible abscissae of the vertices of the square can be	(s)	5
22.		Column – I	Colur	nn – II
	(A)	If $ax + by - 5 = 0$ is the equation of the chord of the circle $(x - 3)^2 + (y - 4)^2 = 4$ , which passes through (2, 3) and at the greatest distance from the centre of the circle, then $ a + b $ is equal to -	<b>(</b> p <b>)</b>	6
	(B)	Let O be the origin and P be a variable point on the circle $x^2 + y^2 + 2x + 2y = 0$ . If the locus of mid-point of OP is $x^2 + y^2 + 2gx + 2fy = 0$ , then the value of $(g + f)$ is equal to -	(q)	3
	(C)	The x-coordinates of the centre of the smallest circle which cuts the circle $x^2 + y^2 - 2x - 4y - 4 = 0$ and $x^2 + y^2 - 10x + 12y + 52 = 0$ orthogonally, is -	(r)	2
	<b>(D)</b>	If $\theta$ be the angle between two tangents which are drawn to the circle $x^2 + y^2 - 6\sqrt{3} x - 6y + 27 = 0$ from the origin, then $2\sqrt{3}$ tan $\theta$ equals to -	(s)	1



#### SECTION - V : COMPREHENSION TYPE 23. Read the following comprehension carefully and answer the questions. Let C be a circle of radius r with centre at O. Let P be a point outside C and D be a point on C. A line through P intersects C at Q and R, S is the midpoint of QR. 1. For different choices of line through P, the curve on which S lies, is (A) a straight line (B) an arc of circle with P as centre (C) an arc of circle with PS as diameter (D) an arc of circle with OP as diameter 2. Let P is situated at a distance 'd' from centre O, then which of the following does not equal the product (PQ) (PR)? (A) $d^2 - r^2$ (B) $PT^2$ , where T is a point on C and PT is tangent to C $(C)(PS)^2 - (QS)(RS)$ (D) $(PS)^2$ Let XYZ be an equilateral triangle inscribed in C. If $\alpha$ , $\beta$ , $\gamma$ denote the distances of D from vertices X, Y, Z respectively, 3. the value of product $(\beta + \gamma - \alpha) (\gamma + \alpha - \beta) (\alpha + \beta - \gamma)$ , is (C) $\frac{\alpha^3 + \beta^3 + \gamma^3 - 3\alpha\beta\gamma}{6}$ (D) None of these (B) $\frac{\alpha\beta\gamma}{2}$ **(A)**0 24. Read the following comprehension carefully and answer the questions. Two variable chords AB and BC of a circle $x^2 + y^2 = a^2$ are such that AB = BC = a, and M and N are The mid points of AB and BC respectively such that line joining MN а intersect the circle at P and Q where P is closer to AB and O is the centre of the circle C ∠OAB is -1. (A) 30° **(B)** 60° (C) 45° **(D)** 15° 2. Angle between tangents at A and C is -(A) 90° **(B)** 120° (C) 60° **(D)** 150° Locus of point of intersection of tangents at A and C is 3. **(B)** $x^2 + y^2 = 2a^2$ (C) $x^2 + y^2 = 4a^2$ (A) $x^2 + y^2 = a^2$ (D) $x^2 + y^2 = 8a^2$ 25. Read the following comprehension carefully and answer the questions. Consider the two quadratic polynomials $C_a: y = \frac{x^2}{4} - ax + a^2 + a - 2$ and $C: y = 2 - \frac{x^2}{4}$ 1. If the origin lies between the zeroes of the polynomial C<sub>a</sub> then the number of integral value(s) of 'a' is **(A)**1 **(B)**2 **(C)**3 (D) more than 3



- 2. If 'a' varies then the equation of the locus of the vertex of  $C_a$ , is (A) x-2y-4=0 (B) 2x-y-4=0 (C) x-2y+4=0 (D) 2x+y-4=0
- 3. For a = 3, if the lines  $y = m_1 x + c_1$  and  $y = m_2 x + c_2$  are common tangents to the graph of  $C_a$  and C then the value of  $(m_1 + m_2)$  is equal to (A)-6 (B)-3 (C) 1/2 (D) none

#### **SECTION - VI : INTEGER TYPE**

- 26. The lines 5x + 12y 10 = 0 and 5x 12y 40 = 0 touch a circle  $C_1$  of diameter 6 unit. If the centre of  $C_1$  lies in the first quadrant, find the equation of the circle  $C_2$  which is concentric with  $C_1$  and cuts of intercepts of length 8 on these lines.
- 27. If  $C_1 : x^2 + y^2 = (3 + 2\sqrt{2})^2$  be a circle and PA and PB are pair of tangents on  $C_1$  where P is any point on the director circle of  $C_1$ , then find the radius of smallest circle which touches  $C_1$  externally and also the two tangents PA and PB.
- 28. A ball moving around the circle  $x^2 + y^2 2x 4y 20 = 0$  in anti-clockwise direction leaves it tangentially at the point P(-2, -2). After getting reflected from a straight line it passes through the centre of the circle. Find the equation of this straight line if its perpendicular distance from P is  $\frac{5}{2}$ . You can assume that the angle of incidence is equal to the angle of reflection.
- 29. S is a circle having centre at (0, a) and radius b(b < a). A variable circle centred at ( $\alpha$ , 0) and touching circle S, meets the X-axis at M and N. A point  $P = \left(0, \pm \lambda \sqrt{a^2 b^2}\right)$  on the Y-axis, such that  $\angle MPN$  is a constant for any choice of  $\alpha$ , then find  $\lambda$ .
- 30. The ends A, B of a fixed straight line of length 'a' and ends A' and B' of another fixed straight line of length 'b' slide upon the axis of X & the axis of Y (one end on axis of X & the other on axis of Y). Find the locus of the centre of the circle passing through A, B, A' and B'.



# **ANSWER KEY**

#### EXERCISE - 1

۲

 1. C
 2. B
 3. A
 4. A
 5. C
 6. C
 7. C
 8. D
 9. A
 10. A
 11. B
 12. D
 13. B

 14. A
 15. A
 16. D
 17. C
 18. C
 19. B
 20. C
 21. D
 22. A
 23. B
 24. C
 25. D
 26. B

 27. D
 28. C
 29. D
 30. A
 31. A
 32. A
 33. A
 34. B
 35. A
 36. A
 37. C
 38. B
 39. A

 40. C

#### EXERCISE - 2 : PART # I

1.	ABCD	2.	AD	3.	AD	4.	BC	5.	AB	6.	ABC	7.	В	8.	ACD
9.	ACD	10.	ABC	11.	ABD	12.	AD	13.	AC	14.	AD	15.	AD	16.	AB
17.	BC	18.	С	19.	AC	20.	AB	21.	BCD	22.	CD				

#### PART - II

1. A 2. D 3. C 4. D 5. A 6. A 7. A 8. C 9. A 10. B 11. C 12. A 13. D

#### EXERCISE - 3 : PART # I

1.  $A \rightarrow r, s \ B \rightarrow s \ C \rightarrow p \ D \rightarrow q$  2.  $A \rightarrow q \ B \rightarrow p \ C \rightarrow r \ D \rightarrow s$  3.  $A \rightarrow p, q \ B \rightarrow p, q \ C \rightarrow q \ D \rightarrow q, s$ 4.  $A \rightarrow r \ B \rightarrow s \ C \rightarrow p \ D \rightarrow q$  5.  $A \rightarrow p, q, r \ B \rightarrow q, r \ C \rightarrow q, r, s \ D \rightarrow p, s$ 

#### PART - II

Comprehension #1: 1.	В	2.	Α	3.	А	<b>4.</b> B	5.	В						
Comprehension #2: 1.	В	2.	С	3.	А				Comprehension #3: 1.	В	2.	С	3.	D
Comprehension #4: 1.	В	2.	D	3.	A				Comprehension #5: 1.	В	2.	D	3.	В
Comprehension #6: 1.	D	2.	D	3.	Α				Comprehension #7: 1.	В	2.	С	3.	С
Comprehension #8: 1.	D	2.	В	3.	А									

#### EXERCISE - 5 : PART # I

 1. C
 2. A
 3. B
 4. D
 5. B
 6. A
 7. A
 8. A
 9. B
 10. D
 11. D
 12. D
 13. C

 14. C
 15. B
 16. C
 17. D
 18. D
 19. C
 20. B
 21. D
 22. C
 23. B
 24. C
 25. D
 26. A

 27. A
 28. C
 29. A
 29. A
 21. D
 22. C
 23. B
 24. C
 25. D
 26. A

#### PART - II

**1.** A **2.**  $OA=3(3+\sqrt{10})$  **3.**  $x^2+y^2+14x-6y+6=0$ ; **4.** 2px+2qy=r **5.** C **6.** A **7.** C **8.**  $2x^2+2y^2-10x-5y+1=0$  **9.** D **10.** B **11.** A **12.** B **13.** C **14.** (i) D, (ii) A, (iii) D **15.** B **16.** 8 **17.** 3 **18.** D **19.** 2 **20.** A **21.** D **22.** A **24.** B,C **25.** A,C **26.** A,B,C



#### **MOCK TEST**

- 1. B 2. D 3. C 4. B 5. B 6. D 7. A 8. C 9. C 10. C 11. ACD 12. B
- 13. BC 14. AC 15. AD 16. C 17. A 18. D 19. A 20. C
- **21.**  $A \rightarrow p,q,r \ B \rightarrow q,r \ C \rightarrow q,r,s \ D \rightarrow p,s$  **22.**  $A \rightarrow r \ B \rightarrow s \ C \rightarrow q \ D \rightarrow p$
- 23. 1. D 2. D 3. A 24. 1. B 2. C 3. C 25. 1. B 2. A 3. B
- **26.**  $x^2 + y^2 10x 4y + 4 = 0$  **27. 1 28.**  $(4\sqrt{3} 3)x (4 + 3\sqrt{3})y (39 2\sqrt{3}) = 0$
- **29.** 1 **30.**  $(2ax 2by)^2 + (2bx 2ay)^2 = (a^2 b^2)^2$

