

(iv) Distance versus time

Plot graph for

Sol.

Ex.1

(i) $V_2 - V_0 =$ Area of a Vs t graph for t = 0 to t = 2 sec $V_2 - 0 = 2 \times 2 \implies V_2 = 4 \text{ m/s}$



(ii) Since slope of a Vs t graph from t = 2 to 6 sec. is constant, we can observe its speed i.e. magnitude of its velocity is zero at = 4 sec. and after that magnitude of velocity increases in negative direction up to 4 m/s at the same rate. (iii) Displacement (x) Vs t

 $x_2 - x_0 = area of v vs t graph for t = 0, t = 2 sec$

Now $V_6 - V_2 = -2 \times 4 \implies V_6 = -4 \text{ m/s}$

$$x_2 - 0 = \frac{1}{2} (2) (4) \implies x_2 = +4 \text{ m}$$

 $x_4 - x_2 = \frac{1}{2} (4) (2) \implies x_4 = 8 \text{ m}$

also
$$x_6 - x_4 = \frac{1}{2} (-4) (2) = -4 m \implies \therefore x_6 = +4 m$$

$$d_2 - d_0 = \frac{1}{2} (2) (4) \implies d_2 = 4m$$

$$d_4 - d_2 = \frac{1}{2} (2) (4) \implies d_4 = 8m$$

Also
$$d_6 - d_4 = \left| \frac{1}{2} (2)(-4) \right| = 4 \implies d_6 = 12 \text{ m}$$

Ex.2 A drunkard walking in a narrow lane takes 5 steps forward and 3 steps backward, followed again by 5 steps forward and 3 steps backward, and so on. Each step is 1 m long and requires 1s. Plot the x-t graph of his motion. Determine graphically or otherwise how long the drunkard takes to fall in a pit 9 m away from the start.



From x-t graph time taken = 21 s Sol.

 $(5m-3m) + (5m-3m) + 5m = 9m \implies \text{total steps} = 21 \implies \text{time} = 21 \text{ s}$



Ex.3 On an open ground a motorist follows a track that turns to his left by an angle of 60° after every 500 m. Starting from a given turn, specify the displacement of the motorist at the third, sixth and eight turn. Compare the magnitude of displacement with the total path length covered by the motorist in each case.



Sol.

At III turn :

Displacement = $\overrightarrow{OA} + \overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{OC} = 500 \cos 60^\circ + 500 + 500 \cos 60^\circ$

$$500 \times \frac{1}{2} + 500 + 500 \times \frac{1}{2} = 1000 \text{ m from O to C}$$

Distance =
$$500 + 500 + 500 = 1500 \text{ m}$$
 So $\frac{\text{Displacement}}{\text{Distance}} = \frac{1000}{1500} = \frac{2}{3}$

At VI turn : + initial and final positions are same so displacement

= 0 and distance =
$$500 \times 6 = 3000 \text{ m}$$
 \therefore $\frac{\text{Displacement}}{\text{Distance}} = \frac{0}{3000} = 0$

At VIII turn : Displacement =
$$(500) \cos\left(\frac{60^{\circ}}{2}\right) = 1000 \times \cos 30^{\circ} = 1000 \times \frac{\sqrt{3}}{2} = 500 \sqrt{3} \text{ m}$$

Distance = $500 \times 84000 \text{ m}$ \therefore $\frac{\text{Displacement}}{\text{Distance}} = \frac{500\sqrt{3}}{4000} = \frac{\sqrt{3}}{8}$

Ex.4 A man walks on a straight road from his to a market 2.5 km away with a speed of 5 km/h. On reaching the market he initially turns and walks back with a speed of 7.5 km/h. What is the

(a) magnitude of average velocity and

(b) average speed of the man, over the interval of time (i) 0 to 30 min. (ii) 0 to 50 min (iii) 0 to 40 min

Sol. Time taken by man to go from his home to market,

$$t_1 = \frac{\text{distance}}{\text{speed}} = \frac{2.5}{5} = \frac{1}{2} \text{h}$$

Time taken by man to go from market to his home, $t_2 = \frac{2.5}{7.5} = \frac{1}{3}h$

Total time taken
$$= t_1 + t_2 = \frac{1}{2} + \frac{1}{3} = \frac{5}{6} h = 50 min$$

(i) 0 to 30 min

Ŀ

Average velocity =
$$\frac{\text{displacement}}{\text{time int erval}} = \frac{2.5}{\frac{30}{60}} = 5 \text{ km/h}$$
 towards market

Average speed = $\frac{\text{dis tan ce}}{\text{time int erval}} = \frac{2.5}{\frac{30}{60}} = 5 \text{ km/h}$



(ii) 0 to 50 min Total displacement = zero So average velocity = 0 So, average speed = $\frac{5}{50/60} = 6$ km/h Total distance travelled = 2.5 + 2.5 = 5 km (iii) 0 to 40 min Distance covered in 30 min (from home to market) = 2.5 km Distance covered in 10 min (from market to home) with speed 7.5 km/h = 7.5 × $\frac{10}{60} = 1.25$ km So, displacement = 2.5 - 1.25 = 1.25 km (towards market) Distance travelled = 2.5 + 1.25 = 3.75 km Average velocity = $\frac{1.25}{\frac{40}{60}} = 1.875$ km/h. (towards market)

Average speed =
$$\frac{3.75}{\frac{40}{60}}$$
 = 5.625 km/h.

Ex.5 If a body travels half its total path in the last second of its fall from rest, find :

(a) The time and

(b) heights of its fall.

Explain the physically unacceptable solution of the quadratic time equation. $(g = 9.8 \text{ m/s}^2)$

Sol. If the body falls a height h in time t, then

$$h = \frac{1}{2}gt^2$$
 [$u = 0$ as the body starts from rest]

....**(i)**

Now, as the distance covered in (t-1) second in $h' = \frac{1}{2}g(t-1)^2$ (ii)

So from Equation (i) and (ii) distance travelled in the last second.

$$h-h' = \frac{1}{2}gt^2 - \frac{1}{2}g(t-1)^2$$
 i.e., $h-h' = \frac{1}{2}g(2t-1)$

But according to given problem as $(h - h') = \frac{h}{2}$

$$\Rightarrow \left(\frac{1}{2}\right)\mathbf{h} = \left(\frac{1}{2}\right)\mathbf{g}(2t-1) \quad \text{or} \quad \left(\frac{1}{2}\right)\mathbf{g}t^2 = \mathbf{g}(2t-1) \text{ [as from equation (i) } \mathbf{h} = \left(\frac{1}{2}\right)\mathbf{g}t^2 \text{]}$$
$$\Rightarrow t^2 - 4t + 2 = 0 \quad \text{or} \quad t = \left[4 \pm \sqrt{(4^2 - 4\times)}\right]/2 \quad \Rightarrow t = 2 \pm \sqrt{2} \Rightarrow t = 0.59 \text{ s or } 3.41 \text{ s}$$

0.50 s is physically unacceptable as it gives the total time t taken by the body to reach ground lesser than one sec while according to the given problem time of motion must be greater than 1 s. so t = 3.41 s and $h = 1/2 \times (9.8) \times (3.41)^2 = 57$ m



- **Ex.6** A passenger is standing d distance away from a bus. The bus begins to move with constant acceleration a. To catch the bus, the passenger runs at a constant speed u towards the bus. What must be the minimum speed of the passenger so that he may catch the bus ?
- **Sol.** Let the passenger catch the bus after time t
 - The distance travelled by the bus, $s_1 = 0 + \frac{1}{2} at^2$...(i) and the distance travelled by the passenger $s^2 = ut + 0$...(ii) Now the passenger will catch the bus if $d + s_1 + s_2$...(iii) $\Rightarrow d + \frac{1}{2}at^2 = ut \Rightarrow \frac{1}{2}at^2 - ut + d = 0 \Rightarrow t = \frac{[u \pm \sqrt{u^2 - 2ad}]}{a}$

So the passenger will catch the bus if t is real, i.e., $u^2 \ge 2ad \implies u \ge \sqrt{2ad}$

So minimum speed of passenger for catching the bus is $\sqrt{2ad}$.

- **Ex.7** A driver takes 0.20 s to apply the brakes after he see a need for it. This is called the reaction time of the driver. If he is driving a car at a speed of 54 km/h and the brakes cause a deceleration of 6.0 m/s2, find the distance travelled by the car after he sees the need to put the brakes on ?
- Sol. Distance covered by the car after during the application of brakes by driver

$$s_1 = ut = \left(54 \times \frac{5}{18}\right) (0.2) = 15 \times 0.2 = 3.0 m$$

After applying the brakes ;

v = 0, u = 15 m/s, a = 6 m/s² s₂ = ?

Using
$$V_2 = u_2 - 2as \implies 0 = (15)^2 - 2 \times 6 \times s_2 \implies 12 s_2 = 255 \implies s_2 = \frac{255}{12} = 18.75 \text{ m}$$

Distance travelled by the car after driver sees the need for it $s = s_1 + s_2 = 3 + 18.75 = 21.75 \text{ m}$

Ex.8 Draw displacement time and acceleration – time graph for the given velocity time - graph



Sol. For
$$0 \le t \le 5 v \propto t^2$$
 and $a_1 = \text{constant } \frac{10}{5} = 2 \text{ ms}^{-2}$

for whole intervals s1 = Area under the curve = $\frac{1}{2} \times 5 \times 10 = 25$ m

For $5 \le t \le 10$, $v = 10 \text{ ms}^{-1} \implies a = 0$

for whole interval $s_2 =$ area under the curve $= 5 \times 10 = 50$ m

For $10 \le t \le 12$ v linearly decreases with time $\Rightarrow a_3 = -\frac{10}{2} = -5 \text{ ms}^{-2}$

for whole interval s_3 = Area under the curve = $\frac{1}{2} \times 2 \times 10 = 10$ m





- **Ex.9** A car accelerates from rest at a constat rate α for some time, after which it decelerates at a constant rate β , to come to rest. If total time elapsed is t evaluate (a) the maximum velocity attained and (b) the total distance travelled.
- Sol. (a) Let the car accelerates for time t_1 and decelerates for time t_2 then $t = t_1 + t_2$...(i) and corresponding velocity- time graph will be as shown in fig.

From the graph a = slope of line OA =
$$\frac{V_{max}}{t_1} \Rightarrow t_1 = \frac{V_{max}}{\alpha}$$

and b = - slope of line AB =
$$\frac{V_{max}}{t_2} \Rightarrow t_2 = \frac{V_{max}}{\beta}$$

$$\frac{V_{max}}{\alpha} + \frac{V_{max}}{\beta} = t \implies V_{max}\left(\frac{\alpha + \beta}{\alpha\beta}\right) = t \implies V_{max} = \frac{\alpha\beta t^2}{\alpha + \beta}$$

(b) Total distance = area under v-t graph =
$$\frac{1}{2} \times t \times v_{max} = \frac{1}{2} \times t \times \frac{\alpha\beta t}{\alpha+\beta} = \frac{1}{2} \left(\frac{\alpha\beta t^2}{\alpha+\beta} \right)$$

ms-1

- Note : This problem can also be solved by using equations of motion (v = u + at, etc).
- **Ex.10** At the height of 500m, a particle A is thrown up with $v = 75 \text{ ms}^{-1}$ and particle B is released from rest. Draw, acceleration time, velocity- time, speed- time and displacement- time graph of each particle.
- Sol.

For particle A:

Time of flight

$$-500 = +75t - \frac{1}{2} \times 10t^{2}$$

$$\Rightarrow t^{2} - 15t - 100 = 0$$

$$\Rightarrow t = 20 s$$

Time taken for A₁A₂

$$v = 0 = 75 - 10t \Rightarrow t = 7.5 s$$

Velocity at A₃, $v = 75 - 10 \times 20 = -125$

Height
$$A_2A_1 = 75 \times 7.5 - \frac{1}{2} (10) (7.5)^2 = 281.25 \text{ m}$$

For particle B

Time of flight

$$500 = \frac{1}{2}(10)t^2 \implies t = 10 s$$

Velocity at B2 $v = 0 - (10) (10) = -100 \text{ ms}^{-1}$







Ex.11 A rocket is fired upwards vertically with a net acceleration of 4 m/s2 and initial velocity zero. After 5 second its fuel is finished and it decelerates with g. At the highest point its velocity becomes zero. Then it accelerates downwards with acceleration g return back to ground. Plot velocity-time and displacement graphs for the complete journey. Take $g = 10 \text{ m/s}^2$.

Sol.





In the graphs, $v_A = at_{OA} = (4)(5) = 20 \text{ m/s } v_B = 0 = v_A - gt_{AB}$

:.
$$t_{AB} = \frac{v_A}{g} = \frac{20}{10} = 2s$$
 :. $t_{OAB} = (5+2)s = 7s$

Now, $s_{OAB} = area under v-t graph between 0 to 7 s = \frac{1}{2} (7) (20) = 70 m$

Now, $s_{OAB} = s_{BC} = \frac{1}{2}gt_{BC}^2$ $\therefore 70 = \frac{1}{2}(10)t_{BC}^2$ $\therefore \tau_{BX} = \sqrt{14} = 3.7 \text{ s}$ $\therefore t_{OABC} = 7 + 3.7 = 10.7 \text{ s}$

Also s_{OA} = area under v-t graph between OA = $\frac{1}{2}$ (5) (20) = 50 m

Ex.12 In the figure shown, the two projectile are fired simultaneously. Find the minimum distance between them during their flight.



Sol. Taking origin at A and x axis along AB Velocity of A w.r.t B

$$= 20\sqrt{3} \left(\cos 60\hat{i} + \sin 60\hat{j} \right) - 20 \left(\cos 150\hat{i} + \sin 150\hat{j} \right)$$

$$= 20\sqrt{3} \left(\frac{1}{2}\hat{i} + \frac{\sqrt{3}}{2}\hat{j} \right) - 20 \left(\frac{\sqrt{3}}{2}\hat{i} + \frac{1}{2}\hat{j} \right) = 20\sqrt{3}\hat{i} + 20\hat{j}$$

$$\tan \theta = \frac{20}{20\sqrt{3}} = \frac{1}{\sqrt{3}} \Longrightarrow \theta = 30^{\circ}$$

So
$$\frac{d_{\min}}{20} = \sin \theta = \sin 30^\circ = \frac{1}{2} \implies d_{\min} = 10m$$





- **Ex.13** Two ships A and B are 10 km apart on a line running south to north. Ship A farther north is streaming west at 20 km/h and hip B is streaming north at 20 km/h. What is their distance of closest approach and how long do they take to reach it ?
- Sol. Ships A and B are moving with the same speed 20 km/h in the direction shown in figure. It is a two dimensional, two body problem with zero acceleration. Let us find $\frac{1}{V_{BA}}$

Here
$$|_{V_{BA}}^{u}| = \sqrt{(20)^2 + (20)^2} = 20\sqrt{2} \text{ km/h}$$

i.e., $\stackrel{\mathbf{u}}{V}_{BA}$ is 20 $\sqrt{2}$ km/h at an angle of 45° from east

towards north. Thus, the given problem can be simplified as :



A is at rest and B is moving with V_{BA}^{u} in the direction shown in figure.

Therefore, the minimum distance between the two is

$$s_{min} = AC = AB \sin 45^\circ = 10 \left(\frac{1}{\sqrt{2}}\right) km = 5\sqrt{2} km$$

and the desired time is $t = \frac{BC}{|V_{BA}|} = \frac{5\sqrt{2}}{20\sqrt{2}} = \frac{1}{4} h = 15 min$

- **Ex.14** Three particles are projected from same point and their paths are as shown. Compare their horizontal and vertical component of velocities of projection
- Sol.

$$\begin{aligned} H_A &= H_B \ H_C \implies (u_y)_A = (u_y)_B > (u_y)_C \\ \therefore \qquad R_B > R_A \\ \text{i.e.,} \qquad (u_x \ u_y)_B > (u_x \ u_y)_A \ [Q \ \text{their} \ u_x \ \text{are equal}] \\ \therefore \qquad (u_x)_B > (u_x)_A \\ Also \qquad R_B = R_C \\ \text{i.e.,} \qquad (u_x \ u_y)_B \ = \ (u_x \ u_y)_C \\ \text{but} \qquad (u_y)_B \ > (u_y)_C \\ \therefore \qquad (u_x)_C > (u_x)_B > (u_x)_C \end{aligned}$$





Ex.15 Find range of projectile on the inclined plane which is projected perpendicular to the incline plane with velocity 20 m/s as shown in figure.
 Sol. β=37°

$$\alpha - \beta = 90^{\circ} \& \alpha = 90^{\circ} + \beta = 90^{\circ} + 37^{\circ}$$

$$\therefore \qquad \text{Range}, R = \frac{2(20)^{2} \sin(90^{\circ}) \cos(90^{\circ} + 37^{\circ})}{10 \times \cos^{2} 37^{\circ}}$$

$$= \frac{2(400)}{10(4/5)^{2}} \times \left(-\frac{3}{5}\right) \qquad \left[Q \cos(90^{\circ} + \theta) = -\sin\theta\right]$$

$$\therefore \qquad R = -75 \text{ m}$$

$$\therefore \qquad |R| = 75 \text{ m}$$
Here negative sign shown that particle strikes the plane along down the incline

Ex.16 A car goes out of control and slides off a steep embankment of height h at θ to the horizontal. It lands in a ditch at a distance T from the base. Find the speed at which the car leaves the slope.

 $(Take h = 12.5 m; R = 10 m; \theta = 45^{\circ})$

$$\Delta x = 10 = (u \cos 45^\circ)t \implies t = \frac{10\sqrt{2}}{u}$$
$$\Delta y = -(u \sin 45\circ)t - \frac{1}{2}gt^2 = -12.5$$
$$\implies \left(\frac{u}{\sqrt{2}} \times \frac{10\sqrt{2}}{u}\right) + \frac{1}{2}(10)\left(\frac{10\sqrt{2}}{u}\right)^2 = 12.5$$

Sol.

Ex.17 A particle is dropped from the top of a high building of height 360 m. The distance travelled by the particle in ninth second is $(g = 10 \text{ m/s}^2)$

(A) 85 m (C) 40 m **(B)** 60 m

(D) Can't be determined

Sol. Total time taken by particle to reach the ground T = $\sqrt{\frac{2H}{g}} = \sqrt{\frac{2 \times 360}{10}} = 6\sqrt{2} = 8.4484$ s

Distance travelled in 8 seconds = $\frac{1}{2}$ gt² = $\frac{1}{2}$ (10)(8)² = 320 m Therefore distance travelled in ninth second = 360 - 320 = 40 m



Ex.18 A particle is thrown vertically upward from the surface of the earth. Let T_p be the time taken by the particle to travel from a point P above the earth to its highest point and back to the point P. Similarly, let T_q be the time taken by the particle to travel from another point Q above the earth to its highest point and back to the same point Q. If the distance between the point P and Q is H, the expression for acceleration due to gravity in terms of T_p , T_Q and H is



Ex.19 An aeroplane is travelling horizontally at a height of 2000 m from ground. The aeroplane, when at a point P, drops a bomb to hit a stationary target Q on the ground. In order that the bomb hits the target, what angle q must the line PQ make with the vertical? $[g = 10 \text{ ms}^{-2}]$



Ex.20 A ball is thrown from the ground to clear a wall 3 m high at a distance of 6 m and falls 18 m away from the wall, the angle of projection of ball is





(D) $a_x v_x + a_y v_y > 0$

Н

g

Ex.21 A particle moves in XY plane such that its position, velocity and acceleration are given by

$$\hat{r} = x\hat{i} + y\hat{j}; \quad v = v_x\hat{i} + v_y\hat{j}; \quad a = a_x\hat{i} + a_y\hat{j}$$

which of the following condition is correct if the particle is speeding down ?

(A)
$$xv_x + yv_y < 0$$
 (B) $xv_x + yv_y > 0$ (C) $a_xv_x + a_yv_y < 0$

- **Sol.** For speeding down $a.v < 0 \implies a_x v_x + a_y v_y < 0$
- **Ex.22** A large number of particles are moving each with speed v having directions of motion randomly distributed. What is the average relative velocity between any two particles averaged over all the pairs ?

(A) v (B)
$$(\pi/4)v$$
 (C) $(4/\pi)v$ (D) Zero

Sol. Relative velocity,
$$\mathbf{v}_{r} = \begin{vmatrix} \mathbf{r} & \mathbf{r} \\ \mathbf{v}_{1} - \mathbf{v}_{2} \end{vmatrix}$$
 where $\mathbf{v}_{1} = \mathbf{v}_{2} = \mathbf{v}$

If angle between them be
$$\theta$$
, then $v_r = \sqrt{v^2 + v^2 - 2v^2 \cos \theta} = \sqrt{2v^2 (1 - \cos \theta)} = 2v \sin \left(\frac{\theta}{2}\right)$

Hence, average relative velocity
$$\mathbf{r}_{r} = \frac{\int_{0}^{2\pi} 2v \sin \frac{\theta}{2} d\theta}{\int_{0}^{2\pi} d\theta} = \frac{4v}{\pi}$$





Sol. Here $\frac{a_x}{a_y} = \frac{g \cot \theta}{g} = \frac{1}{\tan \theta} = \frac{u_x}{u_y} \Rightarrow$ Initial velocity & acceleration are opposite to each other.

 \Rightarrow Ball will return to point O.

- **Ex.24** Three point particles A, B and C are projected from same point with same speed at t = 0 as shown in figure. For this situation select correct statement(s).
 - (A) All of them reach the ground at same time.
 - (B) All of them reach the ground at different time.
 - (C) All of them reach the ground with same speed.
 - (D) All of them have same horizontal displacement when they reach the ground.
- Sol. Vertical component of initial velocities are different \Rightarrow reach the ground at different time.



Ex.25 to 27

A river of which 'd' with straight parallel banks flows due to North with speed u. A boat, whose speed is v relative to water, starts from A and crosses the river. If the boat is steered due West and u varies with y as $u = \frac{y(d-y)v}{d^2}$ then answer the following questions. E Ex.25 The time taken by boat to cross the river is (A) $\frac{d}{\sqrt{2}}$ (C) $\frac{d}{2v}$ 2d **(B)** $\frac{d}{d}$ **(D)** Absolute velocity of boat when it reaches the opposite bank is Ex.26 (C) $\frac{4}{3}$ v, towards West (D) v, towards East (A) $\frac{4}{3}$ v, towards East (B) v, towards West Ex.27 Equation of trajectory of the boat is (C) $y = \frac{x^2}{2d} - \frac{x^2}{3d^2}$ (D) $x = \frac{y^2}{2d} - \frac{y^2}{3d^2}$ (A) $y = \frac{x^2}{2d}$ (B) $x = \frac{y^2}{2d}$ Sol. 25 to 27 25. **(B)** Time taken = $\frac{d}{v_{u}} = \frac{d}{v}$ 26. **(B)** At y = d, u = 0 so absolute velocity of boat = v towards West. 27. **(D)** For boat (w.r.t. ground) $v_y = v$, $v_x = u = \frac{y(d-y)}{d^2}v \implies \frac{dy}{dt} = v$ and $\frac{y(d-y)}{d^2}v \implies \frac{dx}{dt} = \frac{y(d-y)}{d^2}v$ $\frac{dx}{dy} = \frac{y(d-y)}{d^2} \implies \int_{0}^{x} dx = \int_{0}^{y} dx \frac{(yd-y^2)}{d^2} dy \implies x = \frac{y^2}{2d} - \frac{y^3}{3d^2}$ As shown in the figure there is a particle of mass $\sqrt{3}$ kg, is projected with speed 10 m/s at an angle 30° with Ex.28 horizontal (take $g = 10 \text{ m/s}^2$) then match the following 10 m/s





KINEMATICS

Column-I
(A) Average velocity (in m/s) (P)
$$\frac{1}{2}$$

during half of the time of flight, is
(B) The time (in sec) after which the angle (Q) $\frac{5}{2}\sqrt{13} \frac{5}{2}\sqrt{13}$
between velocity vector and initial velocity
vector becomes $\pi/2$, is
(C) Horizontal range (in m), is (R) $5\sqrt{3}$
(D) Change in linear momentum (in N-s) (S) At an angle of $\tan -1\left(\frac{1}{2\sqrt{3}}\right)$ from horizontal
when particle is at highest point, is
(I) 2
Sol. For (A): $v_{sv} = \sqrt{(v_{svx})^2 + (v_{svy})^2} = \sqrt{(10\cos 30^\circ)^2 + (\frac{10\cos 30^\circ + 0}{2})^2} = \sqrt{75 + \frac{25}{4}} = \frac{5}{2}\sqrt{13}$ m/s
Angle with horizontal $\theta = \tan^{-1} = (\frac{v_{svy}}{v_{swx}}) = \tan^{-1}(\frac{5/2}{5\sqrt{3}}) = \tan^{-1}(\frac{1}{2\sqrt{3}})$
For (B): By using $\frac{1}{v} = \frac{u}{u} + \frac{1}{at}$ We have $\frac{u}{gt} = \sin 30^\circ \Rightarrow t = \frac{10}{(10)(1/2)} = 2$
For (C): Horizontal range (R) $= \frac{u^2 \sin 2\theta}{g} = \frac{100 \times \sqrt{3}/2}{10} = 5\sqrt{3}$ m.
For (D): Change in linear momentum $= mu_y = \sqrt{3} \times 10\sin 30^\circ = 5\sqrt{3}$ N-s

- **Ex.29** Find the reaction between acceleration of blocks a_1 , a_2 and a_3 .
- Sol. $x_1 + x_2 + x_3 + x_4 = l$ $\Rightarrow \quad x_1 + x_2 + x_3 + x_4 = l$ $\Rightarrow \quad x_1 + a_2 + a_3 + a_4 = 0$ $\Rightarrow \quad a_1 + 2a_2 + a_3 = 0$ $\Rightarrow \quad \left(\frac{u}{\sqrt{2}} \times \frac{10\sqrt{2}}{u}\right) + \frac{1}{2}(10) \left(\frac{10\sqrt{2}}{u}\right)^2 = 12.5$ $\Rightarrow \quad \left(\frac{u}{\sqrt{2}} \times \frac{10\sqrt{2}}{u}\right) + \frac{1}{2}(10) \left(\frac{10\sqrt{2}}{u}\right)^2 = 12.5$





Ex.30 Two stones A and B are projected simultaneously as shown in figure. It has been observed that both the stones reach the ground at the same place after 7 sec of their projection. Determine difference in their vertical components of initial velocities in m/g. $(g = 9.8 \text{ m/s}^2)$



- Sol. In time of flight i.e., 7 s, the vertical displacement of A is zero and that of B is 49 m so for relative motion of B w.r.t. A $(u_2 \sin \theta_2 - u_1 \sin \theta_1) \times 7 = 49 \implies u_2 \sin \theta_2 - u_1 \sin \theta_1 = 7$ m/s
- **Ex.31** An aeroplane pilot wishes to fly due west. A wind of 100 km/h is blowing toward the south
 - (A) What is the speed of the plane with respect to ground ?
 - (B) If the airspeed of the plane (its speed in still air) is 300 km/h, in which direction should the pilot head?
- Sol. (A) Given,

Velocity of air with respect of ground $V_{A/G} = 100$ km/hr

Velocity of plane with respect to air $V_{P/A}^{u} = 300 \text{ km/hr}$



(B) As the plane is to move towards west, due to air in south direction, air will try drift the plane in south direction., air will try to drift the plane in south direction. Hence, the plane has to make an angle θ towards north-west, south west direction, in order to reach at point on west.

 $\overset{\mathbf{u}}{V}_{P/A} = \overset{\mathbf{u}}{V}_{P/G} - \overset{\mathbf{u}}{V}_{A/G}$ and $\overset{\mathbf{u}}{V}_{P/A} = \sin \theta - V_{AG}$

Ex.32 Ram crossing a 2.5 m wide conveyor belt moves with a speed of 1.6 m/s. The conveyor belt moves at uniform speed of 1.2 m/s.

(A) If the Ram walks straight across the belt, determine the velocity of the Ram relative to an observer standing on ground.

(B) If Shayam has same speed on a still conveyor belt, and is to reach directly across the same moving conveyor belt. At what should he walk ?





Sol. (A) If you walk across a conveyor belt while the conveyor belt takes you along the length, you will not be able to move directly across the conveyor belt, but will end up down the length.

Here the velocity of the Ram will be net effect of his own motion of conveyor belt. The velocity of the Ram relative to the conveyor belt v_{re} , is same as velocity of Ram if conveyor belt was still,

v_c is the velocity of the conveyor belt

we need to find $\boldsymbol{v}_{\boldsymbol{r}}$, the velocity of the Ram relative to the Earth.

Writing Equation of net motion $v_r = v_{rc} + vc$.

three vectors are shown in figure. The quantity v_{rc} is due y ; v_{c} is due x ; and the vector sum of the two v_{r} , is at an angle θ as defined in figure

the speed v_r of the Ram relative to the Earth is

$$\mathbf{v}_r = \sqrt{\mathbf{v}_{s/c}^2 - \mathbf{v}_c^2}$$

(B) To go straight across the conveyor belt he has to walk at some angle.

Writing Equation of net motion $v_s = v_{s/c} + v_c$

three vectors are shown in figure

As in part , we know v_c and the magnitude of the vector $v_{s/c}$ and we want v_c to be directed across the conveyor belt

$$\upsilon_{_{\rm S}}=\sqrt{v_{_{_{\rm S/c}}}^2-v_{_{_{\rm C}}}^2}$$

Ex.33 A river flows due to south with a speed of 2.0 m/s. A man steers a motorboat across the diver: his velocity relative to the water is 4 m/s due east. The river is 800 m wide.

(A) What is his velocity (magnitude direction) relative to the earth?

(B) How much time is required to cross the river?

(C) How far south of his starting point will be reach the opposite bank ?

Sol. Velocity of river (i.e., speed of river w.r.t. earth) $\int_{U_{rc}}^{1} = 2 \text{ m/s}$

Width of the river = 800 m



According to the given statement the diagram will be as given

When two vectors are acting at an angle of 900, their resultant can be obtained by phythagorous theorem,

$$v_{be} = \sqrt{v_{br}^2 + v_{re}^2} = \sqrt{16 + 4} = \sqrt{20} = 4.6 \text{ m/s}$$

To find direction, we have

$$\tan \theta = \frac{\upsilon_{re}}{\upsilon_{br}} = \frac{2}{4} = \frac{1}{2} \qquad \Rightarrow \qquad \theta = \tan^{-1}\left(\frac{1}{2}\right)$$



(A)

- (B) Time taken to cross the river = $\frac{\text{Displacement of boat w.r.t. river}}{\text{Velocity of boat w.r.t. river}} \implies \frac{800}{4} = 200 \text{ s}$
- (C) Desired position on other side is A, but due to current of river boat is drifted to position B. To find out this drift we need time taken in all to cross the river (200s) and speed of current (2 ms⁻¹). So distance AB = Time taken × speed of current = 200 × 2 = 400 m. Hence, the boat is drifted but 400 m away from position A.
- **Ex.34** A person standing on a road has to hold his umbrella at 60° with the vertical to keep the rain away. He throws the umbrella and starts running at 20 ms⁻¹. He finds that rain drops are falling on him vertically. Find the speed of the rain drops with respect to

(A) the road, and

(B) the moving person.

Sol. Given $\theta = 60^{\circ}$ and velocity person

$$V_P = OA = 20 \text{ ms}^-$$

This velocity is the same as the velocity of person w.r.t. ground. First of all let's see how the diagram works out.

 $v_{rP} = OB = velocity of rain w.r.t. the person$



 $v_r = OC = velocity of rain w.r.t. earth <math>v_{rp}$ is along OB OB as a person has to hold umbrella at an angle with vertical which is the angle between velocity of rain and velocity of rain w.r.t. the person.

Values of $\frac{1}{U_{r}}$ and $\frac{1}{U_{r}}$ can be obtained by using simple trigonometric relations.

(A) Speed of rain drops w.r.t. earth = $\frac{1}{U_r} = \frac{U_r}{OC}$

From
$$\triangle OCM$$
, $\frac{CB}{OC} = \sin 60^\circ \Rightarrow OC = \frac{CB}{\sin 60^\circ}$

$$\frac{20}{\sqrt{3/2}} = \frac{40}{\sqrt{3}} = \frac{40\sqrt{3}}{3} \text{ ms}^{-1}$$

Speed of rain w.r.t. the person $U_{rP} = OB$

From
$$\triangle OCM$$
, $\frac{OB}{CB} = \cot 60^{\circ}$

$$\Rightarrow \text{ OB} = \text{CB } \cot 60^\circ = \frac{20}{\sqrt{3}} = \frac{20\sqrt{3}}{3} \text{ ms}^{-1}$$



(B)

- **Ex.35** A person walk up a stationary escalator in t1 second. If he remains stationary on the escalator, then it can take him up t2 second. If the length of the escalators is L, then
 - (A) Determine the speed of man with respect to the escalator.
 - (B) Determine the speed of the escalator
 - (C) How much time would it take him to walk up the moving escalators?

Sol.

(A) As the escalators is stationary, so the distance covered in t_1 second is L which is the length of the escalator.

Speed of the man w.r.t. the escalator $v_e = \frac{L}{t_1}$

(B) When the man is stationary, by taking man as reference point the distance covered by the escalator is L in time t_2 .

Speed of escalator $v_e = \frac{L}{t_2}$

(C) Speed of man w.r.t. the ground

$$\mathbf{v}_{\mathrm{m}} = \mathbf{v}_{\mathrm{me}} + \mathbf{v}_{\mathrm{e}}$$

$$\Rightarrow \qquad \mathbf{v}_{\mathrm{m}} = \frac{\mathrm{L}}{\mathrm{t}_{1}} + \frac{\mathrm{L}}{\mathrm{t}_{2}} = \mathrm{L}\left[\frac{1}{\mathrm{t}_{2}} + \frac{1}{\mathrm{t}_{2}}\right] = \mathrm{L}\left[\frac{\mathrm{t}_{1} + \mathrm{t}_{2}}{\mathrm{t}_{1}\mathrm{t}_{2}}\right] \qquad \Rightarrow \qquad \mathrm{L} = \upsilon_{\mathrm{m}}\left[\frac{\mathrm{t}_{1}\mathrm{t}_{2}}{\mathrm{t}_{1} + \mathrm{t}_{2}}\right]$$

 $\left\lfloor \frac{t_1 t_2}{t_1 + t_2} \right\rfloor$ is the time taken by the man to walk up the moving escalator.

- **Ex.36** A man wants to cross a river 500 m wide. Rowing speed of the man relative to water is 3 km/hr and river flows at the speed of 2 km/hr. If man's walking speed on the shore is 5 km/hr, then in which direction he should starts rowing in order to reach the directly opposite point on the bank in shortest time.
- Sol. Let he should starts at an angle θ with normal hence

$$\hat{v}_{m} = (u - v \sin \theta)\hat{i} + v \cos \theta \hat{j}$$

Here $\frac{1}{V_m}$ = velocity of the man relative to water

$$u = velocity of water$$



Hence time taken by the man to cross the river is $t_1 = \frac{0.5}{v \cos \theta}$

... Drift of the man to cross the river is

 $x = (u - v \sin \theta) t1$

$$\mathbf{x} = (\mathbf{u} - \mathbf{v}\sin\theta) \ \frac{0.5}{\mathbf{v}\cos\theta}$$

Time taken by the man to cover this distance is

$$t_2 = \frac{0.5\left(\frac{u \sec \theta}{v} \tan \theta\right)}{5} = 0.1\left(\frac{u}{v} \sec \theta - \tan \theta\right)$$



Therefore,

total time $T = t_1 + t_2$

$$\Rightarrow \qquad T = \frac{0.5}{v} \sec \theta + \frac{0.1u}{v} \sec \theta - 0.1 \tan \theta$$

Putting the value of u and v, we get

$$T = \frac{0.5}{3} \sec \theta + \frac{0.1 \times 2}{3} \sec \theta - 0.1 \tan \theta$$

 $= \frac{0.7}{3} \sec \theta - 0.1 \tan \theta \qquad \Rightarrow \qquad \frac{dt}{d\theta} = \frac{0.7}{3} \sec \theta \tan \theta - 0.1 \sec^2 \theta$

for T to be minimum

 $\frac{dt}{d\theta} = 0 \implies \sin \theta = (3/7) \implies \theta = \sin^{-1}(3/7)$

Ex.37 A large heavy box is sliding without friction down a smooth plane of inclina-

tion θ . From a point P on the bottom of a box, a particle is projected inside the box. The initial speed of the particle with respect to box. The initial speed of the particle with respect to box is u and the direction of projection makes an angle α with the bottom as shown in figure.



- (a) Find the distance along the bottom of the box between the point of projection P and the point Q where the particle lands. (Assume that the particle does not hit any other surface of the box. Neglect air resistance).
- (b) If the horizontal displacement of the particle as seen by an observer on the ground is zero, find the speed of the box with respect to the ground at the instant when the particle was projected.
- Sol. (a) u is the relative velocity of the particle with respect to the box. Resolve u.

u, is the relative velocity of the particle with respect to the box in x- direction.

uy is the relative velocity with respect to the box in y- direction, therefore this is the vertical velocity of the particle with respect to ground also.

Y- direction motion (Taking relative terms w.r.t. box)

- $u_v = + u \sin \alpha$
- $a_v = -g \cos \theta$

$$s_v = 0$$

$$s_y = u_y t + \frac{1}{2} a_y t^2 \implies 0 = (u \sin \alpha) t - \frac{1}{2} g \cos \theta \times t^2 \implies t = \frac{2u \sin \alpha}{g \cos \theta}$$

x-direction motion (Taking relative terms w.r.t. box)

$$u_{v} = +\cos \alpha$$
; $a_{v} = 0$

$$s_x = u_y t + \frac{1}{2} a_y t^2 \implies s_x = u \cos \alpha \times \frac{2u \sin \alpha}{g \cos \theta} = \frac{u^2 \sin 2\alpha}{g \cos \theta}$$



(b) For the observer (on ground) to see the horizontal displacement to be zero, the distance travelled by the box in time

 $\left(\frac{2u\sin 2\alpha}{g\cos\theta}\right)$ should be equal to the range of the particle w.r.t. box.

Let the speed of the box at the time projection of particle be U. Then for the motion of box with respect to ground.

$$u_{x} = -U; a_{x} = -g\sin\theta; t = \frac{2u\sin\alpha}{g\cos\theta}; s_{x} = \frac{-u^{2}\sin2\alpha}{g\cos\theta} \implies s_{x} = u_{x}t + \frac{1}{2}a_{x}t^{2}$$

$$-u^{2}\sin2\alpha = -u(2u\sin\alpha) - \frac{1}{2}a\sin\theta(2u\sin\alpha)^{2}$$

$$\frac{-U\sin 2\alpha}{g\cos\theta} = -U\left(\frac{2U\sin\alpha}{g\cos\theta}\right) - \frac{1}{2}g\sin\theta\left(\frac{2U\sin\alpha}{g\cos\theta}\right)$$

on solving we get

$$U = \frac{u\cos(\alpha + \theta)}{\cos\theta}$$

Ex	xercise # 1	Sir	ngle Correct Choice	Type Questions]
1.	A particle moves in strai 4 m/s for another 20 sec a particle?	ght line in same direction fo nd finally moves with veloci	r 20 seconds with velocity 3 ty 5 m/s for next 20 seconds.	m/s and then moves with velocity What is the average velocity of the
	(A) 3 m/s	(B) 4 m/s	(C) 5 m/s	(D) Zero
2.	A bird moves from poin the bird is :-	nt $(1, -2, 3)$ to $(4, 2, 3)$. It	f the speed of the bird is 10) m/s, then the velocity vector of
	(A) $5\left(\hat{i}-2\hat{j}+3\hat{k}\right)$		(B) $5\left(4\hat{i}+2\hat{j}+3\hat{k}\right)$	
	(C) $0.6\hat{i} + 0.8\hat{j}$		(D) $6\hat{i} + 8\hat{j}$	
3.	A particle is moving in making 60° with positiv	x–y–plane at 2 m/s along ve x–axis. Its average acce	x-axis. 2 seconds later, its leration for this period of	s velocity is 4 m/s in a direction motion is:-
	(A) $\sqrt{5}$ m/s ² , along y-	axis	(B) $\sqrt{3}$ m/s ² , along y-a	xis
	(C) $\sqrt{5}$ m/s ² , along at	60° with positive x–axis	(D) $3m/s^2$, at 60° with p	ositive x–axis.
4.	The coordinates of a mov	ving particle at time t are give	$en by x = ct^2 and y = bt^2. The$	speed of the particle is given by :-
	(A) $2t(c+b)$	$\textbf{(B)} \ 2t\sqrt{c^2-b^2}$	(C) $t\sqrt{c^2+b^2}$	(D) $2t\sqrt{c^2+b^2}$
5.	The velocity of a parti x–coordinate of the part of particle is zero?	cle moving along x-axis i ticle in metres. Find the n	s given as $v = x^2 - 5x + agnitude$ of acceleration of	4 (in m/s) where x denotes the of the particle when the velocity
	(A) 0 m/s^2	(B) 2 m/s ²	(C) 3 m/s^2	(D) None of these
6 .	A, B, C and D are points descend through AB, BC	in a vertical line such that AI C and CD are in the ratio :-	B = BC = CD. If a body falls	from rest from A, then the times of
	(A) $1:\sqrt{2}:\sqrt{3}$		(B) $\sqrt{2}$: $\sqrt{3}$: 1	
	(C) $\sqrt{3}$:1: $\sqrt{2}$		(D) 1 : $(\sqrt{2} - 1) : (\sqrt{3} - 1)$	$\sqrt{2}$)
7.	A body starts from rest next 10 s is x_2 and the	and is uniformly accelerated as the 10 s is x ₃ . Then x ₁ :	ted for 30 s. The distance $x_2 : x_3$ is the same as:-	travelled in the first 10 s is x_1 ,
	(A) 1 : 2 : 4	(B) $1:2:5$	(C) 1 : 3 : 5	(D) 1 : 3 : 9
8.	A particle has an initial v second will be equal to :	velocity of $(3\hat{i} + 4\hat{j})$ m/s and	d a constant acceleration of	$(4\hat{i} - 3\hat{j})$ m/s ² . Its speed after one
	(A) 0	(B) 10 m/s	(C) $5\sqrt{2}$ m/s	(D) 25 m/s
9.	A particle is projected of the particle at any tir	vertically upwards and in ne t will be :-	t reaches the maximum he	eight H in T seconds . The height
	(A) $H - g(t - T)^2$	(B) $g(t-T)^2$	(C) $H - \frac{1}{2}g(t-T)^2$	(D) $\frac{g}{2}(t-T)^2$
10.	A particle is projected ve continues to move up. If ground is :-	ertically upwards from a point f it takes further t_2 time to re	nt A on the ground. It takes t each the ground from point	time to reach a point B but it still B then height of point B from the
	(A) $\frac{1}{2}g(t_1 + t_2)^2$	(B) gt_1t_2	(C) $\frac{1}{8}g(t_1 + t_2)^2$	(D) $\frac{1}{2}$ gt ₁ t ₂



- 11. A parachutist drops freely from an aeroplane for 10 s before the parachute opens out. Then he descends with a net retardation of 2.5 m/s². If he bails out of the plane at a height of 2495 m and $g = 10 \text{ m/s}^2$, hit velocity on reaching the ground will be :-
- (A) 5 m/s **(B)** 10 m/s (C) 15 m/s (D) 20 m/s 12. With what speed should a body be thrown upwards so that the distances traversed in 5th second and 6th second are equal?

(A) 58.4 m/s

Initially car A is 10.5 m ahead of car B. Both start moving at time t=0 in the same direction along a straight 13. line. The velocity time graph of two cars is shown in figure. The time when the car B will catch the car A, will be:-10 m/s _____ car A

(C) $\sqrt{98}$ m/s

(B) $t = 2\sqrt{5}$ sec (A) t = 21 sec (D) None of these (C) 20 sec

(B) 49 m/s

14. The velocity – time graph of a linear motion is shown in figure. The displacement & distance from the origin after 8 sec. is :-



(A) 5 m, 19m

(B) 16 m,22m

(C) 8 m, 19m

(D) 6 m, 5m

(D) 98 m/s

15. Which of the following velocity-time graph shows a realistic situation for a body in motion :-



16. A man moves in x-y plane along the path shown. At what point is his average velocity vector in the same direction as his instaneous velocity vector. The man starts from point P.

(A)A

- **(B)**B (C) C
- **(D)**D



17. The graph between the displacement x and time t for a particle moving in a straight line is shown in figure. During the interval OA, AB, BC and CD, the acceleration of the particle is :

	OA,	AB,	BC,	CD
A)	+	0	+	+
B)	-	0	+	0
C)	+	0	_	+
D)	- / -	0	0	+





18. A ball is thrown vertically upwards. Which of the following plots represents the speed-time graph of the ball during its flight if the air resistance is ignored :-



19. The velocity – time graph of a body falling from rest under gravity and rebounding from a solid surface is represented by which of the following graphs?



- 20. Which of the following situation is represented by the velocity-time graph as shown in the diagram :-
 - (A) A stone thrown up vertically, returning back to the ground
 - (B) A car decelerating at constant rate and then accelerating at the same rate
 - (C) A ball falling from a height and then bouncing back
 - **(D)** None of the above

21. The acceleration–time graph of a particle moving along a straight line is as shown in figure. At what time the particle acquires its initial velocity?

- (A) 12 sec
- **(B)** 5 sec
- (C) 8 sec
- **(D)** 16 sec
- 22. A river is flowing from west to east at a speed of 5 meters per minute. A man on the south bank of the river, capable of swimming at 10 meters per minute in still water, wants to swim across the river in the shortest time. He should swim in a direction :-

(A) Due north (B) 30° east of north

(C) 30° north of west

(D) 60° east of north

t(s)

 $a(m/s)^2$

10

- 23. A man starts running along a straight road with uniform velocity $\mathbf{u}^{r} = \mathbf{u}\hat{i}$ feels that the rain is falling vertically down along \hat{j} . If he doubles his speed he finds that the rain is coming at an angle θ to the vertical . The velocity of rain with respect to the ground is :-
 - (A) $u\hat{i} u \tan \theta \hat{j}$ (B) $u\hat{i} \frac{u}{\tan \theta} \hat{j}$ (C) $u \tan \theta \hat{i} u\hat{j}$ (D) $\frac{u}{\tan \theta} \hat{i} u\hat{j}$
- 24. Raindrops are falling vertically with a velocity 10m/s. To a cyclist moving on a straight road the rain drops appear to be coming with a velocity of 20m/s. The velocity of cyclist is :-
 - (A) 10m/s (B) $10\sqrt{3}$ m/s (C) 20 m/s (D) $20\sqrt{3}$ m/s



- 25. A boat moving towards east with velocity 4 m/s with respect to still water and river is flowing towards north with velocity 2 m/s and the wind is blowing towards north with velocity 6 m/s. The direction of the flag blown over by the wind hoisted on the boat is :(A) North-west (B) South-east (C) tan⁻¹(1/2) with east (D) North
- 26. A man is crossing a river flowing with velocity of 5m/s. He reaches a point directly across at distance of 60 m in 5s. His velocity in still water should be :-



27. A river is flowing from east to west at a speed of 5 m/min. A man on south bank of river, capable of swimming 10 m/min in still water, wants to swim across the river in shorter time; he should swim :-

(A) Due north	(B) Due north–east
(C) Due north–east with double the speed of river	(D) None of the above

28. A boat which has a speed of 5km per hour in still water crosses a river of width 1km along the shortest possible path in fifteen minutes. The velocity of the river water in km per hour is :-

(A) I (D) 2 (C) 3 (D)	(A) 1	(B) 2	(C) 3	(D) √4
-----------------------	--------------	--------------	--------------	---------------

29. From a motorboat moving downstream with a velocity 2 m/s with respect to river, a stone is thrown. The stone falls on an ordinary boat at the instant when the motorboat collides with the ordinary boat. The velocity of the ordinary boat with respect to the river is equal to zero. The river flow velocity is given to be 1 m/s. The initial velocity vector of the stone with respect to earth is :-



(A) 2i+20j

30.

Two particles P and Q are moving with velocities of $(\hat{i} + \hat{j})$ and $(-\hat{i} + 2\hat{j})$ respectively. At time t = 0,

(C) 3i + 50j

(D) 2i + 50j

P is at origin and Q is at a point with position vector $(2\hat{i} + \hat{j})$. Then the shortest distance between P & Q is :-

(A)
$$\frac{2\sqrt{5}}{5}$$
 (B) $\frac{4\sqrt{5}}{5}$ (C) $\sqrt{5}$ (D) $\frac{3\sqrt{5}}{5}$
The total speed of a projectile at its greatest height is $\sqrt{\frac{6}{5}}$ of its speed when it is at half of i

(B) 3i + 40j

31. The total speed of a projectile at its greatest height is $\sqrt{\frac{6}{7}}$ of its speed when it is at half of its greatest height The angle of projection will be :-(A) 60° (B) 45° (C) 30° (D) 50°



32. Graphs I and II give coordinates x(t) and y(t) of a particle moving in the x-y plane. Acceleration of the particle is constant and the graphs are drawn to the same scale. Which of the vector shown in options best represents the acceleration of the particle :-



- 33. Particle is dropped from the height of 20m on horizontal ground. There is wind blowing due to which horizontal acceleration of the particle becomes 6 ms⁻². Find the horizontal displacement of the particle till it reaches ground. (A) 6m
 (B) 10 m
 (C) 12m
 (D) 24 m
- 34. A particle is projected from a horizontal plane (x-z plane) such that its velocity vector at time t is given by $\stackrel{r}{v} = a\hat{i} + (b ct)\hat{j}$. Its range on the horizontal plane is given by :-

(A)
$$\frac{ba}{c}$$
 (B) $\frac{2ba}{c}$ (C) $\frac{3ba}{c}$ (D) None

35. A projectile is projected at an angle (α >45°) with an initial velocity u. The time t, at which its magnitude of horizontal velocity will equal the magnitude of vertical velocity is :-

(A)
$$t = \frac{u}{g}(\cos\alpha - \sin\alpha)$$
 (B) $t = \frac{u}{g}(\cos\alpha + \sin\alpha)$ (C) $t = \frac{u}{g}(\sin\alpha - \cos\alpha)$ (D) $t = \frac{u}{g}(\sin^2\alpha - \cos^2\alpha)$.

36. A particle is projected from a tower as shown in figure, then the distance from the foot of the tower where it will strike the ground will be :- (take $g = 10 \text{ m/s}^2$)



37. A particle is dropped from a height h. Another particle which was initially at a horizontal distance 'd' from the first, is simultaneously projected with a horizontal velocity 'u' and the two particles just collide on the ground. The three quantities h. d and u are related to :-

(A)
$$d^2 = \frac{u^2 h}{2 g}$$
 (B) $d^2 = \frac{2 u^2 h}{g}$ (C) $d = h$ (D) $g d^2 = u^2 h$

38. A block is dragged on a smooth plane with the help of a rope which moves with a velocity v as shown in figure. The horizontal velocity of the block is :









Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141

- 47.A particle starts moving along a circle of radius $(20/\pi)$ m with constant tangential acceleration. If the velocity of the particle is 50 m/s at the end of the second revolution after motion has began, the tangential acceleration in m/s² is :(A) 1.6(B) 4(C) 15.6(D) 13.2
- 48. A particle is kept fixed on a turntable rotating uniformly. As seen from the ground, the particle goes in a circle, its speed is 20 cm/s and acceleration is 20 cm/s². The particle is now shifted to a new position to make the radius half of the original value. The new values of the speed and acceleration will be
 (A) 10 cm/s, 10 cm/s²
 (B) 10 cm/s, 80 cm/s²
 (C) 40 cm/s, 10 cm/s²
 (D) 40 cm/s, 40 cm/s²
- 49. The second's hand of a watch has length 6 cm. Speed of end point and magnitude of difference of velocities at two perpendicular positions will be

R

(A) $2\pi \& 0 \text{ mm/s}$	(B) $2\sqrt{2} \pi \& 4.44 \text{ mm/s}$	
(C) $2\sqrt{2} \pi \& 2\pi \text{ mm/s}$	(D) $2\pi \& 2\sqrt{2}\pi \text{ mm/s}$	
A particle A moves along a circle of radius R= vector r relative to the point O (figure) rotates	=50 cm so that its radius s with the constant angular	r

velocity ω =0.40 rad/s. Then modulus of the velocity of the particle, and the modulus of its total acceleration will be (A) v= 0.4 m/s, a = 0.4 m/s² (C) v = 0.32 m/s, a = 0.4 m/s² (D) v = 0.4 m/s, a = 0.32 m/s²

51. A spot light S rotates in a horizontal plane with a constant angular velocity of 0.1 rad/s. The spot of light P moves along the wall at a distance 3m. What is the velocity of the spot P when θ =45°?





50.

Exercise # 2 Part # I [Multiple Correct Choice Type Questions]

1. A point moves in a straight line under the retardation av^2 . If the initial velocity is u, the distance covered in 't' seconds is :-

(B) $\frac{1}{a} \bullet n(aut)$ **(C)** $\frac{1}{a} \bullet n(1+aut)$

(**D**) a●n(aut)

2. A particle is moving in a plane with velocity given by $\stackrel{r}{u} = u_0 \hat{i} + (a\omega \cos \omega t)\hat{j}$, where \hat{i} and \hat{j} are unit vectors along x and y axes respectively. If particle is at the origin at t = 0. Calculate the trajectory of the particle:-

(A)
$$y = a \sin\left(\frac{u_0}{\omega x}\right)$$
 (B) $y = a \sin\left(\frac{\omega x}{u_0}\right)$ (C) $y = \frac{1}{a} \cdot \sin\left(\frac{u_0}{\omega x}\right)$ (D) $y = \frac{1}{a} \cdot \sin\left(\frac{\omega x}{u_0}\right)$

- 3. The relation between time t and distance x is $t=\alpha x^2+\beta x$ where α and β are constants. The retardation is :-(A) $2\alpha v^3$ (B) $2\beta v^2$ (C) $2\alpha\beta v^2$ (D) $2\beta^2 v^3$
- 4. A, B & C are three objects each moving with constant velocity. A's speed is 10 m/s in a direction PQ. The velocity of B relative to A is 6 m/s at an angle of, cos⁻¹ (15/24) to PQ. The velocity of C relative to B is 12 m/s in a direction QP. Then the magnitude of the velocity of C is :-
- (A) 5 m/s (B) $2\sqrt{10}$ m/s (C) 3 m/s (D) 4 m/s 5. A particle is moving with uniform acceleration along a straight line. Its velocities at A & B are respectively 7 m/s & 17 m/s. M is mid point of AB. If t₁ is the time taken to go from A to M and t₂ the time taken to go from

```
M to B, the ratio \frac{t_1}{t_2} is equal to :-
```

(A) 3:2
(B) 3:1
(C) 2:1
(D) 2:3
6. Two trains, which are moving along different tracks in opposite directions, are put on the same track due to a mistake. Their drivers, on noticing the mistake, start slowing down the trains when the trains are 300 m apart. Given graphs show their velocities as function of time as the trains slow down. The separation between the trains when both have stopped,, is :-



7. In the diagram shown, the displacement of particles is given as a function of time. The particle A is moving under constant velocity of 9 m/s. The particle B is moving under variable acceleration. From time t = 0 s. to t = 6 s., the average velocity of the particle B will be equal to :-



8. A person drops a stone from a building of height 20 m. At the same instant the front end of a truck passes below the building moving with constant acceleration of 1 m/s² and velocity of 2 m/s at that instant. Length of the truck if the stone just misses to hit its rear part is :
(A) 6 m
(B) 4 m
(C) 5 m
(D) 2 m



The position vector of a particle is given as $\mathbf{\hat{r}} = (t^2 - 4t + 6)\hat{\mathbf{i}} + (t^2)\hat{\mathbf{j}}$. The time after which the velocity vector 9. and acceleration vector becomes perpendicular to each other is equal to :-(D) Not possible (A) 1 sec **(B)** 2 sec (C) 1.5 sec

10. A particle moves with uniform acceleration and v_1 , v_2 and v_3 denote the average velocities in the three successive intervals of timet, t, and t, Which of the following relations is correct? (A) $(v_1 - v_2) : (v_2 - v_3) = (t_1 - t_2) : (t_2 + t_3)$ (C) $(v_1 - v_2) : (v_2 - v_3) = (t_1 - t_2) : (t_1 - t_3)$ **(B)** $(v_1 - v_2) : (v_2 - v_3) = (t_1 + t_2) : (t_2 + t_3)$ **(D)** $(v_1 - v_2) : (v_2 - v_3) = (t_1 - t_2) : (t_2 - t_3)$

A 2m wide truck is moving with a uniform speed of 8 m/s along a straight horizontal road. A pedestrian starts 11. crossing the road at an instant when the truck is 4 m away from him. The minimum constant velocity with which

he should run to avoid an accident is :-

(A)
$$1.6\sqrt{5}$$
 m/s (B) $1.2\sqrt{5}$ m/s (C) $1.2\sqrt{7}$ m/s (D) $1.6\sqrt{7}$ m/s

12. If some function say x varies linearly with time and we want to find its average value in a given time interval we

can directly find it by $\frac{x_i + x_f}{2}$. Here, x_i is the initial value of x and x_f its final value.

x and y co-ordinates of a particle moving in x-y plane at some instant are : $x = 2t^2$ and $y = 3/2t^2$. The average velocity of particle in at time interval from t = 1 s to t = 2s is :-

(B) $(12\hat{i} + 9\hat{j}) \text{ m/s}$ (A) $(8\hat{i} + 5\hat{j})$ m/s (C) $(\hat{6i} + 4.5\hat{j})$ m/s (**D**) $(10\hat{i} + 6\hat{j})$ m/s

- A particle moves along a straight line OX. At a time t (in seconds) the distance x (in metres) of the particle from O is 13. given by $x = 40 + 12t - t^3$. How long would the particle travel before coming to rest? (A) 24 m **(B)** 40 m (C) 56 m **(D)** 16 m
- 14. A particle starts from rest with constant acceleration. The ratio of space-average velocity to the time average velocity is :-
- (A) 1/2 **(B)** 3/4 (C) 4/3 (D) 3/2A particle is thrown upwards from ground. It experiences a constant resistance force which can produce 15. retardation 2 m/s². The ratio of time of ascent to the time of descent is :- $[g = 10 \text{ m/s}^2]$

16.

(B) $\sqrt{\frac{2}{3}}$ A ball is dropped from the top of a building. The ball takes 0.5 s to fall the 3m length of a window some distance from the top of the building. If the velocities of the ball at the top and at the bottom of the window are v_T and $v_{\rm B}$ respectively, then (take g = 10 m/s²) :-

(C) $\frac{2}{3}$

(D) $\sqrt{\frac{3}{2}}$

(A)
$$v_T + v_B = 12 \text{ ms}^{-1}$$
 (B) $v_B - v_T = 4.9 \text{ ms}^{-1}$ (C) $v_B v_T = 1 \text{ ms}^{-1}$ (D) $\frac{v_B}{v_T} = 1 \text{ ms}^{-1}$

- 17. Drops of water fall from the roof of a building 9m. high at regular intervals of time, the first drop reaching the ground at the same instant fourth drop starts to fall. What are the distances of the second and third drops from the roof?
- (A) 6 m and 2 m **(B)** 6 m and 3 m (C) 4 m and 1 m (**D**) 4 m and 2 m 18. Two boats A and B are moving along perpendicular paths in a still lake at night. Boat A move with a speed of 3 m/ s and boat B moves with a speed of 4 m/s in the direction such that they collide after sometime. At t = 0, the boats are 300 m apart. The ratio of distance travelled by boat A to the distance travelled by boat B at the instant of collision is:-(A) 1 **(B)** 1/2 (C) 3/4 **(D)** 4/3



Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91 - 9350679141

19. A disc in which several grooves are cut along the chord drawn from a point 'A', is arranged in a vertical plane, several particles starts slipping from 'A' along the grooves simultaneously. Assuming friction and resistance negligible, the time taken in reaching the edge of disc will be :-



(A) Maximum in groove AB(C) Same in all groove

(B) Maximum in groove AD(D) According to the heights of B, C, D, E, F

20.

A particle P is projected from a point on the surface of smooth inclined plane (see figure). Simultaneously another particle Q is released on the smooth inclined plane from the same position. P and Q collide after t = 4 second. The speed of projection of P is :-

(A) 5 m/s (B) 10 m/s (C) 15 m/s



21. A trolley is moving horizontally with a constant velocity of v m/s w.r.t. earth. A man starts running from one end of the trolley with a velocity 1.5v m/s w.r.t. to trolley. After reaching the opposite end, the man return back and continues running with a velocity of 1.5 v m/s w.r.t. the trolley in the backward direction. If the length of the trolley is L then the displacement of the man with respect to earth during the process will be :-

(A) 2.5L (B) 1.5L (C) $\frac{5L}{3}$ (D) $\frac{4L}{3}$

- 22. A body is thrown horizontally with a velocity $\sqrt{2}$ gh from the top of a tower of height h. It strikes the level ground through the foot of the tower at a distance x from the tower. The value of x is :-(A) h (B) h/2 (C) 2h (D) 2h/3
- 23. A particle is projected from a point P(2,0,0)m with a velocity 10m/s making an angle 45° with the horizontal. The plane of projectile motion passes through a horizontal line PQ which makes an angle of 37° with positive x-axis, xy plane is horizontal. The coordinates of the point where the particle will strike the line PQ is :- (take g = 10 m/s²)
 (A) (10,6,0)m
 (B) (8,6,0)m
 (C) (10,8,0)m
 (D) (6,10,0)m

24. A ball is projected from a certain point on the surface of a planet at a certain angle with the horizontal surface. The horizontal and vertical displacements x and y vary with time t in second as : $x = 10\sqrt{3}$ t; $y = 10t - t^2$ the maximum height attained by the ball is :-(A) 100m (B) 75m (C) 50 m (D) 25m

- 25. A particle A is projected with speed v_A from a point making an angle 60° with the horizontal. At the same instant, a second particle B is thrown vertically upward from a point directly below the maximum height point of parabolic path of A with velocity v_B . If the two particles collide then the ratio of v_A/v_B should be :-
 - (A) 1 (B) $\frac{2}{\sqrt{3}}$ (C) $\frac{\sqrt{3}}{2}$ (D) $\sqrt{3}$



Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141

A particle moves in the xy plane and at time t is at the point $(t^2, t^3 - 2t)$. Then :-26. (A) At t = 2/3 s, directions of velocity and acceleration are perpendicular **(B)** At t = 0, directions of velocity and acceleration are perpendicular (C) At $t = \sqrt{\frac{2}{3}}$ s, particle is moving parallel to x-axis (D) Acceleration of the particle when it is at point (4, 4) is $2\hat{i} + 24\hat{j}$ An object may have :-27. (A) Varying speed without having varying acceleration (B) Varying velocity without having varying speed (C) Non-zero acceleration without having varying velocity (D) Non-zero acceleration without having varying speed. 28. The figure shows the velocity time graph of a particle which moves along a straight line starting with velocity at 5 m/sec and coming to rest at t = 30s. Then :-(15.10)(20,10) (A) Distance travelled by the particle is 212.5 m (0.5)(B) Distance covered by the particle when it moves with constant velocity is 100 m (C) Velocity of the particle at t = 25s is 5 m/sec (30,0) (D) Velocity of the particle at t = 9s is 8 m/sec. 29. A particle moves along x-axis according to the law $x = (t^3-3t^2-9t+5)m$. Then :-(A) In the interval $3 \le t \le 5$, the particle is moving in +x direction (B) The particle reverses its direction of motion twice in entire motion if it starts at t=0(C) The average acceleration from $1 \le t \le 2$ seconds is $6m/s^2$. (D) In the interval $5 \le t \le 6$ seconds, the distance travelled is equal to the displacement. **30**. A particle moves with constant speed y along a regular hexagon ABCDEF in the same order. Then the magnitude of the average velocity for its motion from A to :-(B) D is $\frac{v}{3}$ (C) C is $\frac{v\sqrt{3}}{2}$ (A) F is $\frac{v}{5}$ **(D)** B is v. 31. The co-ordinate of the particle in x-y plane are given as $x = 2 + 2t + 4t^2$ and $y = 4t + 8t^2$:-The motion of the particle is :-(A) Along a straight line (B) Uniformly accelerated (D) Non-uniformly accelerated (C) Along a parabolic path A particle moving along a straight line with uniform acceleration has velocities 7m/s at A and 17m/s at C. B is the 32. mid point of AC. Then :-(A) The velocity at B is 12m/s (B) The average velocity between A and B is 10m/s (C) The ratio of the time to go from A to B to that from B to C is 3 : 2 (D) The average velocity between B and C is 15m/s 33. A particle moves along the X-axis as $x = u(t-2s) + a(t-2s)^2$:-(A) The initial velocity of the particle is u (B) The acceleration of the particle is a (C) The acceleration of the particle is 2a (D) At t = 2s particle is at the origin.

- 34. Pick the correct statements :-
 - (A) Average speed of a particle in a given time is never less than the magnitude of the average velocity.
 - (B) It is possible to have a situation in which $\left|\frac{d\vec{u}}{dt}\right| \neq 0$ but $\frac{d}{dt} |\vec{u}| = 0$.
 - (C) The average velocity of a particle is zero in a time interval. It is possible that the instantaneous velocity is never zero in the interval.
 - (D) The average velocity of a particle moving on a straight line is zero in a time interval. It is possible that the instantaneous velocity is never zero in the interval. (infinite acceleration is not allowed)
- A particle leaves the origin with an initial velocity $\overset{r}{u} = 3\hat{i}$ m/s and a constant acceleration 35. $\stackrel{r}{a} = (-1.0\hat{i} - 0.5\hat{j}) \text{ m/s}^2$. Its velocity $\stackrel{r}{v}$ and position vector $\stackrel{r}{r}$ when it reaches its maximum x-co-ordinate are ·-
 - **(B)** $\hat{v} = -1.5\hat{j}$ m/s (A) $\hat{v} = -2\hat{i}$ (**D**) $r = (3\hat{i} - 2\hat{j}) m$ (C) $r = (4.5\hat{i} - 2.25\hat{i}) m$
- If velocity of the particle is given by $v = \sqrt{x}$, where x denotes the position of the particle and initially particle **36**. was at x = 4m, then which of the following are correct.
 - (A) At t = 2 s, the position of the particle is at x = 9m
 - **(B)** Particle acceleration at t = 2 s. is 1 m/s^2
 - (C) Particle acceleration is $1/2 \text{ m/s}^2$ through out the motion
 - (D) Particle will never go in negative direction from it's starting position
- 37. Which of the following statements are true for a moving body?
 - (A) If its speed changes, its velocity must change and it must have some acceleration
 - (B) If its velocity changes, its speeds must change and it must have some acceleration
 - (C) If its velocity changes, its speed may or may not change, and it must have some acceleration
 - (D) If its speed changes but direction of motion does not change, its velocity may remain constant
- 38. The figure shows the v-t graph of a particle moving in straight line. Find the time when particle returns to the starting point.



(A) 30 sec

(C) 36.2 sec

(D) 35.4 sec

The velocity - time graph of the particle moving along a straight line is shown. The rate of acceleration and 39. deceleration is constant and it is equal to 5 ms^{-2} . If the average velocity during the motion is 20 ms⁻¹, then the value of t is





Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91 - 9350679141

- A particle is projected from a point P with a velocity v at an angle θ with horizontal. At a certain point Q it moves **40**. at right angle to its initial direction. Then :-(A) Velocity of particle at Q is $vsin\theta$ **(B)** Velocity of particle at Q is $vcot\theta$ (C) Time of flight from P to Q is (v/g)cosec θ (D) Time of flight from P to Q is $(v/g)sec\theta$ 41. In a projectile motion assuming no air drag let $t_{0A} = t_1$ and $t_{AB} = t_2$. The horizontal displacement from O to A is R, and from A to B is R₂. Maximum height is H and time of flight is T. Now if air drag is to be considered, then choose the correct alternative(s). (A) t, will decrease while t, will increase (B) H will increase (C) R_1 will decrease while R_2 will increase (D) T may increase or decrease A gun is set up in such a way that the muzzle is at ground level as in figure. The hoop A is located at a horizontal **42**. distance 40m from the muzzle and is 50m above the ground level. Shell is fired with initial horizontal component of velocity as 40m/s. Which of the following is/are correct?
 - (A) The vertical component of velocity of the shell just after it is fired is 55m/s,
 - if the shell has to pass through the hoop A.
 - (B) The shell will pass through both the hoops if x=40m.
 - (C) The shell will pass through both the hoops if x=20m.
 - (D) The vertical component of velocity of the shell just after it is fired is 45m/s, if the shell is to pass through both the hoops.
- **43**. If T is the total time of flight, H the maximum height and R is the horizontal range of a projectile. Then x and y co-ordinates at any time t are related as :-

(A)
$$y = 4H\left(\frac{t}{T}\right)\left(1-\frac{t}{T}\right)$$

(B) $y = 4H\left(\frac{T}{t}\right)\left(1-\frac{T}{t}\right)$
(C) $y = 4H\left(\frac{x}{R}\right)\left(1-\frac{x}{R}\right)$
(D) $y = 4H\left(\frac{R}{x}\right)\left(1-\frac{R}{x}\right)$

- 44. Two particles P & Q are projected simultaneously from a point O on a level ground in the same vertical plane with the same speed in directions making angle of 30^o and 60^o respectively with the horizontal.
 - (A) Both reach the ground simultaneously
 - (B) P reaches the ground earlier than Q
 - (C) Both strike the same point on the level ground
 - (D) The maximum height attained by Q is thrice that attained by P
- 45. Two particles A & B projected along different directions from the same point P on the ground with the same velocity of 70 m/s in the same vertical plane. They hit the ground at the same point Q such that PQ = 480m. Then :- (g = 9.8m/s²]
 - (A) Ratio of their times of flight is 4:5
 - (B) Ratio of their maximum heights is 9 : 16
 - (C) Ratio of their minimum speeds during flights is 4 : 3
 - (D) The bisector of the angle between their directions of projection makes 45^o with horizontal





46. A ball is projected on smooth inclined plane in direction perpendicular to line of greatest slope with velocity of 8m/s. Find it's speed after 1 sec.



(D) 20 m/s

- 47. A particle of mass m moves along a curve $y = x^2$. When particle has x co-ordinate as 1/2 and x-component of velocity as 4m/s. Then :-
 - (A) The position coordinate of particle are (1/2, 1/4)

(A) 10 m/s

(B) The velocity of particle will be along the line 4x - 4y - 1 = 0

(B) 12 m/s

- (C) The magnitude of velocity at that instant is $4\sqrt{2}$ m/s
- (D) The magnitude of angular momentum of particle about origin at that position is 0.
- 48. Balls are thrown vertically upward in such a way that the next ball is thrown when the previous one is at the maximum height. If the maximum height is 5m, the number of balls thrown per minute will be :(A) 40
 (B) 50
 (C) 60
 (D) 120
- 49. The horizontal range of a projectile is R and the maximum height attained by it is H. A strong wind now begins to blow in the direction of motion of the projectile, giving it a constant horizontal acceleration = g/2. Under the same conditions of projection. Find the horizontal range of the projectile.
 (A) R +H
 (B) R + 2H
 (C) R
 (D) R +H/2
- 50. In the figure shown the acceleration of A is, ${}_{A}^{I} = 15\hat{i} + 15\hat{j}$ then the acceleration of B is (A remains in contact with B)



51. Acceleration versus velocity graph of a particle moving in a straight line starting from a

rest is as shown in figure. The corresponding velocity – time graph would be:-



52.

If block A is moving with an acceleration of 5 m/s², the acceleration of B w.r.t. ground is





Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141

53. Block B has a downward velocity in m/s and given by $v_B = \frac{t^2}{2} + \frac{t^3}{6}$, where t is in s. Acceleration of A at t = 2 second is



54. Block A and C start from rest and move to the right with acceleration $a_A = 12t \text{ m/s}^2$ and $a_C = 3 \text{ m/s}^2$. Here t is in seconds. The time when block B again comes to rest is



55. In the figure acceleration of A is 1 m/s² upwards, acceleration of B is 7 m/s² upwards and acceleration of C is 2m/s² upwards. Then acceleration of D will be



56. A particle moves along an arc of a circle of radius R. Its velocity depends on the distance covered as $v=a\sqrt{s}$, where a is a constant then the angle α between the vector of the total acceleration and the vector of velocity as a function of s will be

(A) $\tan \alpha = \frac{R}{2s}$ (B) $\tan \alpha = \frac{2s}{R}$ (C) $\tan \alpha = \frac{2R}{s}$ (D) $\tan \alpha = \frac{s}{2R}$

A particle moves with deceleration along the circle of radius R so that at any moment of time its tangential and normal accelerations are equal in moduli. At the initial moment t = 0 the speed of the particle equals v_0 , then the speed of the particle as a function of the distance covered S will be

(A) $v = v_0 e^{-S/R}$ (B) $v = v_0 e^{S/R}$ (C) $v = v_0 e^{-R/S}$ (D) $v = v_0 e^{R/S}$



57.

Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141

	Part # II	\geq	>	[Assertion & Reason Type Questions]
	These questio (A) Statement (B) Statement (C) Statement	ns co 1 is 1 is 1 is	ntains, Stat True, Stater True, Stater True, Stater	ement 1 (assertion) and Statement 2 (reason). ment–2 is True ; Statement–2 is a correct explanation for Statement–1 ment–2 is True ; Statement–2 is not a correct explanation for Statement–1 ment–2 is False.
	(D) Statement	-1 is	False, State	ment-2 is True.
	(E) Statement	- I 1S İ	alse, Staten	hent-II is false.
1.	Statement–I Statement–II	:	Path of a p Trajectory be neglector	rojectile is a parabola irrespective of its velocity of projection. of a projectile is a parabola when variation of g, (acceleration due to gravity) can ed.
2.	Statement-I	:	A projectile then $a = b$	e is thrown with an initial velocity of $(a\hat{i} + b\hat{j})$ m/s. If range of projectile is maximum
	Statement-II	:	In projecti	le motion, angle of projection is equal to 45° for maximum range condition.
3.	Statement I Statement II	:	When velo Acceleratio	ocity of a particle is zero then acceleration of particle is zero. on is equal to rate of change of velocity.
4.	Statement-I	:	If two part along the	icles, moving with constant velocities are to meet, the relative velocity must be line joining the two particles.
	Statement-II	:	Relative v	elocity means motion of one particle as viewed from the other.
5.	Statement I	:	A particle particle ca	moves in a straight line with constant acceleration. The average velocity of this nnot be zero in any time interval.
	Statement II	:	For a parti	cle moving in straight line with constant acceleration, the average velocity in a
			time interv given time	ral is $\frac{u+v}{2}$, where u and v are initial and final velocity of the particle of the interval.
6 .	Statement-I	:	When a bo at the sam	dy is dropped or thrown horizontally from the same height, it reaches the ground e time.
	Statement-II	:	They have	same acceleration and same initial speed in vertical direction.
7.	Statement-I	:	A man can time to cro	cross river of width d in minimum time t. On increasing river velocity, minimum oss the river by man will remain unchanged.
	Statement-II	:	Velocity of of velocity	Friver is perpendicular to width of river. So time to cross the river is independent of river.
8.	Statement-I	:	Two balls increases l hits groun	are dropped one after the other from a tall tower. The distance between them inearly with time (elapsed after the second ball is dropped and before the first d).
	Statement-II	:	Relative ad	eccleration is zero, whereas relative velocity is non-zero in the above situation.
9.	Statement-I	:	The maxin when throw	num range along the inclined plane, when thrown downward is greater than that wn upward along the same inclined plane with same velocity.
	Statement-II	:	The maxin	num range along inclined plane is independent of angle of inclination.



10.	Statement-I	: A positive acceleration can be associated with a 'slowing down' of the body.
	Statement-II	: The origin and the positive direction of an axis are a matter of choice.
11.	Statement-I	: Two particles of different mass, are projected with same velocity at same angles. The maximum height attained by both the particles will be same.
	Statement-II	: The maximum height of projectile is independent of particle mass
12.	Statement-I	: The speed of a body can be negative.
	Statement-II	: If the body is moving in the opposite direction of positive motion, then its speed is negative.
13.	Statement-I	: Graph (1) represent one dimensional motion of a particle. While graph (2) can not represent 1– D motion of the particle. (here x is position and t is time)
		$ \begin{array}{c} \uparrow x \\ \uparrow \\ (1) \end{array} \\ (1) \end{array} \\ (2) \end{array} $
	Statement-II	: Particle can have only one position at an instant.
14.	Statement-I	: In a free fall, the initial velocity of a body may or may not be zero.
	Statement-II	: A heavy body falls at a faster rate as compared to a lighter body.
15.	Statement-I	: When the direction of motion of a particle moving in a circular path is reversed the direction of radial acceleration still remains the same (at the given point).
	Statement-II	Particle revolves on circular path in any direction such as clockwise or anticlockwise the direction of radial acceleration is always towards the centre of the circular path.
16.	Statement-I	: When a body is dropped or thrown horizontally from the same height, it would reach the ground at the same time.
	Statement-II	: Horizontal velocity has no effect on the vertical direction.
17.	Statement-I	: A cyclist must adopt a zig-zag path while ascending a steep hill.
	Statement-II	: The zig-zag path prevent the cyclist to slip down.
18.	Statement-I	: When a particle is thrown obliquely from the surface of the Earth, it always moves in a parabolic path, provided the air resistance is negligible.
	Statement-II	: A projectile motion is a two dimensional motion.
19.	Statement-I	: Mountain roads rarely go straight up the slope.
	Statement-11	. Stope of mountains are large increase more chances of venicle to shp from foads.



Exercise # 3

[Matrix Match Type Questions]

Following question contains statements given in two columns, which have to be matched. The statements in Column-I are labelled as A, B, C and D while the statements in Column-II are labelled as p, q, r and s. Any given statement in Column-I can have correct matching with one or more statement(S) in Column-II.

1. Column I

- (A) Zero acceleration
- (B) Infinite acceleration
- (C) Constant positive acceleration, with zero initial velocity

Part # I

- (D) Constant positive acceleration, with non-zero initial velocity
- (E) Constant Negative acceleration
- (F) Increasing acceleration
- (G) Decreasing acceleration



2. The equation of one dimensional motion of the particle is described in column I. At t= 0, particle is at origin and at rest. Match the column I with the statements in Column II.

	Column I		Column II
(A)	$\mathbf{x} = (3t^2 + 2)\mathbf{m}$	(P)	Velocity of particle at $t = 1s$ is 8 m/s
(B)	v = 8t m/s	(Q)	Particle moves with uniform acceleration
(C)	a = 16 t	(R)	Particle moves with variable acceleration
(D)	$v = 6t - 3t^2$	(S)	Particle will change its direction some time.

3. For the velocity-time graph shown in figure, in a time interval from t = 0 to t = 6 s, match the following:



	Column I		Column II
(A)	Change in velocity	(P)	– 5/3 SI unit
(B)	Average acceleration	(Q)	– 20 SI unit
(C)	Total displacement	(R)	- 10 SI unit
(D)	Acceleration at t=3s	(S)	– 5 SI unit

4. A balloon rises up with constant net acceleration of 10m/s^2 . After 2 s a particle drops from the balloon. After further 2 s match the following : (Take g = 10 m/s^2)

	Column I		Column II
(A)	Height of particle from ground	(P)	Zero
(B)	Speed of particle	(Q)	10 SI units
(C)	Displacement of Particle	(R)	40 SI units
(D)	Acceleration of particle	(S)	20 SI units


PHYSICS FOR JEE MAIN & ADVANCED

5. A particle is rotating in a circle of radius 1m with constant speed 4 m/s. In time 1 s, match the following (in SI units)

	Column I		Column II		
(A)	Displacement	(P)	8 sin 2		
(B)	Distance	(Q)	4		
(C)	Average velocity	(R)	2 sin 2		
(D)	Average acceleration	(S)	4 sin 2		

6. In the figure shown, acceleration of 1 is x (upwards). Acceleration of pulley P_3 , w.r.t. pulley P_2 is y (downwards) and acceleration of 4 w.r.t. to pulley P_3 is z (upwards). Taking upward +ve and downward -ve then



A car is moving on a straight road. The velocity of the car varies with time as shown in the figure. Initially (at t = 0), the car was at x = 0, where, x is the position of the car at any time 't'.



1. The variation of acceleration (A) with time (T) will be best represented by :-





KINEMATICS



(B) t = 30s to t = 50s(A) t = 50s to t = 70s(D) t = 10s to t = 20s(C) t = 30s to t = 60s

Comprehension #2

Distance is a scalar quantity. Displacement is a vector quantity. The magnitude of displacement is always less than or equal to distance. For a moving body displacement can be zero but distance cannot be zero. Same concept is applicable regarding velocity and speed. Acceleration is the rate of change of velocity. If acceleration is constant, then equations of kinematics are applicable for one dimensional motion under the gravity in which air resistance is considered, then the value of acceleration depends on the density of medium. Each motion is measured with respect of frame of reference. Relative velocity may be greater/smaller to the individual velocities.

1. A particle moves from A to B. Then the ratio of distance to displacement is :-





Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91 - 9350679141

Comprehension #3



Comprehension #4

A circus wishes to develop a new clown act. Fig. (1) shows a diagram of the proposed setup. A clown will be shot out of a cannon with velocity v_0 at a trajectory that makes an angle $\theta = 45^\circ$ with the ground. At this angle, the clown will travel a maximum horizontal distance. The cannon will accelerate the clown by applying a constant force of 10,000N over a very short time of 0.24s. The height above the ground at which the clown begins his trajectory is 10m.

(C) 4m



A large hoop is to be suspended from the ceiling by a massless cable at just the right place so that the clown will be able to dive through it when he reaches a maximum height above the ground. After passing through the hoop he will then continue on his trajectory until arriving at the safety net. Fig. (2) shows a graph of the vertical component of the clown's velocity as a function of time between the cannon and the hoop. Since the velocity depends on the mass of the particular clown performing the act, the graph shows data for several different masses.

(B) 1m





Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91 - 9350679141

1.	If the angle the cannon makes with the horizontal is increased from 45°, the hoop will have to be :- (A) Moved farther away from the cannon and lowered (B) Moved farther away from the cannon and raised (C) Moved closer to the cannon and lowered (D) Moved closer to the cannon and raised			
2.	If the clown's mass is 80 k	g, what initial velocity v_0 wi	II he have as he leaves the c	annon?
	(A) 3 m/s	(B) 15 m/s	(C) 30 m/s	(D) 300 m/s
3.	The slope of the line segment does this slope represent $(A) - g$	ents plotted in figure 2 is a (B) v ₀	constant. Which one of the f (C) $y - y_0$	following physical quantities (D) sinθ
4.	From figure 2, approximate 10 m below the height of t	ly how much time will it take f the cannon ?	for clown with a mass of 60 kg	to reach the safety net located
	(A) 4.3s	(B) 6.4s	(C) 5.9s	(D) 7.2s
5.	If the mass of a clown dou	bles, his initial kinetic energy	$v_{\rm r}$, mv_2/2, will :-	
	(A) Remain the same	(B) Be reduce to half	(C) Double	(D) Four times
6 .	If a clown holds on to hoop instead of passing through it, what is the position of the cable so that he doesn't hit his head on the ceiling as he swings upward ?			
	$(\mathbf{A}) \; \frac{2 \mathbf{v}_0^2}{\mathbf{g}}$	(B) $\frac{V_0^2}{g}$	(C) $\frac{v_0^2}{2g}$	(D) $\frac{\mathbf{v}_0^2}{4\mathrm{g}}$

Comprehension #5

A student performs an experiment to determine how the range of a ball depends on the velocity with which it is projected. The "range" is the distance between the points where the ball lands and from where it was projected, assuming it lands at the same height from which it was projected. It each trial, the student uses the same baseball, and launches it at the same angle. Table shows the experimental results.

Trail	Launch speed (m/s)	Range (m)
1	10	8
2	20	31.8
3	30	70.7
4	40	122.5

Based on this data, the student then hypothesizes that the range, R, depends on the initial speed v_o, according to the following equation : $R=Cv_0^n$, where C is a constant and n is another constant.

- Based on this data, the best guess for the value of n is :-1.
- (A) $\frac{1}{2}$ **(B)** 1 **(C)**2 **(D)** 3 2. The student speculates that the constant C depends on :-(i) The angle at which the ball was launched (ii) The ball's mass (iii) The ball's diameter If we neglect air resistance, then C actually depends on :-(A) I only (B) I and II (C) I and III (D) I, II and III
- The student performs another trial in which the ball is launched at speed 5.0 m/s. Its range is approximately: 3. **(B)** 2.0 m (A) 1.0 m (C) 3.0 m (D) $4.0 \, \text{m}$



Comprehension #6

A projectile is projected with some initial velocity and some initial angle of projection. A wind is also blowing, due to which constant horizontal retardation a is imparted to the particle in the plane of motion. It is found that, the particle is at same height at two different time $t_1 & t_2$ and particle is at same horizontal distance at two different time $t_3 & t_4$.

1. Angle of projection of particle is :-

(A)
$$\tan^{-1}\left(\frac{t_1 + t_2}{t_3 + t_4}\right)$$
 (B) $\tan^{-1}\left[\frac{a(t_1 + t_2)}{g(t_3 + t_4)}\right]$ (C) $\tan^{-1}\left[\frac{g(t_1 + t_2)}{a(t_3 + t_4)}\right]$ (D) None of these

2. Maximum height of the particle is :-

(A)
$$\frac{g}{8}(t_1 + t_2)^2$$
 (B) $\frac{g}{4}(t_1 + t_2)^2$ (C) $\frac{g}{2}(t_1 + t_2)^2$ (D) None of these

3. Range of the projectile is :-

(A)
$$\frac{1}{2}g(t_1 + t_2)^2$$
 (B) $\frac{1}{2}g(t_1 + t_2)(t_3 + t_4 - t_1 - t_2)$ (C) $\frac{1}{2}g(t_3 + t_4 - t_1 - t_2)^2$ (D) can't determined

Comprehension #7

When a particle is undergoing motion, the displacement of the particle has a magnitude that is equal to or smaller than the total distance travelled by the particle. In many cases the displacement of the particle may actually be zero, while the distance travelled by it is non-zero. Both these quantities, however depend on the frame of reference in which motion of the particle is being observed. Consider a particle which is projected in the earth's gravitational

field, close to its surface, with a speed of $100\sqrt{2}$ m/s, at an angle of 45° with the horizontal in the eastward direction. Ignore air resistance and assume that the acceleration due to gravity is 10 m/s².

- 1. The motion of the particle is observed in two different frames : one in the ground frame (A) and another frame (B), in which the horizontal component of the displacement is always zero. Two observers located in these frames will agree on :-
 - (A) The total distance travelled by the particle
- (B) The horizontal range of the particle
- (C) The maximum height risen by the particle
- (D) None of the above
- 2. "A third observer (C) close to the surface of the earth reports that particle is initially travelling at a speed of $100 \sqrt{2}$ m/s making on angle of 45° with the horizontal, but its horizontal motion is northward". The third observer is moving in :-
 - (A) The south–west direction with a speed of 100 $\sqrt{2}$ m/s
 - (B) The south–east direction with a speed of $100 \sqrt{2}$ m/s
 - (C) The north–west direction with a speed of 100 $\sqrt{2}$ m/s
 - (D) The north–east direction with a speed of $100 \sqrt{2}$ m/s
- 3. There exists a frame (D) in which the distance travelled by the particle is a minimum. This minimum distance is equal to :-
 - (A) 2 km

(B) 1 km

(C) 0 km

(D) 500 m

4. Consider an observer in frame D (of the previous question), who observes a body of mass 10 kg accelerating in the upward direction at 30 m/s² (w.r.t. himself). The net force acting on this body, as observed from the ground is :(A) 400 N in the upward direction
(B) 300 N in the upward direction
(D) 500 N in the upward direction



Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141

Exercise # 4 [Subjective Type Questions]

- The position-time (x-t) graphs for two children A and B returning from their school O to their homes P and Q respectively are shown in fig. Choose correct entries in the brackets below :
 - (i) (A/B) lives closer to the school than (B/A)
 - (ii) (A/B) starts from the school earlier than (B/A)
 - (iii) (A/B) walks faster than (B/A)
 - (iv) A and B reach home at the (same / different) time
 - (v) (A/B) overtakes (B/A) on the road (once/ twice).
- 2. For shown situation in which interval is the average speed greatest ? (Given each interval is of equal duration)



- 3. A particle is moving along x axis with acceleration $a = a_0 (1 t/T)$ where a_0 and T are constants. The particle at t = 0 has zero velocity. Calculate the average velocity between t = 0 and the instant when a = 0.
- 4. A lift accelerates downwards from rest at rate of 2 m/s², starting 100 m above the ground. After 3 sec, an object falls out of the lift. Which will reach the ground first ? What is the time interval between their striking the ground ?
- 5. A body moving with uniform acceleration, covers a distance of 20 m in the 7th second and 24 m in the 9th second. How much shall it cover in 15th second ?
- 6. A driver travelling at speed 36 kmh⁻¹ sees the light turn red at the intersection. If his reaction time is 0.6s, and then the car can deaccelerate at 4ms⁻². Find the stopping distance of the car.
- 7. A train, travelling at 20 km/hr is approaching a platform. A bird is sitting on a pole on the platform. When the train is at a distance of 2 km from pole, brakes are applied which produce a uniform deceleration in it. At that instant the bird flies towards the train at 60 km/hr and after touching the nearest point on the train flies back to the pole and then flies towards the train and continues repeating itself. Calculate how much distance will the bird have flown before the train stops ?
- 8. A parachutist after bailing out falls 52 m without friction . When the parachute opens, she decelerates at 2.1 ms^{-2} & reaches the ground with a speed of 2.9 ms⁻¹.
 - (i) How long has been the parachutist in the air? (ii) At what height did the fall begin?
- 9. A ball is thrown vertically upwards with a velocity of 20 ms^{-1} from the top of a tower. The height of the tower is 25 m from the ground.
 - (i) How high will the ball rise ?
 - (ii) How long will it be before the ball hits the ground ? (Take $g = 10 \text{ ms}^{-2}$)
- 10. A particle starts motion from rest and moves along a straight line. Its acceleration-time graph is shown. Find out speed of particle at t = 2s and at t = 3s.







- 11. A balloon is going upwards with a constant velocity 15 m/s. When the balloon is at 50 m height, a stone is dropped outside from the balloon. How long will stone take to reach at the ground ? (take $g = 10 \text{ m/s}^2$)
- 12. A train moves from one station to another in two hours time. Its speed during the motion is shown in the graph. Calculate



(i) Maximum acceleration during the journey.

- (ii) Distance covered during the time interval from 0.75 hour to 1 hour.
- 13. Two trains A and B 100 m and 60 m long are moving in opposite directions on the parallel tracks. The speed of shorter train is 3 times that of the longer one. If the train take 4 seconds to cross each other then find the velocities of the trains ?
- 14. Two cars travelling towards each other on a straight road at velocity 10 m/s and 12 m/s respectively. When they are 150 metre apart, both drivers apply their brakes and each car decelerates at 2 m/s² until it stops. How far apart will they be when they have both come to a stop ?
- 15. Two motor cars start from A simultaneously & reach B after 2 hour. The first car travelled half the distance at a speed of $v_1 = 30$ km hr⁻¹ & the other half at a speed of $v_2 = 60$ km hr⁻¹. The second car covered the entire distance with a constant acceleration. At what instant of time, were the speeds of both the vehicles same? Will one of them overtake the other in route?
- 16. In a harbour, wind is blowing at the speed of 72 km/hr and the flag on the mast of a boat anchored in the harbour flutters along the N E direction. If the boat starts moving at a speed of 51 km/hr to the north, what is the direction of flag on the mast of the boat ?
- 17. 'n' number of particles are located at the vertices of a regular polygon of n sides having the edge length 'a'. They all start moving simultaneously with equal constant speed 'v' heading towards each other all the time. How long will the particles take to collide ?
- 18. A particle is projected with a speed v and an angle θ to the horizontal. After a time t, the magnitude of the instantaneous velocity is equal to the magnitude of the average velocity from 0 to t. Find t.
- 19. A man crosses a river in a boat. If he crosses the river in minimum time he takes 10 minutes with a horizontal drift 120 m. If he crosses the river taking shortest path in 12.5 minutes then find width of the river, velocity of the boat w.r.t. water and speed of flow of river.
- 20. A particle is projected horizontally as shown from the rim of a large hemispherical bowl. The displacement of the particle when it strikes the bowl the first time is R. Find the velocity of the particle at that instant and the time taken.



- 21. A projectile is thrown with speed u making angle θ with horizontal at t=0. It just crosses the two points at equal height at time t=1 s and t = 3 sec respectively. Calculate maximum height attained by it. (g=10m/s²)
- 22. A Bomber flying upward at an angle of 53° with the vertical releases a bomb at an altitude of 800 m. The bomb strikes the ground 20s after its release. Find : [Given sin $53^{\circ} = 0.8$; $g = 10 \text{ ms}^{-2}$]
 - (i) The velocity of the bomber at the time of release of the bomb.

(ii) The maximum height attained by the bomb.

- (iii) The horizontal distance travelled by the bomb before it strikes the ground
- (iv) The velocity (magnitude & direction) of the bomb just when it strikes the ground .
- 23. A food package was dropped from an aircraft flying horizontally. 6 s before it hit the ground, it was at a height of 780 m, and had travelled a distance of 1 km horizontally. Find the speed and the altitude of the aircraft.



24. Two particles are projected from the two towers simultaneously, as shown in the figure. What should be value of 'd' for their collision.



25. A body falls freely from some altitude H. At the moment the first body starts falling another body is thrown from the earth's surface which collides with the first at an altitude h = H/2. The horizontal distance is \bullet . Find the initial velocity and the angle at which it was thrown.....



26. Calculate the relative acceleration of A w.r.t. B if B is moving with acceleration a₀ towards right.



- 27. A particle is performing circular motion of radius 1 m. Its speed is $v = (2t^2)$ m/s. What will be magnitude of its acceleration at t = 1s.
- 28. A ring rotates about z axis as shown in figure. The plane of rotation is xy.

At a certain instant the acceleration of a particle P (shown in figure) on the ring is $(6\hat{i} - 8\hat{j})m/s^2$. Find the angular acceleration of the ring & the angular velocity



at that instant. Radius of the ring is 2m.

- 29. A particle is moving in a circular orbit with a constant tangential acceleration. After a certain time t has elapsed after the beginning of motion, the angle between the total acceleration a and the direction along the radius r becomes equal to 45°. What is the angular acceleration of the particle.
- 30. Two particles A and B start at O and travel in opposite directions along the circular path at constant speed $v_A = 0.7$ m/s and $v_B = 1.5$ m/s respectively. Determine the time when they collide and the magnitude of the acceleration

of B just before this happening. (radius = 5m)



31. Two particles A and B move anticlockwise with the same speed v in a circle of radius R and are $72v^2$

diametrically opposite to each other. At t=0, A is given a constant acceleration (tangential) $a_t = \frac{72v^2}{25\pi R}$.

Calculate the time in which A collides with B, the angle traced by A, its angular velocity and radial acceleration at the time of collision.



32. A particle moves clockwise in a circle of radius 1m with centre at (x,y) = (1m,0). It starts at rest at the origin at time t=0. Its speed increases at the constant rate of $\left(\frac{\pi}{2}\right)$ m/s². (i) How long does it take to travel halfway

around the circle? (ii) What is the speed at that time?

- **33**. Figure shows the total acceleration and velocity of a particle moving clockwise in a circle of radius 2.5m at a given instant of time. At this instant, Find :(i) the radial acceleration, (ii) the speed of the particle and (iii) its tangential acceleration.
- 34. The brakes of a train which is travelling at 30 m/s are applied as the train passes point A. The brakes produce a constant retardation of magnitude 3λ m/s² until the speed of the train is reduced to 10 m/s. The train travels at this speed for a distance and is then uniformly accelerated at λ m/s² until it again reaches a speed of 30 m/s as it passes point B. The time taken by the train in travelling from A to B, a distance of 4 km, is 4 min. Sketch the speed time graph for this motion and calculate :-(i) The value of λ (ii) Distance travelled at 10 m/s.
- 35. Two towns A and B are connected by a regular bus service with a bus leaving in either direction every T minutes. A man cycling with a speed of 20 km h⁻ in the direction A to B notices that a bus goes past him every 18 min. in the direction of his motion, and every 6 min. in the opposite direction. What is the period T of the bus service and with what speed (assumed constant) do the buses ply on the road ?
- 36. A balloon starts ascending from the ground at a constant speed of 25 m/s. After 5 s, a bullet is shot vertically upwards from the ground.
 - (i) What should be the minimum speed of the bullet so that it may reach the balloon ?
 - (ii) The bullet is shot at twice the speed calculated in (i). Find the height at which it passes the balloon.
- 37. A ship is moving at a constant speed of 10 km/hr in the direction of the unit vector \hat{i} . Initially, its position vector, relative to a fixed origin is $10(-\hat{i} + \hat{j})$ where $\hat{i} & \hat{j}$ are perpendicular vectors of length 1 km. Find its position vector relative to the origin at time t hours later. A second ship is moving with constant speed u km/hr parallel to the vector $\hat{i} + 2\hat{j}$ and is initially at the origin.

(i) If $u=10\sqrt{5}$ km / h. Find the minimum distance between the ships and the corresponding value of t (ii) Find the value of u for which the ships are on a collision course and determine the value of t at which the collision would occur if no avoiding action were taken.

- 38. A helicopter takes off along the vertical with an acceleration of 3 m/sec² & zero initial velocity. In a certain time, the pilot switches off the engine. At the point of take-off, the sound dies away in 30 sec. Determine the velocity of the helicopter at the moment when its engine is switched off, assuming the velocity of sound is 320 m/s.
- **39.** Two bodies move towards each other in a straight line at initial velocities $v_1 & v_2 & with constant accelerations a_1 & a_2 directed against the corresponding velocities at the initial instant. What must be the maximum initial separation <math>\bullet_{max}$ between the bodies for which they meet during the motion ?
- 40. A swimmer starts to swim from point A to cross a river. He wants to reach point B on the opposite side of the river. The line AB makes an angle 60° with the river flow as shown. The velocity of the swimmer in still water is same as that of the water



- (i) In what direction should he try to direct his velocity? Calculate angle between his velocity and river velocity.
- (ii) Find the ratio of the time taken to cross the river in this situation to the minimum time in which he can cross this river.





41. Two swimmers start a race. One who reaches the point C first on the other bank wins the race. A makes his strokes in a direction of 37° to the river flow with velocity 5km/hr relative to water. B makes his strokes in a direction 127° to the river flow with same relative velocity. River is flowing with speed of 2km/hr and is 100m wide. Who will win the race? Compute the time taken by A and B to reach the point C if the speeds of A and B on the ground are 8 km/hr and 6 km/hr respectively.



- 42. A shell is fired from a point O at an angle of 60° with a speed of 40 m/s & it strikes a horizontal plane through O, at a point A. The gun is fired a second time with the same angle of elevation but a different speed v. If it hits the target which starts to rise vertically from A with a constant speed $9\sqrt{3}$ m/s at the same instant as the shell is fired, find v.
- 43. Hailstones falling vertically with speed of 10 m/s hit the wind screen of a moving car and rebound elastically . Find the velocity of the car if the driver find the hailstones rebound vertically after striking . Wind screen makes an angle 30° with the horizontal.
- 44. A stone is projected from the point of a ground in such a direction so as to hit a bird on the top of a telegraph post of height h and then attain the maximum height 2h above the ground. If at the instant of projection, the bird were to fly away horizontally with a uniform speed, find the ratio between the horizontal velocities of the bird and the stone, if the stone still hits the bird while descending.
- 45. Two inclined planes OA and OB having inclinations 30° and 60° respectively, intersect each other at O as in figure. A particle is projected from point P with velocity $u = 10\sqrt{3} \text{ m/s}$ along a direction perpendicular to plane OA. If the particle strikes the plane OB perpendicularly at Q. Calculate (i) Time of flight.
 - (ii) Velocity with which particle strikes the plane OB.
 - (iii) Vertical height of P from O.

(iv) Maximum height from O attained by the particle.

- (v) Distance PQ.
- 46. A particle is projected with a velocity $2\sqrt{ag}$ so that it just clears two walls of equal height 'a'. which are at a distance 2a apart. Show that the time of passing between the walls is $2\sqrt{a/g}$.
- 47. A projectile is launched at an angle α from a cliff of height H above the sea level. If it falls into the sea at a distance

D from the base of the cliff, show that its maximum height above the sea level is $\left[H + \frac{D^2 \tan^2 \alpha}{4 \left(H + D \tan \alpha\right)}\right].$





E	xercise # 5	Part # I [Pre	evious Year Questions] [AIEEE/JEE-MAIN]
1.	A ball whose kinetic en the highest point of its f	ergy is E, is projected at ar light will be-	n angle of 45° to the horizont	al. The kinetic energy of the ball at [AIEEE - 2002]
	(1) E	(2) $E/\sqrt{2}$	(3) E/2	(4) zero
2.	A boy playing on the ro horizontal. How far from	of of a 10 m high building m the throwing point will t	throws a ball with a speed of he ball be at the height of 10	f 10 m/s at an angle of 30° with the m from the ground ?
	$[g = 10 \text{ m/s}^2, \sin 30^\circ = 1]$	$(2, \cos 30^\circ = \sqrt{3}/2]$		[AIEEE - 2003]
	(1) 5.20 m	(2) 4.33 m	(3) 2.60 m	(4) 8.66 m
3.	A ball is thrown from a	point with a speed v_0 at a	an angle of projection θ . Fro	om the same point and at the same
	instant, a person starts	running with a constant sp	beed $\frac{v_0}{2}$ to catch the ball. W	Vill the person be able to catch the
	ball ? If yes, what shou	ld be the angle of projection	on?	[AIEEE - 2004]
	(1) Yes, 60°	(2) Yes, 30°	(3) No	(4) Yes, 45°
4.	A projectile can have the cases, then the product	he same range R for two ar of the two times of flights	ngles of projection. If t ₁ and s is proportional to-	t ₂ be the times of flights in the two [AIEEE - 2005]
	(1) R^2	(2) $\frac{1}{R^2}$	(3) $\frac{1}{R}$	(4) R
5.	A particle is projected a	at 60° to the horizontal wit	h a kinetic energy K. The kir	netic energy at the highest point is-
				[AIEEE - 2007]
	(1) K	(2) zero	(3) K/4	(4) K/2
6.	A particle is moving wit	h velocity $\vec{v} = K(v $ + x	\$), where K is a constant.	The general equation for its path
	is:			[AIEEE - 2010]
	(1) $y^2 = x^2 + constant$		(2) $y = x^2 + constant$	
	(3) $y^2 = x + constant$		(4) $xy = constant$	
7.	A water fountain on the is v, the total area arou	e ground sprinkles water and the fountain that gets	all around it. If the speed of wet is :-	water coming out of the fountain [AIEEE - 2011]
	(1) $\frac{\pi}{2} \frac{v^4}{g^2}$	$(2) \pi \frac{v^2}{g^2}$	$(3) \pi \frac{v^2}{g}$	(4) $\pi \frac{v^4}{g^2}$
8.	A boy can throw a stone the same stone up to w	up to a maximum height of	f 10 m. The maximum horizo	ntal distance that the boy can throw [AIEEE - 2012]
	(1) 20 m	(2) $20\sqrt{2}$ m	(3) 10 m	(4) $10\sqrt{2}$ m
9.	A projectile is given an	initial velocity of $(\hat{i} + 2\hat{i})$	m/s , where \hat{i} is along the g	round and \hat{i} is along the vertical.
	If $g = 10 \text{ m/s}^2$, the equa	tion of its trajectory is :	, , <u> </u>	[JEE (Main) - 2013]
	(1) $y = x - 5x^2$	(2) $y = 2x - 5x^2$	(3) $4y = 2x - 5x^2$	(4) $4y = 2x - 25x^2$



- From a tower of height H, a particle is thrown vertically upwards with a speed U. The time taken by the particle, to hit the ground, is n times that taken by it to reach the highest point of its path. The relation between H, u and n is:
- (1) $2 g H = nu^2(n-2)$ (2) $g H = (n-2)u^2$ (3) $2 g H = n^2u^2$ (4) $g H = (n-2)^2u^2$ 11. Two stones are thrown up simultaneously from the edge of a cliff 240 m high with initial speed of 10 m/s and 40 m/ s respectively. Which of the following graph best represents the time variation of relative position of the second stone with respect to the first ?

(Assume stones do no rebound after hitting the ground and neglect air resistance, take $g = 10 \text{ m/s}^2$)





Previous Year Questions][IIT-JEE ADVANCED]

 In 1.0s, a particle goes from point A to point B, moving in a semicircle (see figure). The magnitude of the average velocity is :- [IIT-JEE 1999]
 (A) 3.14 m/s

(B) 2.0 m/s

(C) 1.0 m/s

Part # II

- (D) zero
- 2. A ball is dropped vertically from a height d above the ground. It hits the ground and bounces up vertically to a height d/2. Neglecting subsequent motion and air resistance, its velocity v varies with height h above the ground as :

 [IIT-JEE 2000]



3.

A particle starts from rest. Its acceleration (A) versus time (t) is as shown in the figure. The maximum speed of the particle will be :- [IIT-JEE 2004]





Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141 4. A small block slides without friction down an inclined plane starting from rest. Let s_n be the distance travelled



5. The given graph shows the variation of velocity with displacement. Which one of the graph given below correctly represents the variation of acceleration with displacement :- [IIT-JEE 2005]



Assertion-Reasoning Type Questions

1. **Statement-I**: For an observer looking out through the window of a fast moving train, the nearby objects appear to move in the opposite direction to the train, while the distant objects appear to be stationary.

[IIT-JEE 2008]

Statement-2: If the observer and the object are moving at velocities V_1 and V_2 respectively with reference

to a laboratory frame, the velocity of the object with respect to the observer is $\dot{V}_2 - \dot{V}_1$.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

MCQ's with one or More Than One Correct Answer

- 1. The coordinates of a particle moving in a plane are given by $x(t) = a\cos(pt)$ and $y(t) = b\sin(pt)$ where a, b (<a) and p are positive constants of appropriate dimensions. Then : [IIT-JEE 1999]
 - (A) The path of the particle is an ellipse.
 - (B) The velocity and acceleration of the particle are normal to each other at $t = \pi/2p$
 - (C) The acceleration of the particle is always directed towards a focus.

(**D**) The distance travelled by the particle in time interval t = 0 to $t = \frac{\pi}{2p}$ is a

Subjective Questions

2.

3.

1. A large heavy box is sliding without friction down a smooth plane of inclination θ . From a point P on the bottom of the box, a particle is projected inside the box. The initial speed of the particle with respect to the box is u and the direction of projection makes an angle α with the bottom as shown in the figure.

Q D D D

(i) Find the distance along the bottom of the box between the point of projection P and the point Q where the particle lands (Assume that the particle does not hit any other surface of the box. Neglect air resistance)
(ii) If the horizontal displacement of the particle as seen by an observer on the ground is zero, find the speed of the box with respect to the ground at the instant when the particle was projected. [IIT-JEE 1998]

An object A is kept fixed at the point x = 3m and y=1.25 m on a plank P raised above the ground. At time t = 0 the plank starts moving along the +x direction with an acceleration 1.5 m/s². At the same instant a stone is projected from the origin with a velocity $\frac{1}{u}$ as shown. A stationary person on the ground observes the stone hitting the object during its downward motion at an angle of 45° to the horizontal. All the motions are in x-y plane. Find $\frac{1}{u}$ and the time after which the stone hits the object. Take $g = 10 m/s^2$.



[IIT-JEE 2000]

[IIT-JEE 2002]

- On a frictionless horizontal surface, assumed to be the x-y plane, a small trolley A is moving along a straight line parallel to the y-axis (see figure) with a constant velocity of $(\sqrt{3} - 1)$ m/s. At a particular instant when the line OA makes an angle of 45° with the x-axis, a ball is thrown along the surface from the origin O. Its velocity makes on angle ϕ with the x-axis and it hits the trolly.
 - (i) The motion of the ball is observed from the frame of the trolley. Calculate the angle θ made by the velocity vector of the ball with the x-axis in this frame.
 - (ii) Find the speed of the ball with respect to the surface, if $\phi = \frac{4\theta}{3}$.

Integer Type Questions

- A train is moving along a straight line with a constant acceleration 'a'. A boy standing in the train throws a ball forward with a speed of 10 m/s, at an angle of 60° to the horizontal. The boy has to move forward by 1.15 m inside the train to catch the ball back at the initial height. The acceleration of the train, in m/s², is
- 2. Airplanes A and B are flying with constant velocity in the same vertical plane at angles 30° and 60° with respect to the horizontal respectively as shown in the figure. The speed of A is $100\sqrt{3} ms^{-1}$. At time t = 0 s, an observer in A finds B at a distance of 500 m. This observer sees B moving with a constant velocity perpendicular to the line of motion of A. If at t = t₀, A just escapes being hit by B, t₀ in seconds is



[[]IIT-JEE 2014]



Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141



3. A rocket is moving in a gravity free space with a constant acceleration of $2 \text{ ms}^{-2} \text{ along} + x$ direction (see figure). The length of a chamber inside the rocket is 4 m. A ball is thrown from the left end of the chamber in + x direction with a speed of 0.3 ms^{-1} relative to the rocket. At the same time, another ball is thrown in -x direction with a speed of 0.2 ms^{-1} from its right end relative to the rocket. The time in seconds when the two balls hit each other is



4. A horizontal circular platform of radius 0.5 m and mass 0.45 kg is free to rotate about its axis. Two massless spring toy-guns, each carrying a steel ball of mass 0.05 kg are attached to the platform at a distance 0.25 m from the centre on its either sides along its diameter (see figure). Each gun simultaneously fires the balls horizontally and perpendicular to the diameter in opposite directions. After leaving the platform, the balls have horizontal speed of 9 ms-1 with respect to the ground. The rotational speed of the platform in rad s-1 after the balls leave the platform is

[IIT-JEE 2014]

Comprehension

A frame of reference that is accelerated with respect to an inertial frame of reference is called a non-inertial frame of reference. A coordinate system fixed on a circular disc rotating about a fixed axis with a constant angular velocity ω is an example of a non-inertial frame of reference.

The relationship between the force \mathbf{F}_{rot} experienced by a particle of mass m moving on the rotating disc and the

force $\stackrel{\mathbf{i}}{F_{in}}$ experienced by the particle in an inertial fame of reference is $\stackrel{\mathbf{i}}{F_{rot}} = \stackrel{\mathbf{i}}{F_{in}} + 2m(\stackrel{\mathbf{i}}{v_{rot}} \times \stackrel{\mathbf{u}}{\omega}) + m(\stackrel{\mathbf{u}}{\omega} \times \stackrel{\mathbf{i}}{r}) \times \stackrel{\mathbf{u}}{\omega}$ where $\stackrel{\mathbf{i}}{v_{rot}}$ is the velocity of the particle in the rotating frame of reference and $\stackrel{\mathbf{i}}{r}$ is the position vector of the particle with respect to the centre of the disc.

Now consider a smooth slot along a diameter of a disc of radius R rotating counter-clockwise with a constant angular speed w about its vertical axis through its center. We assign a coordinate system with the origin at the centre of the disc, the x-axis along the slot, the y-axis perpendicular to the slot and the z-axis along the rotation axis

 $\begin{pmatrix}
\mathbf{u} \\
(\omega = \omega \hat{k})
\end{pmatrix}$. A small block of mass m is gently placed in the slot at $\stackrel{\mathbf{r}}{\mathbf{r}} = (\mathbf{R} / 2)\hat{\mathbf{i}}$ at $\mathbf{t} = 0$ and is constrained to move only along the slot. [IIT-JEE 2016]

1. The distance r of the block at time t is

(A)
$$\frac{R}{4}(e^{\omega t}+e^{-\omega t})$$
 (B) $\frac{R}{2}\cos\omega t$

(C)
$$\frac{R}{4}(e^{2\omega t} + e^{-2\omega t})$$
 (D) $\frac{R}{2}\cos 2\omega t$

2. The net reaction of the disc on the block is

(A)
$$\frac{1}{2}$$
 m $\omega^2 R(e^{2\omega t} - e^{-2\omega t})\hat{j} + mg\hat{k}$
(C) $-m\omega^2 R\cos\omega t\hat{j} - mg\hat{k}$

(B)
$$\frac{1}{2}$$
 m $\omega^2 R(e^{\omega t} - e^{-\omega t})\hat{j} + mg\hat{k}$
(D) m $\omega^2 R\cos\omega t\hat{j} - mg\hat{k}$







Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141

A particle moves along x-axis with initial position x = 0. Its velocity varies with x-coordinate as shown in 8. graph. The acceleration 'a' of this particle varies with x as :



9. Initially car A is 10.5 m ahead of car B. Both start moving at time t = 0 in the same direction along a straight line. The velocity time graph of two cars is shown in figure. The time when the car B will catch the car A, will be :

car B



10. A semicircle of radius R = 5m with diameter AD is shown in figure. Two particles 1 and 2 are at points A and B on shown diameter at t = 0 and move along segments AC and BC with constant speeds u, and u, respectively.

Then the value of $\frac{u_1}{u_2}$ for both particles to reach point C simultaneously will be :



- 11. Two bikes A and B start from a point. A moves with uniform speed 40 m/s and B starts from rest with uniform acceleration 2 m/s². If B starts at t = 0 and A starts from the same point at t = 10 s, then the time during the journey in which A was ahead of B is : **(B)** 8 s
 - (A) 20 s

(C) 10 s

(D) A was never ahead of B

12. Graphs I and II give coordinates x(t) and y(t) of a particle moving in the x-y plane Acceleration of the particle is constant and the graphs are drawn to the same scale. Which of the vector shown in options best represents the acceleration of the particle :





13. An insect moving along a straight line, (without returning) travels in every second distance equal to the magnitude of time elapsed. Assuming acceleration to be constant, and the insect starts at t = 0. Find the magnitude of initial velocity of insect. (A) $\frac{1}{2}$ unit **(B)** $\frac{1}{4}$ unit (C) $\frac{3}{2}$ unit (D) 1 unit 14. A stone is dropped from the top of building and at the same time a second stone is thrown vertically upward from the bottom of the building with a speed of 20 ms⁻¹. They pass each other 3 seconds later. Find the height of the building. **(B)** 60 m (C) 65 m (A) 40 m (D) 80 m The position vector of a particle is given as $\vec{\mathbf{k}} = (t^2 - 4t + 6)\hat{\mathbf{j}} + (t^2)\hat{\mathbf{j}}$. The time after which the velocity vector 15. and acceleration vector becomes perpendicular to each other is equal to (A) 1sec **(B)** 2 sec (C) 1.5 sec (D) not possible 16 Each of the four particles move along x axis. Their coordinates (in metres) as function of time (in seconds) are given by Particle 1 : $x(t) = 3.5 - 2.7t^3$ Particle 2: $x(t) = 3.5 + 2.7t^3$ Particle 3 : $x(t) = 3.5 + 2.7t^2$ Particle 4 : $x(t) = 3.5 - 3.4t - 2.7t^2$ which of these particles is speeding up for t > 0? (A) All four (B) only 1 (C) only 1, 2 and 3. (D) only 2, 3 and 4. **SECTION - II : MULTIPLE CORRECT ANSWER TYPE**

SECTION - II: MULTIPLE CORRECT ANSWER

17. Mark the correct statements:

- (A) Average velocity of a particle moving on a straight line was zero in a time interval but its instantaneous velocity was never zero in that interval (Consider finite acceleration only)
- (B) Average velocity of a particle was zero in a time interval but its speed was never zero in that interval (Consider finite acceleration only)
- (C) We can have a situation where magnitude of acceleration is non-zero, but rate of change of speed is zero
- (D) Average speed of particle in a given time interval is always greater than or equal to the magnitude of the average velocity in the same time interval.

18. Which of the following statements are true for a moving body?

- (A) if its speed changes, its velocity must change and it must have some acceleration.
- (B) if its velocity changes, its speed must change and it must have some acceleration.
- (C) if its velocity changes, its speed may or may not change, and it must have some acceleration.
- (D) if its speed changes but direction of motion does not change, its velocity may remain constant.
- 19. If velocity of the particle is given by $v = \sqrt{x}$, where x denotes the position of the particle and initially particle was at x = 4, then which of the following are correct?
 - (A) at t = 2 sec, the position of the particle is at x = 9.
 - **(B)** Particle's acceleration at t = 2 sec. is 1 m/s^2 .
 - (C) Particle's acceleration is $\frac{1}{2}$ m/s² throughout the motion.
 - (D) Particle will never go in negative direction from it's starting position.

20. A particle has a rectilinear motion and the figure gives its displacement as a function of time . Which of the following statements are true with respect to the motion ?



- (A) In the motion between O and A the velocity is positive and acceleration is negative.
- (B) Between A and B the velocity and acceleration are positive .
- (C) Between B and C the velocity is negative and acceleration is positive .
- (D) Between D and E the acceleration is positive .
- 21. A rabbit is moving in straight line towards a carrot, slowing down its speed so that in each second it moves half the remaining distance from his nose to a carrot. If the total distance travelled by the rabbit in time t is X, its instantaneous speed V and magnitude of its instantaneous acceleration 'a' then which of the following graph(S) is/are best representing the motion ?



- 22. A particle moves with an initial velocity v_0 and retardation βv , where v is its velocity at any time t (β is a positive constant).
 - (A) the particle will cover a total distance of v_0/β
 - (B) the particle will continue to move for a very long time
 - (C) the particle will stop shortly
 - (**D**) the velocity of particle will become $v_0/2$ after time $1/\beta$.

SECTION - III : ASSERTION AND REASON TYPE

23. Statement 1 : Magnitude of average velocity is equal to average speed.

Statement 2 : Magnitude of instantaneous velocity is equal to instantaneous speed.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1.
- (C) Statement-1 is True, Statement-2 is False.
- (D) Statement-1 is False, Statement-2 is True.

24. **Statement 1**: When velocity of a particle is zero then acceleration of particle is also zero.

Statement 2 : Acceleration is equal to rate of change of velocity.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1.
- (C) Statement-1 is True, Statement-2 is False.
- (D) Statement-1 is False, Statement-2 is True.



25. Statement-1 : A particle moves in a straight line with constant acceleration. The average velocity of this particle cannot be zero in any time interval

Statement-2: For a particle moving in straight line with constant acceleration, the average velocity in a time 11 + V

interval is $\frac{u+v}{2}$, where u and v are initial and final velocity of the particle of the given time interval.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1.
- (C) Statement-1 is True, Statement-2 is False.
- (D) Statement-1 is False, Statement-2 is True.
- 26. Statement-1 : For a particle moving in a straight line, velocity (v in m/s) of the particle in terms of time (t in sec) is given by $v = t^2 6t + 8$. Then the speed of the particle is minimum at t = 2 sec.

Statement-2: For a particle moving in a straight line the velocity v at any time t may be minimum or may be maximum when $\frac{dv}{dt} = 0$.

(A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.

- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- **(D)** Statement-1 is False, Statement-2 is True

SECTION - IV : COMPREHENSION TYPE

Comprehension - 1

A particle moves along x-axis and its acceleration at any time t is $a = 2 \sin(\pi t)$, where t is in seconds and a is in m/s². The initial velocity of particle (at time t = 0) is u = 0.

27. Then the distance travelled (in meters) by the particle from time t = 0 to t = 1 s will be :

(A)
$$\frac{2}{\pi}$$
 (B) $\frac{1}{\pi}$ (C) $\frac{4}{\pi}$ (D) None of these

- 28. Then the distance travelled (in meters) by the particle from time t = 0 to t = t will be :
 - (A) $\frac{2}{\pi^2} \sin \pi t \frac{2t}{\pi}$ (B) $-\frac{2}{\pi^2} \sin \pi t + \frac{2t}{\pi}$ (C) $\frac{2t}{\pi}$ (D) None of these
- 29. Then the magnitude of displacement (in meters) by the particle from time t = 0 to t = t will be :

(A)
$$\frac{2}{\pi^2} \sin \pi t - \frac{2t}{\pi}$$
 (B) $-\frac{2}{\pi^2} \sin \pi t + \frac{2t}{\pi}$ (C) $\frac{2t}{\pi}$ (D) None of these

Comprehension - 2

A car is moving on a straight road. The velocity of the car varies with time as shown in the figure. Initially (at t = 0), the car was at x = 0, where, x is the position of the car at any time 't'.





PHYSICS FOR JEE MAIN & ADVANCED

30. The variation of acceleration (A) with time (t) will be best represented by :



31. The displacement time graph will be best represented by :



32.The maximum displacement from the starting position will be :
(A) 200 m(B) 250 m(C) 160 m(D) 165 m

SECTION - V : MATRIX - MATCH TYPE

33. The velocity time graph for a particle moving along a straight line is given in each situation of column-I. In the time interval $\infty > t > 0$, match the graph in column-I with corresponding statements in column-II.



Column-II

(p) speed of particle is continuously decreasing.

(q) magnitude of acceleration of particle is decreasing with time.





(r) direction of acceleration of particle does not change.

(s) magnitude of acceleration of particle is increasing with time.

(t) magnitude of acceleration of particle does not change.

34. The equation of motion of the particle is described in column I. At t = 0, particle is at origin and at rest. Match the column I with the statements in column II.
 Column I

$(\mathbf{A}) \mathbf{x} = (3t^2 + 2t)\mathbf{m}$	(p) velocity of particle at $t = 1$ s is 8 m/s.
(B) $v = 8t m/s$	(q) particle moves with uniform acceleration.
(C) $a = 16 t$	(r) particle moves with variable acceleration.
(D) $v = 6t - 3t^2$	(s) acceleration of the particle at $t = 1 \sec is 2m/s$
	(t) particle will change its direction some time.

SECTION - VI : INTEGER TYPE

- 35. A railway track runs parallel to a road until a turn brings the road to railway crossing. A cyclist rides along the road everyday at a constant speed 20 km/hr. He normally meets a train that travels in same direction at the crossing. One day he was late by 25 minutes and met the train 10 km before the railway crossing. Find the speed of the train (in km/hr).
- 36. A particle starts moving rectilinearly at time t = 0 such that its velocity 'v' changes with time 't' according to the equation $v = t^2 t$ where t is in seconds and v is in m/s. Find the time interval for which the particle retards. (in m/s)
- 37. A bird is at a point P(4m, -1m, 5m) and sees two points $P_1(-1m, -1m, 0m)$ and $P_2(3m, -1m, -3m)$. At time t = 0, it starts flying in a plane of the three positions, with a constant speed of 2 m/s in a direction perpendicular to

the straight line
$$P_1P_2$$
 till it sees $P_1 \& P_2$ collinear at time t. If t is $\frac{x}{10}$ sec then x is



\rightarrow mock test : projectile motion \langle

SECTION-I: STRAIGHT OBJECTIVE TYPE

- 1. A body is thrown horizontally with a velocity $\sqrt{2gh}$ from the top of a tower of height h. It strikes the level ground through the foot of the tower at a distance x from the tower. The value of x is:
 - (A) h (B) $\frac{h}{2}$ (C) 2h (D) $\frac{2h}{3}$
- 2. It was calculated that a shell when fired from a gun with a certain velocity and at an angle of elevation $\frac{5\pi}{36}$ rad should strike a given target. In actual practice, it was found that a hill just prevented the trajectory. At what angle of elevation should the gun be fired to hit the target?

(A)
$$\frac{5\pi}{36}$$
 rad (B) $\frac{11\pi}{36}$ rad (C) $\frac{7\pi}{36}$ rad (D) $\frac{13\pi}{36}$ rad.

3. A ball is projected horizontally with a speed v from the top of a plane inclined at an angle 45° with the horizontal. How far from the point of projection will the ball strike the plane?

(A)
$$\frac{v^2}{g}$$
 (B) $\sqrt{2} \frac{v^2}{g}$ (C) $\frac{2v^2}{g}$ (D) $\sqrt{2} \left[\frac{2v^2}{g}\right]$

4. Consider a boy on a trolley who throws a ball with speed 20 m/s at an angle 37° with respect to trolley in direction of motion of trolley which moves horizontally with speed 10 m/s then what will be maximum distance travelled by ball parallel to road :

(A) 20.2 m
(B) 12 m
(C) 31.2 m
(D) 62.4 m

- 5. A particle is projected up the inclined plane such that its component of velocity along the incline is 10 m/s. Time of flight is 2 sec and maximum height above the incline is 5 m. Then velocity of projection will be :
 - (A) 10 m/s
- 6. Two men A and B, A standing on the extended floor nearby a building and B is standing on the roof of the building. Both throw a stone each towards each other. Then which of the following will be correct:

(C) $5\sqrt{5}$ m/s

(D) none

- (A) stone will hit A, but not B
- (B) stone will hit B, but not A
- (C) stone will not hit either of them, but will collide with each other

(B) $10\sqrt{2}$ m/s

- (D) none of these.
- 7. A particle P is projected from a point on the surface of smooth inclined plane (see figure). Simultaneously another particle Q is released on the smooth inclined plane from the same position. P and Q collide on the inclined plane after t = 4 second. The speed of projection of P is : (in m/s)

	Pro		
	mmmmmmmm	60°	
(A) 5 m/s	(B) 10 m/s	(C) 15 m/s	(D) 20 m/s



Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141 8. A particle is projected from a point (0, 1) on Y-axis (assume + Y direction vertically upwards) aiming towards a point (4, 9). It fell on ground along x axis in 1 sec. Taking $g = 10 \text{ m/s}^2$ and all coordinate in metres. Find the X-coordinate where it fell:

(A) (3,0)	(B) (4,0)	(C) (2,0)	(D) $(2\sqrt{5}, 0)$
------------------	------------------	------------------	-----------------------------

9. Velocity of a stone projected, 2 second before it reaches the maximum height, makes angle 53° with the horizontal then the velocity at highest point will be : (in m/s)
 (A) 20 m/s
 (B) 15 m/s
 (C) 25 m/s
 (D) 80/3 m/s

10. A stone is projected from a horizontal plane. It attains maximum height 'H' & strikes a stationary smooth wall & falls on the ground vertically below the maximum height. Assume the collision to be elastic, the height of the point on the wall where ball will strike is :



11. A particle is projected from a tower as shown in figure, then the distance from the foot of the tower where it will strike the ground will be : (take $g = 10 \text{ m/s}^2$)



- 12. Distance between a frog and an insect on a horizontal plane is 10 m. Frog can jump with a maximum speed of $\sqrt{10}$ m/s. g =10 m/s². Minimum number of jumps required by the frog to catch the insect is : (A) 5 (B) 10 (C) 100 (D) 50
- 13. A particle starts from the origin at t = 0 and moves in the x-y plane with constant acceleration 'a' in the y direction. Its equation of motion is $y = bx^2$. The x component of its velocity (at t = 0) is :

(A) variable (B)
$$\sqrt{\frac{2a}{b}}$$
 (C) $\frac{a}{2b}$ (D) $\sqrt{\frac{a}{2b}}$

14. A projectile is fired at an angle θ with the horizontal. Find the condition under which it lands perpendicular on an inclined plane inclination α as shown in figure.



```
(A) \sin \alpha = \cos (\theta - \alpha)
(C) \tan \theta = \cot (\theta - \alpha)
```





SECTION - II : MULTIPLE CORRECT ANSWER TYPE

15. A particle is projected at an angle θ from ground with speed u (g = 10 m/s²) :

(A) if u = 10 m/s and $\theta = 30^{\circ}$, then time of flight will be 1 sec.

(B) if $u = 10\sqrt{3}$ m/s and $\theta = 60^{\circ}$, then time of flight will be 3 sec.

- (C) if $u = 10\sqrt{3}$ m/s and $\theta = 60^{\circ}$, then after 2 sec velocity becomes perpendicular to initial velocity.
- (D) if u = 10 m/s and $\theta = 30^{\circ}$, then velocity never becomes perpendicular to initial velocity during its flight.
- **16.** A particle is projected vertically upwards with a velocity u from a point O. When it returns to the point of projection :
 - (A) its average velocity is zero

(B) its displacement is zero

(C) its average speed is u/2

- (D) its average speed is u.
- 17. A particle of mass m moves along a curve $y = x^2$. When particle has x co-ordinate as 1/2 and x-component of velocity as 4m/s then:
 - (A) the position coordinate of particle are (1/2, 1/4)
 - (B) the velocity of particle will be along the line 4x 4y 1 = 0.
 - (C) the magnitude of velocity at that instant is $4\sqrt{2}$ m/s
 - (D) the magnitude of angular momentum of particle about origin at that position is 0.
- 18. A stone is projected from level ground at time t=0. Let v_x and v_y are the horizontal and vertical components of velocity at any time t; x and y are displacements along horizontal and vertical from the point of projection at any time t. Then:
 - (A) $v_v t$ graph is a straight line
 - (B) x t graph is a straight line passing through origin
 - (C) y t graph is a straight line passing through origin
 - (**D**) $v_x t$ graph is a straight line
- 19. Two balls are thrown from an inclined plane at angle of projection α with the plane one up the incline plane and other down the incline as shown in the figure. If $R_1 \& R_2$ be their respective ranges and $h_1 \& h_2$ be their respective maximum height then

(A) $h_1 = h_2$ (B) $R_2 - R_1 = T_1^2$

- (C) $R_2 R_1 = g \sin \theta T_2^2$
- **(D)** $R_2 R_1 = g \sin \theta T_1^2$

[here $T_1 \& T_2$ are times of flight in the two cases respectively]



20. A ball is projected horizontally from top of a 80 m deep well with velocity 10 m/s. Then particle will fall on the bottom at a distance of (all the collisions with the wall are elastic) :



SECTION - III : ASSERTION AND REASON TYPE

21. Statement-1 : Two stones are simultaneously projected from level ground from same point with same speed but different angles with horizontal. Both stones move in same vertical plane. Then the two stones may collide in mid air.

Statement-2: For two stones projected simultaneously from same point with same speed at different angles with horizontal, their trajectories may intersect at some point.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True
- 22. Statement-1 : During flight under action of gravity, the change in velocity of a projectile in same time intervals is same. (Neglect air friction)

Statement-2 : Neglecting air friction, the acceleration of projectile is constant during flight.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True.
- 23. Statement-1 : The velocity of a particle depends on its position vector.

Statement-2: If f' is position vector of a particle at any time t, then velocity is given by $v = \frac{df'}{dt}$.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- **(D)** Statement-1 is False, Statement-2 is True.



SECTION - IV : COMPREHENSION TYPE

Comprehension #1

The direction of velocity of a particle at time t=0 is as shown in the figure and has magnitude u = 20m/s. The acceleration of particle is always constant and has magnitude 10m/s². The angle between its initial velocity and acceleration is 127° . (Take sin $37^{\circ} = 3/5$)



(A) 0.6 sec. (B) 1.2 sec. (C) 2.4 sec.

Comprehension # 2

We know how by neglecting the air resistance, the problems of projectile motion can be easily solved and analysed. Now we consider the case of the collision of a ball with a wall. In this case the problem of collision can be simplified by considering the case of elastic collision only. When a ball collides with a wall we can divide its velocity into two components, one perpendicular to the wall and other parallel to the wall. If the collision is elastic then the perpendicular component of velocity of the ball gets reversed with the same magnitude.

(D) None of these



The other parallel component of velocity will remain constant if wall is given smooth.

Now let us take a problem. Three balls 'A' and 'B' & 'C' are projected from ground with same speed at same angle with the horizontal. The balls A,B and C collide with the wall during their flight in air and

all three collide perpendicularly with the wall as shown in figure.





27. Which of the following relation about the maximum height H of the three balls from the ground during their motion in air is correct :

(A) $H_A = H_C > H_B$ (B) $H_A > H_B = H_C$ (C) $H_A > H_C > H_B$ (D) $H_A = H_B = H_C$

28. If the time taken by the ball A to fall back on ground is 4 seconds and that by ball B is 2 seconds. Then the time taken by the ball C to reach the inclined plane after projection will be :
(A) 6 sec. (B) 4 sec. (C) 3 sec. (D) 5 sec.

29.The maximum height attained by ball 'A' from the ground is :(A) 10 m(B) 15 m(C) 20 m

(D) Insufficient information

SECTION - V : MATRIX - MATCH TYPE

30. In the column-I, the path of a projectile (initial velocity 10 m/s and angle of projection with horizontal 60° in all cases) is shown in different cases. Range 'R' is to be matched in each case from column-II. Take $g = 10 \text{ m/s}^2$. Arrow on the trajectory indicates the direction of motion of projectile.

Match each entry of column-I with its corresponding entry in column-II



31. A ball is thrown at an angle 75° with the horizontal at a speed of 20 m/s towards a high wall at a distance d. If the ball strikes the wall, its horizontal velocity component reverses the direction without change in magnitude and the vertical velocity component remains same. Ball stops after hitting the ground. Match the statement of column I with the distance of the wall from the point of throw in column II.

Column I		Column II
(A) Ball strikes the wall directly	(p)	d = 8 m
(B) Ball strikes the ground at $x = 12$ m from the wall	(q)	d = 10 m
(C) Ball strikes the ground at $x = 10$ m from the wall	(r)	d = 15 m
(D) Ball strikes the ground at $x = 5$ m from the wall	(s)	d = 25 m
	(t)	d = 30 m



SECTION - VI : INTEGER TYPE

32. Two graphs of the same projectile motion (in the xy plane) projected from origin are shown. X axis is along horizontal direction & Y axis is vertically upwards. Take $g = 10 \text{ m/s}^2$.



- 33. A stone is dropped from a height of 45 m from horizontal level ground. There is horizontal wind blowing due to which horizontal acceleration of the stone becomes 10 m/s². (Take g= 10m/s²)
 (A) Find time taken by stone to reach the ground. (in m/s)
- 34. A projectile is fired into the air from the edge of a 100 m high cliff at an angle of 37° above the horizontal. The projectile hits a target 400 m away from the base of the cliff. If initial velocity of the projectile, v_0 is $x\sqrt{5}$ m/s then x is ? (Neglect air friction and assume x-axis to be horizontal and y-axis to be vertical).

00m





4. Two particles A & B separated by a distance 2 R are moving counter clockwise along the same circular path of radius R each with uniform speed v. At time t = 0, A is given a tangential acceleration of magnitude a

$$=\frac{72 \text{ v}^2}{25 \pi \text{ R}}$$

(A) the time lapse for the two bodies to collide is $\frac{6\pi R}{5V}$

B) the angle covered by A is
$$\frac{11\pi}{6}$$

) angular velocity of A is
$$\frac{11V}{5R}$$

(D) radial acceleration of A is
$$\frac{289 \text{ v}^2}{5\text{R}}$$



(C



- (B) should turn the car in a circle of radius r.
- (C) apply the brakes and also turn the car in a circle of radius r.
- **(D)** jump on the back seat.

KINEMATICS



+91-9350679141

PHYSICS FOR JEE MAIN & ADVANCED

18. A simple pendulum is oscillating in a vertical plane. If resultant acceleration of bob of mass m at a point A is in horizontal direction, find the tangential force at this point in terms of tension T and mg.

(B) $\frac{\text{mg}}{\text{T}} \sqrt{\text{T}^2 - (\text{mg})^2}$ **(C)** $\frac{\text{mg}}{\text{T}} \sqrt{(\text{mg})^2 + \text{T}^2}$ **(D)** $\frac{\text{T}}{\text{mg}} \sqrt{(\text{mg})^2 + \text{T}^2}$

(A) mg

19. Objects A and B each of mass m are connected by light inextensible cord. They are constrained to move on a frictionless ring in a vertical plane as shown in figure. The objects are released from rest at the positions shown. The tension in the cord just after release will be

(B) $\frac{\text{mg}}{\sqrt{2}}$ (C) $\frac{\text{mg}}{2}$ (A) mg $\sqrt{2}$

20. A circular curve of a highway is designed for traffic moving at 72 km/h. If the radius of the curved path is 100 m, the correct angle of banking of the road should be given by :

 $(C) \sqrt{5gR}$

(C) $\frac{2\pi}{3}$

(B) 37°

(D) $180 + 37^{\circ}$

(A)
$$\tan^{-1} \frac{2}{3}$$
 (B) $\tan^{-1} \frac{3}{5}$ (C) $\tan^{-1} \frac{2}{5}$

A ring of radius R lies in vertical plane. A bead of mass 'm' can move along the ring 21. without friction. Initially the bead is at rest at the bottom most point on ring. The minimum constant horizontal speed v with which the ring must be pulled such that the bead completes the vertical circle

A car moves around a curve at a constant speed. When the car goes 22. around the arc subtending 60° at the centre, then the ratio of magnitude of instantaneous acceleration to average acceleration over the 60°

A bus is moving with a constant acceleration a = 3g/4 towards right. In

in the figure. The tension in the rope will be minimum, when the

(B) $\frac{\pi}{6}$

(B) $\sqrt{4qR}$

arc is :

(A) $\frac{\pi}{3}$

(A) 53°

(C) $180 - 53^{\circ}$

23.

60 5π **(D)**

mq

R

m

С

mg

mg

(D) $\tan^{-1} \frac{1}{4}$

(D) $\sqrt{5.5 \text{ gR}}$

(D)



Natural length of spring is same as that of length of rod.

ring

- 24. A ring attached with a spring is fitted in a smooth rod. The spring is fixed at the outer end of the rod. The mass of the ring is 3kg & spring constant of spring is 300 N/m. The ring is given a velocity 'V' towards the outer end of the rod. And the rod is set to be rotating with an angular velocity ω . Then ring will move with constant speed with respect to the rod if :
 - **(A)** angular velocity of rod is increased continuously
 - $\omega = 10 \text{ rad/s}$ **(B)**

rope makes an angle $\theta =$

- angular velocity of rod is decreased continuously. **(C)**
- constant velocity of ring is not possible. **(D)**



25. A uniform rod of mass m and length ● is rotating with constant angular velocity ω about an axis which passes through its one end and perpendicular to the length of rod. The area of cross section of the rod is A and its Young's modulus is Y. Neglect gravity. The strain at the mid point of the rod is :

(A)
$$\frac{m\omega^2 \lambda}{8AY}$$

26. A body moves along an uneven surface with constant speed at all points. The normal reaction of the road on the body is:

(B) $\frac{3m\omega^2 \lambda}{8AY}$ (C) $\frac{3m\omega^2 \lambda}{4AY}$ (D) $\frac{m\omega^2 \lambda}{4AY}$

(A) maximum at A

(B) maximum at B

(C) minimum at C

(D) the same at A, B & C

SECTION - II : MULTIPLE CORRECT ANSWER TYPE

- 27. A particle is describing circular motion in a horizontal plane in contact with the smooth inside surface of a fixed right circular cone with its axis vertical and vertex down. The height of the plane of motion above the vertex is h and the semivertical angle of the cone is α . The period
 - of revolution of the particle:
 - (A) increases as h increases(C) increases as α increases

- (B) decreases as h increases
- (D) decreases as α increases
- 28. A machine, in an amusement park, consists of a cage at the end of one arm, hinged at O. The cage revolves along a vertical circle of radius r

(ABCDEFGH) about its hinge O, at constant linear speed $v = \sqrt{gr}$. The

cage is so attached that the man of weight 'w' standing on a weighing machine, inside the cage, is always vertical. Then which of the following is/are correct

- (A) the weight reading at A is greater than the weight reading at E by 2 w.
- **(B)** the weight reading at G = w
- (C) the ratio of the weight reading at E to that at A = 0
- (D) the ratio of the weight reading at A to that at C = 2.
- 29. A small sphere of mass m suspended by a thread is first taken aside so that the thread forms the right angle with the vertical and then released, then :
 - (A) total acceleration of sphere as a function of θ is $g\sqrt{1+3\cos^2\theta}$
 - (B) thread tension as a function of θ is T = 3mg cos θ
 - (C) the angle θ between the thread and the vertical at the moment when the total acceleration vector of the sphere is directed horizontally is $\cos^{-1} 1/\sqrt{3}$
 - (D) thread tension at the moment when the vertical component of the sphere's velocity is maximum will be mg



PHYSICS FOR JEE MAIN & ADVANCED

30. On a circular table, A and B are moving on the circumference. Man A runs behind man B to catch him. A runs with constant angular speed ω_1 with respect to table and B runs at constant tangential speed v_2 with respect to ground. If it is found that the table rotates 30° in the opposite direction in every one second and the initial angular separation between A and B is 30°, then A catches B after : (Radius of table is 3 m).

(A) 0.5 sec, if
$$\omega_1 = \frac{5\pi}{6}$$
 rad/s and $v_2 = 3.14$ m/s

(B) 0.5 sec, if
$$\omega_1 = \frac{4\pi}{3}$$
 rad/s and $v_2 = 3.14$ m/s

(C) 0.5 sec, if
$$\omega_1 = \frac{4\pi}{3}$$
 rad/s and $v_2 = 6.28$ m/s

(D) A can not catch B within 0.5 s, if $\omega_1 = \frac{\pi}{6}$ rad/s and $v_2 = 6.28$ m/s

SECTION - III : ASSERTION AND REASON TYPE

31. Statement-1: Two small spheres are suspended from same point O on roof with strings of different lengths. Both spheres move along horizontal circles as shown. Then both spheres may move along circles in same horizontal plane.

Statement-2: For both spheres in statement-1 to move in circular paths in same horizontal plane, their angular speeds must be same.

(A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.

- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True
- 32. Statement-1 : A car moves along a road with uniform speed. The path of car lies in vertical plane and is shown in figure. The radius of curvature(R) of the path is same everywhere. If the car does not loose contact with road at the highest point, it can travel the shown

path without loosing contact with road anywhere else.



<u>//////</u>

Statement-2 : For car to loose contact with road, the normal reaction between car and road should be zero.

(A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.

- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True
- 33. Statement-1: A ball tied by thread is undergoing circular motion (of radius R) in a vertical plane. (Thread always remains in vertical plane). The difference of maximum and minimum tension in thread is independent of speed (u) of ball at the lowest position ($u > \sqrt{5gR}$)

Statement-2: For a ball of mass m tied by thread undergoing vertical circular motion (of radius R), difference in maximum and minimum magnitude of centripetal acceleration of the ball is independent of speed (u) of ball at the lowest position $(u \ge \sqrt{5\alpha R})$

at the lowest position ($u > \sqrt{5gR}$).

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True



KINEMATICS

O C

(D) zero

SECTION - IV : COMPREHENSION TYPE

Comprehension #1

One end of a light string of length L is connected to a ball and the other end is connected to a fixed point O. The ball is released from rest at t = 0 with string horizontal and just taut. The ball then moves in vertical circular path as shown. The time taken by ball to go from position A to B is t_1 and from B to lowest position C is t_2 . Let the velocity of ball at B is \breve{V}_B and at C is \breve{V}_C respectively.

34. If $|\breve{V}_{C}| = 2|\breve{V}_{B}|$ then the value of θ as shown is (A) $\cos^{-1}\frac{1}{4}$ (B) $\sin^{-1}\frac{1}{4}$ (C) $\cos^{-1}\frac{1}{2}$ (D) $\sin^{-1}\frac{1}{2}$ 35. If $|\breve{V}_{C}| = 2|\breve{V}_{B}|$ then : (A) $t_{1} > t_{2}$ (B) $t_{1} < t_{2}$ (C) $t_{1} = t_{2}$ (D) Information insufficient

36. If
$$|\nabla_{C} - \nabla_{B}| = |\nabla_{B}|$$
, then the value of θ as shown is :
(A) $\cos^{-1}\left(\frac{1}{4}\right)^{1/3}$
(B) $\sin^{-1}\left(\frac{1}{4}\right)^{1/3}$
(C) $\cos^{-1}\left(\frac{1}{2}\right)^{1/3}$
(D) $\sin^{-1}\left(\frac{1}{2}\right)^{1/3}$

Comprehension #2

A particle of mass M attached to an inextensible string is moving in a vertical circle of radius R about fixed point O. It is imparted a velocity u in horizontal direction at lowest position as shown in figure.

Following information is being given :

- (i) Velocity at a height h can be calculated by using formula $v^2 = u^2 2gh$.
- (ii) Particle will complete the circle if $u \ge \sqrt{5gR}$,
- (iii) Particle will oscillates in lower half $(0^{\circ} < \theta \le 90^{\circ})$ if $0 < u \le \sqrt{2gR}$
- (iv) The magnitude of tension at a height 'h' is calculated by using formula $T = \frac{M}{R} \{u^2 + gR 3gh\}$.



37. If R = 2m, M = 2 kg and u = 12 m/s. Then value of tension at lowest position is :(A) 120 N(B) 164 N(C) 264 N

38.Tension at highest point of its trajectory in above question will be :(A) 100 N(B) 44 N(C) 144 N(D) 264 N

39. If M = 2 kg, R = 2m and u = 10 m/s the velocity of particle when $\theta = 60^{\circ}$,

(A) $2\sqrt{5}$ m/s (B) $4\sqrt{5}$ m/s (C) $5\sqrt{2}$ m/s (D) 5 m/s


	SECTION - V : MATRIX - N	IATCH	ТҮРЕ
40.	In column-I condition on velocity, force and acceleration described in column-II. \mathbf{U} = initial velocity, \mathbf{F} = resultar	of a part nt force a	ticle is given. Resultant motion is nd \bigvee_{ν}^{μ} = instantaneous velocity.
	Column-I		Column-II
	(A) $\mathbf{\hat{u}} \times \mathbf{\hat{F}} = 0$ and $\mathbf{\hat{F}} = \text{constant}$	(p)	path will be circular path
	(B) $\mathbf{\hat{u}} \cdot \mathbf{\hat{F}} = 0$ and $\mathbf{\hat{F}} = \mathbf{constant}$	(q)	speed will increase
	(C) $\bigvee_{v} \stackrel{P}{H} = 0$ all the time and $ \stackrel{P}{H} = \text{constant}$ and the particle always remains in one plane.	(r)	path will be straight line
	(D) $\hat{\mathbf{u}} = 2\hat{\mathbf{i}} - 3\hat{\mathbf{j}}$ and acceleration at all time $\hat{\mathbf{a}} = 6\hat{\mathbf{i}} - 9\hat{\mathbf{j}}$	(s) (t)	path will be parabolic Particle may retrace back
41.	A particle is moving with speed $v = 2t^2$ on the circumference in column-I with corresponding results in column-II	le of radius R. Match the quantities given	
	(A) Magnitude of tangential acceleration of particle	(p)	decreases with time.
	(B) Magnitude of Centripetal acceleration of particle	(q)	increases with time
	(C) Magnitude of angular speed of particle with respect to centre of circle	(r)	remains constant
	(D) Value of tan θ, where θ is angle between the total acceleration vector and centripetal acceleration vector	(8)	proportional to R
	of particle		incomplete and a set in the D
42.	Each situation in column I gives graph of a particle moving	(t) in circul	inversely proportional to K ar path. The variables $\omega \theta$ and t represent

42. Each situation in column I gives graph of a particle moving in circular path. The variables ω, θ and t represent angular speed (at any time t), angular displacement (in time t) and time respectively. Column II gives certain resulting interpretation. Match the graphs in column I with statements in column II and indicate your answer by darkening appropriate bubbles in the 4 × 4 matrix given in the OMR.







SECTION - VI : INTEGER TYPE

- 43. A ball is projected making an angle θ with the vertical. Consider a small part of the trajectory near the highest position and take it approximately to be a circular arc. What is the radius of this circle? This radius is called the radius of curvature of the curve at the point.if u = 20 then ROC in m.
- 44. A rod AB is moving on a fixed circle of radius R with constant velocity 'v' as shown in figure. P is the point of intersection of the rod and the circle. At an instant the rod is at a distance $x = \frac{3R}{5}$ from centre of the circle. The velocity of the rod is perpendicular to the rod and the rod is always parallel to the diameter CD.



- (A) the speed of point of intersection P is xv/4 m./sec. then x is
- (B) Find the angular speed of point of intersection P with respect to centre of the circle.
- 45. A toy car of mass m can travel at a fixed speed. It moves in a circle

on a fixed horizontal table. A string is connected to car and attached to a block of mass M that hangs as shown in figure (the portion of string below the table is always vertical). The coefficient of friction between the surface of table and tyres of the toy car is μ . Find the ratio of the maximum radius to the minimum radius for which the toy car can move in a circular

path with centre O on table.





46. A particle of mass m is attached at one end of a light, inextensible string of length \bullet whose other end is fixed at the point C. The particle is given minimum velocity at the lowest point to complete the circular path in the vertical plane. As it moves in the circular path the tension in the string changes with θ . θ is defined in the figure. As θ varies from 'O' to '2\pi' (i.e. the particle completes one revolution) plot the variation of tension 'T' against ' θ '.



47. A small wedge whose base is horizontal is fixed to a vertical rod as shown in the figure $\theta = 45^{\circ} l = \frac{1}{5\sqrt{2}}$ m. The sloping side of the wedge is frictionless and the wedge is spun with a constant angular speed ω about vertical axis as shown in the figure. Find



(A) The value of angular speed ω for which the block of mass m just does not slide down the wedge?





- (A) 2 sec
- **(B)** 3 sec
- (C) 1 sec
- (D) 6 sec



7. A man in a balloon, throws a stone downwards with a speed of 5 m/s with respect to balloon. The balloon is moving upwards with a constant acceleration of 5 m/s². Then velocity of the stone relative to the man after 2 second is :





Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035 +91-9350679141

PHYSICS FOR JEE MAIN & ADVANCED

- 8. Three stones A, B and C are simultaneously projected from same point with same speed. A is thrown upwards, B is thrown horizontally and C is thrown downwards from a building. When the distance between A and C becomes 10 m, then distance between A and B will be:
 - (A) 10 m (B) 5 m (C) $5\sqrt{2}$ m (D) $10\sqrt{2}$ m
- 9. A projectile A is projected from ground. An observer B running on ground with uniform velocity of magnitude 'v' observes A to move along a straight line. The time of flight of A as measured by B is T. Then the range R of projectile on ground is :

(B) R < vT

(A) R = vT(C) R > vT

(D) information insufficient to draw inference

- 10. Two aeroplanes fly from their respective position 'A' and 'B' starting at the same time and reach the point 'C' (along straight line) simultaneously when wind was not blowing. On a windy day they head towards 'C' but both reach the point 'D' simultaneously in the same time which they took to reach 'C'. Then the wind is blowing in:
 - (A) North-East direction
 - (B) North-West direction
 - (C) Direction making an angle $0 < \theta < 90$ with North towards West.
 - **(D)** North direction



- 11. A particle is thrown up inside a stationary lift of sufficient height. The time of flight is T. Now it is thrown again with same initial speed v_0 with respect to lift. At the time of second throw, lift is moving up with speed v_0 and uniform acceleration g upward (the acceleration due to gravity). The new time of flight is:
 - (A) $\frac{T}{4}$ (B) $\frac{T}{2}$ (C) T (D) 2T

12. A swimmer crosses a river with minimum possible time 10 second. And when he reaches the other end starts swimming in the direction towards the point from where he started swimming. Keeping the direction fixed the swimmer crosses the river in 15 sec. The ratio of speed of swimmer with respect to water and the speed of river flow is (Assume constant speed of river & swimmer):

- (A) $\frac{3}{2}$ (B) $\frac{9}{4}$ (C) $\frac{2}{\sqrt{5}}$ (D) $\frac{\sqrt{5}}{2}$
- A taxi leaves the station X for station Y every 10 minutes. Simultaneously, a taxi also leaves the station Y for station X every 10 minutes. The taxis move at the same constant speed and go from X to Y or vice versa in 2 hours. How many taxis coming from the other side will meet each taxi enroute from Y to X:

 (A) 11
 (B) 12
 (C) 23
 (D) 24
- 14. Consider a collection of a large number of particles each with speed v. The direction of velocity is randomly distributed in the collection. The magnitude of the relative velocity between a pair of particles averaged over all the pairs in the collection is :

```
(A) greater than v (B) less than v (C) equal to v (D) we can't say anything
```

- 15. An aeroplane is flying in geographic meridian vertical plane at an angle of 30° with the horizontal (north) and wind is blowing from west. A package is dropped from an aeroplane. The velocity of the wind if package hits
 - a kite flying in the space with a position vector $\vec{R} = (400\sqrt{3}\hat{i} + 80\hat{j} + 200\hat{k})$ m with respect to the point of

dropping.(Here \hat{j} and \hat{j} are the unit vectors along north and vertically up respectively and \hat{k} be the unit vector due east. Assume that the bag is light enough to get carried away by the wind) : (A) 50 m/sec (B) 25 m/sec (C) 20 m/sec (D) 10 m/sec



16. P is a point moving with constant speed 10 m/s such that its velocity vector always maintains an angle 60° with line OP as shown in figure (O is a fixed point in space). The initial distance between O and P is 100 m. After what time shall P reach O.



SECTION - II : MULTIPLE CORRECT ANSWER TYPE

17. A river is flowing with a speed of 1 km/hr. A swimmer wants to go to point 'C' starting from 'A'. He swims with a speed of 5 km/hr, at an angle θ w.r.t. the river. If AB = BC = 400 m. Then :



(A) the value of θ is 53°

(A) 10 sec.

- (C) time taken by the man is 8 min
- **(D)** the value of θ is 45°
- 18. A swimmer who can swim in a river with speed *mv* (with respect to still water) where *v* is the velocity of river current, jumps into the river from one bank to cross the river:
 - (A) If $m \le 1$ he can not reach a point on other bank directly opposite to his starting point
 - **(B)** If m < 1 he can not cross the river
 - (C) If m > 1 then only he can reach a point on other bank
 - (D) He can reach the other bank at some point, whatever be the value of m.
- 19. A man is standing on a road and observes that rain is falling at angle 45° with the vertical. The man starts running on the road with constant acceleration 0.5 m/s². After a certain time from the start of the motion, it

appears to him that rain is still falling at angle 45° with the vertical, with speed $2\sqrt{2}$ m/s. Motion of the man is in the same vertical plane in which the rain is falling. Then which of the following statement(s) are true:

- (A) It is not possible
- (B) Speed of the rain relative to the ground is 2 m/s.
- (C) Speed of the man when he finds rain to be falling at angle 45° with the vertical, is 4m/s.
- (D) The man has travelled a distance 16m on the road by the time he again finds rain to be falling at angle 45°.
- 20. Two stones are thrown vertically upwards simultaneously from the same point on the ground with initial speed $u_1 = 30$ m/sec and $u_2 = 50$ m/sec. Which of the curve represents correct variation (for the time interval in which both reach the ground) of
 - $(x_2 x_1) =$ the relative position of second stone with respect to first with time (t).

 $(v_2 - v_1) =$ the relative velocity of second stone with respect to first with time (t).

Assume that stones do not rebound after hitting the ground





SECTION - III : ASSERTION AND REASON TYPE

- 21. Statement-1: The magnitude of velocity of two boats relative to river is same. Both boats start simultaneously from same point on one bank may reach opposite bank simultaneously moving along different paths.
 Statement-2: For boats to cross the river in same time. The component of their velocity relative to river in direction normal to flow should be same.
 - (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
 - (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
 - (C) Statement-1 is True, Statement-2 is False
 - (D) Statement-1 is False, Statement-2 is True
- 22. Statement-1 : Three projectiles are moving in different paths in the air. Vertical component of relative velocity between any of the pair does not change with time as long as they are in air. Neglect the effect of air friction. Statement-2 : Relative acceleration between any of the pair of projectiles is zero.
 - (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
 - (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
 - (C) Statement-1 is True, Statement-2 is False
 - (D) Statement-1 is False, Statement-2 is True
- 23. STATEMENT-1 : Two stones are projected with different velocities from ground from same point and at same instant of time. Then these stones cannot collide in mid air. (Neglect air friction)

STATEMENT-2: If relative acceleration of two particles initially at same position is always zero, then the distance between the particle either remains constant or increases continuously with time.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True.

SECTION - IV : COMPREHENSION TYPE

A swimmer can swim with a speed v in still water.



24. If the swimmer crosses a swimming pool 'd' from A to directly opposite point B on other side in time t_1 as shown in figure (i) and in a flowing river (river velocity 'u') of same width d from A' to directly opposite point B' on other bank in time t_2 , then (t_1/t_2) is equal to : (Assume v > u)

(A)
$$\sqrt{1 - \frac{v^2}{u^2}}$$
 (B) $\sqrt{1 + \frac{u^2}{v^2}}$ (C) $\sqrt{1 - \frac{u^2}{v^2}}$ (D) 1

25.

If the minimum time taken in swimming pool to reach opposite bank is t_1' and minimum. time to reach opposite bank

in river is t_2' , then the ratio $\frac{t_1'}{t_2'}$ will have a value :

(B)
$$\frac{\sqrt{v^2 + u^2}}{v}$$
 (C) $\frac{\sqrt{v^2 - u^2}}{u}$ (D) $\frac{u}{v}$



(A) 1

26. If the time taken by swimmer to reach opposite point on other bank in river is T_1 and the time taken to travel an equal

distance upstream (against the water current) in the river is T_2 , then ratio $\frac{T_2}{T_1}$ will have a value :

(A)
$$\sqrt{\frac{1-u/v}{1+u/v}}$$
 (B) $\sqrt{\frac{1+u/v}{1-u/v}}$ (C) $\frac{\sqrt{v^2-u^2}}{(v+u)}$ (D) $\frac{\sqrt{v^2-u}}{v}$

Comprehension - 2

Raindrops are falling with a velocity $10\sqrt{2}$ m/s making an angle of 45° with the vertical. The drops appear to be falling vertically to a man running with constant velocity. The velocity of rain drops change such that the rain drops now appear to be falling vertically with $\sqrt{3}$ times the velocity it appeared earlier to the same person running with same velocity.

27. The magnitude of velocity of man with respect to ground is:

(A) 10√2 m/s	(B) 10√3 m/s	(C) 20 m/s	(D) 10 m/s
--------------	---------------------	------------	-------------------

28. After the velocity of rain drops change, the magnitude of velocity of raindrops with respect to ground is:

	(A) 20 m/s	(B) $20\sqrt{3}$ m/s	(C) 10 m/s	(D) 10√3 m/s
--	------------	-----------------------------	------------	---------------------

29. The angle (in degrees) between the initial and the final velocity vectors of the raindrops with respect to the ground is:

(A) 8 (B) 15 (C) 22.5 (D) 37

SECTION - V : MATRIX - MATCH TYPE

30. Two particles A and B moving in x-y plane are at origin at t = 0 sec. The initial velocity vectors of A and B are $\ddot{u}_A = 8\hat{j}$ m/s and $\ddot{u}_B = 8\hat{j}$ m/s. The acceleration of A and B are constant and are $\ddot{u}_A = -2\hat{j}$ m/s² and $\ddot{u}_B = -2\hat{j}$ m/s². Column I gives certain statements regarding particle A and B. Column II gives corresponding results. Match the statements in column I with corresponding results in Column II.

	Column I		Column II
(A)	The time (in seconds) at which velocity	(p)	16 √2
	of A relative to B is zero		
(B)	The distance (in metres) between A and B	(q)	8√2
	when their relative velocity is zero.		
(C)	The time (in seconds) after $t = 0$ sec, at which A and B are at same position	(r)	8
(D)	The magnitude of relative velocity of A and B	(s)	4
	at the instant they are at same position.		
		(t)	6 secconds

31. Two particles A & B are projected as shown in fig in x–y plane. Under the effect of force which provide a constant acceleration a =11 m/s² in negative y-direction. Then match situation in column-I with the corresponding results in column-II (All positions are given in metre) (\vec{v}_{AB} = velocity of A w.r.t. B; \vec{F}_{AB} = Position of A w.r.t. B).





32. When two bodies move uniformly towards each other, the distance between them diminishes by 16 m every 10 s. If velocity of one body is reversed the distance between them will decrease 3 m every 5 s. Calculate the

speed of second body is $\frac{10}{x}$ m/s then x is.

33. A swimmer jumps from a bridge over a canal and swims 1km upstream. After that first km, he passes a floating cork. He continues swimming for half an hour and then turns around and swims back to the bridge. The swimmer and the cork reach the bridge at the same time. The swimmer has been swimming at a constant speed. How fast does the water in the canal flow?



ANSWER KEY

EXERCISE - 1

1. B 2. D 3. B 4. D 5. A 6. D 7. C 8. C 9. C 10. D 11. A 12. B 13. A 14. A 15. B 16. C 17. D 18. C 19. A 20. A 21. C 22. A 23. B 24. B 25. A 26. B 27. A 28. C 29. C **30.** B **31.** C **32.** B **33.** C **34.** B **35.** C **36.** A **37.** B **38.** B **39.** B 40. D 41. C 42. C 43. B 44. C 45. C 46. A 47. C 48. A 49. D 50. D 51. A

EXERCISE - 2 : PART # I

1.	С	2.	В	3.	А	4.	А	5.	А	6.	D	7.	С	8.	А	
9.	А	10.	В	11.	А	12.	С	13.	D	14.	C	15.	В	16.	A,B	
17.	С	18.	С	19.	С	20.	В	21.	D	22.	С	23.	Α	24.	D	
25.	В	26.	A,B,C	,D	27.	A,B,D	28.	A,C,D	29.	AD	30.	A,C,D	31.	A,B	32.	B,C,D
33.	C,D	34.	A,B,C	35.	B,C	36.	A,C,D	37.	A,C	38.	С	39.	В	40.	B,C	
41.	A,D	42.	A,B	43.	A,C	44.	B,C,D	45.	B,C,D	46.	А	47.	A,B,C	48.	С	
49.	В	50.	D	51.	D	52.	С	53.	А	54.	D	55.	С	56.	B 57	. A

PART # II

1. D 2. A 3. D 4. B 5. D 6. A 7. A 8. A 9. C 10. A 11. A 12. D 13. A 14. C 15. A 16. B 17. A 18. D 19. A

EXERCISE - 3 : PART # I

1. $A \rightarrow P$; $B \rightarrow V$, $C \rightarrow S$, D, $\rightarrow Q$, $E \rightarrow U$, $F \rightarrow T$, $G \rightarrow R$ **2.** (A) \rightarrow Q, (B)–p, q (C)–p, r, (D)–r, s 3. A-r, B-p, C-r, D-s 4. (A)-r, (B)-p, (C)-s, (D)-q5. A-r, B-q, C-r, D-p 6. (A) -s, (B)-r, (C)-q

PART # II

Comp. #1 : 1. (D) 2. (B) 3. (A) 4. (B) 5. (C) Comp. #3 : 1. (B) 2. (B) 3. (A) 4. (A) 5. (D) Comp. #5 : 1. (C) 2. (A) 3. (B) Comp. #7 : 1. (C) 2. (B) 3. (C) 4. (C)

Comp. #2 : 1. (A) 2. (D) 3. (A) **Comp.** #4: **1.** (D) **2.** (C) **3.** (A) **4.** (C) **5.** (B) **6.** (D) **Comp. #6:** 1. (C) 2. (A) 3. (B)

EXERCISE - 4

1. (i) A, B (ii) A, B (iii) B, A (iv) same (v) B, A, once **2**.3 3. $a_0 T/3$ 4. object, 3.3 s **6**. 18.5 m 7.12 km **8**. (i) 17.1s (ii) 293.8 m **9**. (i) 20 m (ii) 5s **10.** 1.5 m/s, 1 m/s 5.36 m **11**. 5s **12**. (i) 160 km/hr² (ii) 10 km **13**. $v_{A} = 10 \text{ ms}^{-1}$, $v_{B} = 30 \text{ ms}^{-1}$ 14.89m

15. 0.75 hr, 1.5 hr after the moment 16. east of departure, No overtaking

17. $\frac{a}{v\left(1-\cos\frac{2\pi}{r}\right)}$ 18. $\frac{4}{3} \frac{v\sin\theta}{g}$

19. 200m, $v_{\text{boat}} = 1.20 \text{ km/hr}$, $v_{\text{water}} = 0.72 \text{ km/hr}$, **20**. $u_{\hat{i}} + \sqrt{\sqrt{3}\text{Rg}} \hat{j}$, $\sqrt{\frac{\sqrt{3}\text{R}}{g}}$ **21**. 20 m

22. (i) 100 m/s(ii) 980 m(iii) 1600 m(iv) $\left(80\,\hat{i} - 140\,\hat{j}\right)$ **23**. 360 km/hr, 1.28 km **24**.20 m **25**. $v_0 = \sqrt{gH\left(1 + \frac{1^2}{H^2}\right)}$, $\tan \alpha = \frac{H}{1}$



26.	$\frac{a_0}{2}$ 27. $-3\hat{k}$ rad/ s^2 , $-2\hat{k}$ rad	d/s 27.4 $\sqrt{2}$ ms ⁻²	29 . $\frac{1}{t^2}$	30. 14.3s,0.45 ms ⁻²	
31.	$\frac{5\pi R}{6v}$, $\frac{11\pi}{6}$, $\frac{17v}{5R}$, $\frac{289v^2}{25R}$	32. (i) t= 2s, (ii) 3.14 m/s	33. (i) $25\frac{\sqrt{3}}{2}$ m/s	$(125\frac{\sqrt{3}}{4})^{1/2}$ m/s (iii) $\frac{25}{2}$ m/s	\mathbf{s}^2
34.	(i) $\lambda = \frac{1}{6}$, (ii) 800m 35 . 9	0 min, 40 km hr ⁻¹ 36 . 75 ms	s^{-1} , (437.5 ± 62.5	√21)m	
37.	${}_{s}^{r} = 10(t-1)\hat{i} + 10\hat{j}(a)t = \frac{1}{2}$	h,10km (b) $\frac{10\sqrt{5}}{2}$, t = 2 hr	38 . 80 ms ⁻¹ 3	9. $\frac{(v_1 + v_2)^2}{2(a_1 + a_2)}$ 40. (a) 120° (b) 2/ $\sqrt{2}$	3
41.	B, $t_A = 165 \text{ s}, t_B = 150 \text{ s}$ 42 .	50 ms ⁻¹ 43 .10 $\sqrt{3}$ ms ⁻¹ 44 .	$\frac{2}{\sqrt{2}+1}$ 45 . (i) 2s ((ii) 10 ms ⁻¹ (iii) 5m(iv) 16.25 m (v) 20	m
		EVEDCISE 5	• DADT # I		

1. 3 **2.** 4 **3.** 1 **4.** 4 **5.** 3 **6.** 1 **7.** 4 **8.** 1 **9.** 2 **10.** 1 **11.** 1

PART # II

MCQ's with one correct answer	1. B 2. A 3	в. В	4. C	5. A		
Assertion-Reasoning	1. B					
MCO's with one or more than one correct answer 1. ABC						

Subjective Questions 1. (i) $\frac{u^2 \sin 2\alpha}{g \cos \theta}$ (ii) $v = \frac{u \cos(\alpha + \theta)}{\cos \theta}$ 2. $u=7.29 \text{ ms}^{-1}, t=1s$ 3. (i) 45° (ii) 2 ms^{-1}

Integer Type Question1.52.53.2 or 84.4Comprehension Type Questions1.A2.B

MOCK TEST : RECTILINEAR MOTION

 1. B
 2. C
 3. A
 4. D
 5. D
 6. C
 7. B
 8. A
 9. A
 10. D
 11. D
 12. B
 13. A

 14. B
 15. A
 16. A
 17. B,C,D
 18. A,C,D
 19. A,C,D
 20. A,C,D
 21. A,C,D
 22. A,B
 23. D

 24. D
 25. D
 26. B
 27. A
 28. B
 29. B
 30. D

 31.
 31.
 31.
 33. A
 33. A
 33. A
 33. A
 33. A



AB = Slope increasing, BC = Slope constant,

CD = Slope increasing

DE = Slope decreasing, EF = Slope increasing

F = Slope is 0

32. (A) **33.** (A) r,t (B) r,t (C) p,q,r (D) p,q,r **34.** (A) p, q (B) p, q (C) p, r (D) r, t **35.** 120 **36.** 500 **37.** 35



MOCK TEST : PROJECTILE MOTION

 1. C
 2. D
 3. D
 4. D
 5. C
 6. D
 7. B
 8. C
 9. B
 10. C
 11. A
 12. B
 13. D

 14. D
 15. A,B,C,D
 16. A,B,C
 17. A,B,C
 18. A,B,D
 19. A,C,D
 20. B,C
 21. D
 22. A
 23. D

 24. B
 25. B
 26. C
 27. A
 28. C
 29. C
 30. A-r; B-p; C-s; D-q

 31. A-p,q,r; B-p; C-q,t; D-r,s
 32. 4
 33. 3
 34. 25

MOCK TEST : CIRCULAR MOTION

 1. D
 2. C
 3. A
 4. B
 5. C
 6. C
 7. D
 8. C
 9. D
 10. D
 11. A
 12. A
 13. C

 14. D
 15. C
 16. A
 17. C
 18. B
 19. B
 20. C
 21. B
 22. A
 23. A
 24. B
 25. B
 26. A

 27. A,C
 28. A,B,C,D
 29. A,B,C
 30. A,D
 31. A
 32. D
 33. A
 34. B
 35. B
 36. B
 37. B

 38. B
 39. B
 40. A - r, t; B - q,s; C - p; D - q,r
 31. A
 32. D
 33. A
 34. B
 36. B
 37. B

41. A - q; B - q, t; C - q, t; D - p, s **42.** A - q,s; B - p,t; C - p,t; D - q,r **43.** 20

44. (A)(5) (B) $\omega = \frac{V_P}{R} = \frac{5V}{4R}$ 45. $\frac{r_{max}}{r_{min}} = \frac{M + \mu m}{M - \mu m}$ 46. $3mg (1 + \cos\theta) = 6 mg \cos^2(\theta/2)$

47. $\omega = \sqrt{\frac{g\sin\theta}{\lambda\cos^2\theta}}$

MOCK TEST : RELATIVE MOTION

 1. C
 2. A
 3. A
 4. C
 5. C
 6. B
 7. D
 8. C
 9. A
 10. B
 11. B
 12. C
 13. C

 14. A
 15. B
 16. C
 17. A,B
 18. A,D
 19. C,D
 20. A,D
 21. A
 22. A
 23. A
 24. C

 25. A
 26. B
 27. D
 28. A
 29. B
 30. A-s; B-p; C-r; D-q
 31. A-r; B-t; C-p, s; D-p

 32. 5
 33. 1

