




SOLVED EXAMPLES

Ex. 1 Which of the following pictorial diagrams represent the function



Sol. B and D. In (A) one element of domain has no image, while in (C) one element of 1st set has two images in 2nd set

Ex. 2 Find the Domain of the following function :

(i) $y = \log_{(x-4)}(x^2 - 11x + 24)$

(ii) $f(x) = \sqrt{x^2 - 5}$

(iii) $\sin^{-1}(2x - 1)$

(iv) $f(x) = \sqrt{\sin x} - \sqrt{16 - x^2}$

Sol. (i) $y = \log_{(x-4)}(x^2 - 11x + 24)$

Here 'y' would assume real value if,

$$x - 4 > 0 \text{ and } \neq 1, x^2 - 11x + 24 > 0$$

$$\Rightarrow x > 4 \text{ and } \neq 5, x < 3 \text{ or } x > 8$$

$$\Rightarrow \text{Domain}(y) = (8, \infty)$$

$$x > 4 \text{ and } \neq 5, (x - 3)(x - 8) > 0$$

$$\Rightarrow x > 8$$

(ii) $\sqrt{x^2 - 5} \quad f(x) = \text{is real iff } x^2 - 5 \geq 0$

$$\Rightarrow |x| \geq \sqrt{5} \Rightarrow x \leq -\sqrt{5} \text{ or } x \geq \sqrt{5}$$

$$\therefore \text{the domain of } f \text{ is } (-\infty, -\sqrt{5}] \cup [\sqrt{5}, \infty)$$

(iii) $\sin^{-1}(2x - 1)$ is real iff $-1 \leq 2x - 1 \leq +1$

$$\therefore \text{domain is } x \in [0, 1]$$

(iv) $\sqrt{\sin x}$ is real iff $\sin x \geq 0 \Leftrightarrow x \in [2n\pi, 2n\pi + \pi], n \in \mathbb{I}$.

$$\sqrt{16 - x^2} \text{ is real iff } 16 - x^2 \geq 0 \Leftrightarrow -4 \leq x \leq 4.$$

Thus the domain of the given function is $\{x : x \in [2n\pi, 2n\pi + \pi], n \in \mathbb{I}\} \cap [-4, 4] = [-4, -\pi] \cup [0, \pi]$.

Ex. 3 Find the range of following functions :

(i) $f(x) = \frac{1}{8 - 3\sin x}$

(ii) $f(x) = \frac{x^2 - 4}{x - 2}$

Sol. (i) $f(x) = \frac{1}{8 - 3\sin x}$


$$-1 \leq \sin x \leq 1$$

$$\therefore \text{Range of } f = \left[\frac{1}{11}, \frac{1}{5} \right]$$

(ii) $f(x) = \frac{x^2 - 4}{x - 2} = x + 2; x \neq 2$

\therefore graph of $f(x)$ would be

Thus the range of $f(x)$ is $\mathbb{R} - \{4\}$



Ex. 4 Find the range of following functions :

(i) $y = \bullet n(2x - x^2)$

(ii) $y = \sec^{-1}(x^2 + 3x + 1)$

Sol. (i) Step - 1

We have $2x - x^2 \in (-\infty, 1]$

Step - 2

Let $t = 2x - x^2$

For $\bullet nt$ to be defined accepted values are $(0, 1]$

Now, using monotonicity of $\bullet nt$,


$\bullet n(2x - x^2) \in (-\infty, 0]$

\therefore range is $(-\infty, 0]$

(ii) $y = \sec^{-1}(x^2 + 3x + 1)$

Let $t = x^2 + 3x + 1$ for $x \in \mathbb{R}$, then $t \in \left[-\frac{5}{4}, \infty\right)$

but $y = \sec^{-1}(t) \Rightarrow t \in \left[-\frac{5}{4}, -1\right] \cup [1, \infty)$



from graph the range is $\left[0, \frac{\pi}{2}\right) \cup \left[\sec^{-1}\left(-\frac{5}{4}\right), \pi\right]$

Ex. 5 (i) Let $\{x\}$ and $[x]$ denote the fractional and integral part of a real number x respectively.

Solve $4\{x\} = x + [x]$

(ii) Draw graph of $f(x) = \text{sgn}(\bullet n x)$

Sol. (i)


As $x = [x] + \{x\}$

\therefore Given equation $\Rightarrow 4\{x\} = [x] + \{x\} + [x] \Rightarrow \{x\} = \frac{2[x]}{3}$

As $[x]$ is always an integer and $\{x\} \in [0, 1)$, possible values are

$[x]$	$\{x\}$	$x = [x] + \{x\}$
0	0	0
1	$\frac{2}{3}$	$\frac{5}{3}$

\therefore There are two Solution of given equation $x = 0$ and $x = \frac{5}{3}$



Ex. 6 Find the domain $f(x) = \frac{1}{\sqrt{[\lceil x \rceil - 5] - 11}}$ where $[\cdot]$ denotes greatest integer function.

Sol. $[\lceil x \rceil - 5] > 11$

So $[\lceil x \rceil - 5] > 11$ or $[\lceil x \rceil - 5] < -11$

$[\lceil x \rceil] > 16$ $[\lceil x \rceil] < -6$

$|x| \geq 17$ or $[\lceil x \rceil] < -6$ (Not Possible)

$\Rightarrow x \leq -17$ or $x \geq 17$

So $x \in (-\infty, -17] \cup [17, \infty)$

Ex. 6 Examine whether following pair of functions are identical or not ?

(i) $f(x) = \frac{x^2 - 1}{x - 1}$ and $g(x) = x + 1$

(ii) $f(x) = \sin^2 x + \cos^2 x$ and $g(x) = \sec^2 x - \tan^2 x$

Sol. (i) No, as domain of $f(x)$ is $\mathbb{R} - \{1\}$
while domain of $g(x)$ is \mathbb{R}

(ii) No, as domain are not same. Domain of $f(x)$ is \mathbb{R}

while that of $g(x)$ is $\mathbb{R} - \left\{ (2n+1)\frac{\pi}{2}; n \in \mathbb{I} \right\}$

Ex. 7 Find the value of $\left\lfloor \frac{1}{2} \right\rfloor + \left\lfloor \frac{1}{2} + \frac{1}{1000} \right\rfloor + \dots + \left\lfloor \frac{1}{2} + \frac{2946}{1000} \right\rfloor$ where $[\cdot]$ denotes greatest integer function ?

Sol. $\left\lfloor \frac{1}{2} \right\rfloor + \left\lfloor \frac{1}{2} + \frac{1}{1000} \right\rfloor + \dots + \left\lfloor \frac{1}{2} + \frac{499}{1000} \right\rfloor + \left\lfloor \frac{1}{2} + \frac{500}{1000} \right\rfloor + \dots + \left\lfloor \frac{1}{2} + \frac{1499}{1000} \right\rfloor + \left\lfloor \frac{1}{2} + \frac{1500}{1000} \right\rfloor + \dots$
 $+ \left\lfloor \frac{1}{2} + \frac{2499}{1000} \right\rfloor + \left\lfloor \frac{1}{2} + \frac{2500}{1000} \right\rfloor + \dots + \left\lfloor \frac{1}{2} + \frac{2946}{1000} \right\rfloor$
 $= 0 + 1 \times 1000 + 2 \times 1000 + 3 \times 447 = 3000 + 1341 = 4341$

Ex. 8 Find the range of $f(x) = \frac{x - [x]}{1 + x - [x]}$, where $[.]$ denotes greatest integer function.

Sol. $y = \frac{x - [x]}{1 + x - [x]} = \frac{\{x\}}{1 + \{x\}}$

$$\therefore \frac{1}{y} = \frac{1}{\{x\}} + 1 \Rightarrow \frac{1}{\{x\}} = \frac{1-y}{y} \Rightarrow \{x\} = \frac{y}{1-y}$$

$$0 \leq \{x\} < 1 \Rightarrow 0 \leq \frac{y}{1-y} < 1$$

$$\text{Range} = [0, 1/2)$$

Ex. 9 Let $f(x) = e^x; \mathbb{R}^+ \rightarrow \mathbb{R}$ and $g(x) = \sin^{-1} x; [-1, 1] \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$. Find domain and range of $\text{fog}(x)$

Sol. Domain of $f(x) : (0, \infty)$ Range of $g(x) : \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

values in range of $g(x)$ which are accepted by $f(x)$ are $\left(0, \frac{\pi}{2}\right]$

$$\Rightarrow 0 < g(x) \leq \frac{\pi}{2} \Rightarrow 0 < \sin^{-1} x \leq \frac{\pi}{2} \Rightarrow 0 < x \leq 1$$

Hence domain of $\text{fog}(x)$ is $x \in (0, 1]$



Ex. 10 Let $A = \{x : -1 \leq x \leq 1\} = B$ be a mapping $f: A \rightarrow B$. For each of the following functions from A to B , find whether it is surjective or bijective.

(A) $f(x) = |x|$ (B) $f(x) = x|x|$ (C) $f(x) = x^3$

(D) $f(x) = [x]$ (E) $f(x) = \sin \frac{\pi x}{2}$

Sol. (A) $f(x) = |x|$


Graphically;

Which shows many one, as the straight line is parallel to x -axis and cuts at two points. Here range for $f(x) \in [0, 1]$

Which is clearly subset of co-domain i.e., $[0, 1] \subseteq [-1, 1]$ Thus, into.

Hence, function is many-one-into

\therefore Neither injective nor surjective




(B) $f(x) = x|x| = \begin{cases} -x^2, & -1 < x < 0 \\ x^2, & 0 \leq x < 1 \end{cases}$,

Graphically,

The graph shows $f(x)$ is one-one, as the straight line parallel to x-axis cuts only at one point.

Here, range



$$f(x) \in [-1, 1]$$

Thus, range = co-domain


Hence, onto.

Therefore, $f(x)$ is one-one onto or (Bijective).

(C) $f(x) = x^3$,

Graphically;

Graph shows $f(x)$ is one-one onto
(i.e. Bijective)




[as explained in above example]

(D) $f(x) = [x]$,

Graphically;

Which shows $f(x)$ is many-one, as the straight line parallel to x-axis meets at more than one point.




Here, range

$$f(x) \in \{-1, 0, 1\}$$

which shows into as range \subseteq co-domain

Hence, many-one-into

- (E) $f(x) = \sin$
Graphically;



Which shows $f(x)$ is one-one and onto as range
= co-domain.

Therefore, $f(x)$ is bijective.

Ex. 11 Composition of piecewise defined functions :

$$\text{If } f(x) = ||x-3|-2| \quad 0 \leq x \leq 4$$

$$g(x) = 4 - |2-x| \quad -1 \leq x \leq 3$$

then find $f \circ g(x)$ and draw rough sketch of $f \circ g(x)$.

Sol. $f(x) = ||x-3|-2| \quad 0 \leq x \leq 4$

$$= \begin{cases} |x-1| & 0 \leq x < 3 \\ |x-5| & 3 \leq x \leq 4 \end{cases} = \begin{cases} 1-x & 0 \leq x < 1 \\ x-1 & 1 \leq x < 3 \\ 5-x & 3 \leq x \leq 4 \end{cases}$$

$$g(x) = 4 - |2-x| \quad -1 \leq x \leq 3$$

$$= \begin{cases} 4-(2-x) & -1 \leq x < 2 \\ 4-(x-2) & 2 \leq x \leq 3 \end{cases} = \begin{cases} 2+x & -1 \leq x < 2 \\ 6-x & 2 \leq x \leq 3 \end{cases}$$

$$\therefore f \circ g(x) = \begin{cases} 1-g(x) & 0 \leq g(x) < 1 \\ g(x)-1 & 1 \leq g(x) < 3 \\ 5-g(x) & 3 \leq g(x) \leq 4 \end{cases} = \begin{cases} 1-(2+x) & 0 \leq 2+x < 1 \text{ and } -1 \leq x < 2 \\ 2+x-1 & 1 \leq 2+x < 3 \text{ and } -1 \leq x < 2 \\ 5-(2+x) & 3 \leq 2+x \leq 4 \text{ and } -1 \leq x < 2 \\ 1-6+x & 0 \leq 6-x < 1 \text{ and } 2 \leq x \leq 3 \\ 6-x-1 & 1 \leq 6-x < 3 \text{ and } 2 \leq x \leq 3 \\ 5-6+x & 3 \leq 6-x \leq 4 \text{ and } 2 \leq x \leq 3 \end{cases}$$

$$= \begin{cases} -1-x & -2 \leq x < -1 \text{ and } -1 \leq x < 2 \\ 1+x & -1 \leq x < 1 \text{ and } -1 \leq x < 2 \\ 3-x & 1 \leq x \leq 2 \text{ and } -1 \leq x < 2 \\ x-5 & -6 \leq -x < -5 \text{ and } 2 \leq x \leq 3 \\ 5-x & -5 \leq -x < -3 \text{ and } 2 \leq x \leq 3 \\ x-1 & -3 \leq -x \leq -2 \text{ and } 2 \leq x \leq 3 \end{cases} = \begin{cases} -1-x & -2 \leq x < -1 \text{ and } -1 \leq x < 2 \\ 1+x & -1 \leq x < 1 \text{ and } -1 \leq x < 2 \\ 3-x & 1 \leq x \leq 2 \text{ and } -1 \leq x < 2 \\ x-5 & 5 < x \leq 6 \text{ and } 2 \leq x \leq 3 \\ 5-x & 3 < x \leq 5 \text{ and } 2 \leq x \leq 3 \\ x-1 & 2 \leq x \leq 3 \text{ and } 2 \leq x \leq 3 \end{cases}$$

$$= \begin{cases} 1+x & -1 \leq x < 1 \\ 3-x & 1 \leq x < 2 \\ x-1 & 2 \leq x \leq 3 \end{cases}$$

Ex. 12 (i) Find whether $f(x) = x + \cos x$ is one-one.

(ii) Identify whether the function $f(x) = -x^3 + 3x^2 - 2x + 4$ for $f: \mathbb{R} \rightarrow \mathbb{R}$ is ONTO or INTO

(iii) $f(x) = x^2 - 2x + 3; [0, 3] \rightarrow A$. Find whether $f(x)$ is injective or not. Also find the set A, if $f(x)$ is surjective.

Sol. (i) The domain of $f(x)$ is \mathbb{R} . $f'(x) = 1 - \sin x$.


$\therefore f'(x) \geq 0 \forall x \in \text{complete domain}$ and equality holds at discrete points only

$\therefore f(x)$ is strictly increasing on \mathbb{R} . Hence $f(x)$ is one-one.

(ii) As range \equiv codomain, therefore given function is ONTO

(iii) $f(x) = 2(x-1); 0 \leq x \leq 3$

$$\therefore f'(x) = \begin{cases} -ve & ; 0 \leq x < 1 \\ +ve & ; 1 < x \leq 3 \end{cases}$$



$\therefore f(x)$ is non monotonic. Hence it is not injective.

For $f(x)$ to be surjective, A should be equal to its range. By graph range is $[2, 6]$

$\therefore A \equiv [2, 6]$

Ex. 13 If f be the greatest integer function and g be the modulus function, then $(g \circ f)\left(-\frac{5}{3}\right) - (f \circ g)\left(-\frac{5}{3}\right) =$

(A) 1

(B) -1

(C) 2

(D) 4

Sol. Given $(g \circ f)\left(-\frac{5}{3}\right) - (f \circ g)\left(-\frac{5}{3}\right) = g\left\{f\left(-\frac{5}{3}\right)\right\} - f\left\{g\left(-\frac{5}{3}\right)\right\} = g(-2) - f\left(\frac{5}{3}\right) = 2 - 1 = 1$ **Ans.(A)**

Ex. 14 Show that $\log(x + \sqrt{x^2 + 1})$ is an odd function.

Sol. Let $f(x) = \log(x + \sqrt{x^2 + 1})$.

$$\text{Then } f(-x) = \log(-x + \sqrt{(-x)^2 + 1})$$

$$= \log\left(\frac{(\sqrt{x^2 + 1} - x)(\sqrt{x^2 + 1} + x)}{\sqrt{x^2 + 1} + x}\right) = \log\frac{1}{\sqrt{x^2 + 1} + x} = -\log(x + \sqrt{x^2 + 1}) = -f(x)$$

$$\text{or } f(x) + f(-x) = 0$$

Hence $f(x)$ is an odd function.

Ex. 15 Show that $\cos^{-1} x$ is neither odd nor even.

Sol. Let $f(x) = \cos^{-1} x$. Then $f(-x) = \cos^{-1}(-x) = \pi - \cos^{-1} x$ which is neither equal to $f(x)$ nor equal to $-f(x)$.

Hence $\cos^{-1} x$ is neither odd nor even

Ex. 161 Which of the following functions is (are) even, odd or neither :

(i) $f(x) = x^2 \sin x$

(ii) $f(x) = \sqrt{1+x+x^2} - \sqrt{1-x+x^2}$

(iii) $f(x) = \log\left(\frac{1-x}{1+x}\right)$

(iv) $f(x) = \sin x - \cos x$

(v) $f(x) = \frac{e^x + e^{-x}}{2}$

Sol. (i) $f(-x) = (-x)^2 \sin(-x) = -x^2 \sin x = -f(x).$

Hence $f(x)$ is odd.

(ii) $f(-x) = \sqrt{1+(-x)+(-x)^2} - \sqrt{1-(-x)+(-x)^2}$
 $= \sqrt{1-x+x^2} - \sqrt{1+x+x^2} = -f(x).$

Hence $f(x)$ is odd.

(iii) $f(-x) = \log\left(\frac{1-(-x)}{1+(-x)}\right) = \log\left(\frac{1+x}{1-x}\right) = -f(x).$

Hence $f(x)$ is odd

(iv) $f(-x) = \sin(-x) - \cos(-x) = -\sin x - \cos x.$

Hence $f(x)$ is neither even nor odd.

(v) $f(-x) = \frac{e^{-x} + e^{-(-x)}}{2} = \frac{e^{-x} + e^x}{2} = f(x).$

Hence $f(x)$ is even

Ex. 17 Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = (e^x - e^{-x})/2$. Is $f(x)$ invertible? If so, find its inverse.

Sol. Let us check for invertibility of $f(x)$:

(A) **One-One :**

Let $x_1, x_2 \in \mathbb{R}$ and $x_1 < x_2$

$\Rightarrow e^{x_1} < e^{x_2}$ (Because base $e > 1$)(i)

Also $x_1 < x_2 \Rightarrow -x_2 < -x_1$

$\Rightarrow e^{-x_2} < e^{-x_1}$ (Because base $e > 1$)(ii)

(i) + (ii) $\Rightarrow e^{x_1} + e^{-x_2} < e^{x_2} + e^{-x_1}$

$\Rightarrow \frac{1}{2}(e^{x_1} - e^{-x_1}) < \frac{1}{2}(e^{x_2} - e^{-x_2}) \Rightarrow f(x_1) < f(x_2)$ i.e. f is one-one.

(B) **Onto :**

As x tends to larger and larger values so does $f(x)$ and

when $x \rightarrow \infty, f(x) \rightarrow \infty$.

Similarly as $x \rightarrow -\infty, f(x) \rightarrow -\infty$ i.e. $-\infty < f(x) < \infty$ so long as $x \in (-\infty, \infty)$

Hence the range of f is same as the set \mathbb{R} . Therefore $f(x)$ is onto.

Since $f(x)$ is both one-one and onto, $f(x)$ is invertible.

(C) To find f^{-1} :

Let f^{-1} be the inverse function of f , then by rule of identity $\text{fof}^{-1}(x) = x$

$$\frac{e^{f^{-1}(x)} - e^{-f^{-1}(x)}}{2} = x \Rightarrow e^{2f^{-1}(x)} - 2xe^{f^{-1}(x)} - 1 = 0$$

$$\Rightarrow e^{f^{-1}(x)} = \frac{2x \pm \sqrt{4x^2 + 4}}{2} \Rightarrow e^{f^{-1}(x)} = x \pm \sqrt{1+x^2}$$

Since $e^{f^{-1}(x)} > 0$, hence negative sign is ruled out and

$$\text{Hence } e^{f^{-1}(x)} = x + \sqrt{1+x^2}$$

Taking logarithm, we have $f^{-1}(x) = \ln(x + \sqrt{1+x^2})$.

Ex. 18 Find the periods (if periodic) of the following functions, where $[.]$ denotes the greatest integer function

(i) $f(x) = e^{\bullet n(\sin x)} + \tan^3 x - \text{cosec}(3x - 5)$ (ii) $f(x) = x - [x - b]$, $b \in \mathbb{R}$

(iii) $f(x) = \frac{|\sin x + \cos x|}{|\sin x| + |\cos x|}$ (iv) $f(x) = \tan \frac{\pi}{2} [x]$

(v) $f(x) = \cos(\sin x) + \cos(\cos x)$ (vi) $f(x) = \frac{(1 + \sin x)(1 + \sec x)}{(1 + \cos x)(1 + \csc x)}$

(vii) $f(x) = e^{x - [x] + \cos \pi x + \cos 2\pi x + \dots + \cos n\pi x}$

Sol.(i) $f(x) = e^{\bullet n(\sin x)} + \tan^3 x - \text{cosec}(3x - 5)$

Period of $e^{\bullet n \sin x} = 2\pi$, $\tan^3 x = \pi$

$$\text{cosec}(3x - 5) = \frac{2\pi}{3}$$

$$\therefore \text{Period} = 2\pi$$

(ii) $f(x) = x - [x - b] = b + \{x - b\}$

$$\therefore \text{Period} = 1$$

(iii) $f(x) = \frac{|\sin x + \cos x|}{|\sin x| + |\cos x|}$

Since period of $|\sin x + \cos x| = \pi$ and period of $|\sin x| + |\cos x|$ is $\frac{\pi}{2}$. Hence $f(x)$ is periodic with π as its period

(iv) $f(x) = \tan \frac{\pi}{2} [x]$

$$\tan \frac{\pi}{2} [x + T] = \tan \frac{\pi}{2} [x] \Rightarrow \frac{\pi}{2} [x + T] = n\pi + \frac{\pi}{2} [x]$$

$$\therefore T = 2$$

$$\therefore \text{Period} = 2$$

(v) Let $f(x)$ is periodic then $f(x + T) = f(x)$

$$\Rightarrow \cos(\sin(x + T)) + \cos(\cos(x + T)) = \cos(\sin x) + \cos(\cos x)$$

$$\text{If } x = 0 \text{ then } \cos(\sin T) + \cos(\cos T) = \cos(0) + \cos(1) = \cos\left(\cos\frac{\pi}{2}\right) + \cos\left(\sin\frac{\pi}{2}\right)$$

$$\text{On comparing } T = \frac{\pi}{2}$$

$$(vi) \quad f(x) = \frac{(1 + \sin x)(1 + \sec x)}{(1 + \cos x)(1 + \csc x)} = \frac{(1 + \sin x)(1 + \sec x)}{(1 + \cos x)(1 + \csc x)}$$

$$\Rightarrow f(x) = \tan x$$

Hence $f(x)$ has period π .

$$(vii) \quad f(x) = e^{x - [x] + |\cos \pi x| + |\cos 2\pi x| + \dots + |\cos n\pi x|}$$

$$\text{Period of } x - [x] = 1$$

$$\text{Period of } |\cos \pi x| = 1$$

$$\text{Period of } |\cos 2\pi x| = \frac{1}{2}$$

.....

$$\text{Period of } |\cos n\pi x| = \frac{1}{n}$$

So period of $f(x)$ will be L.C.M. of all period = 1

Ex. 19 Find the periods (if periodic) of the following functions, where $[.]$ denotes the greatest integer function

$$(i) \quad f(x) = e^{x - [x]} + \sin x$$

$$(ii) \quad f(x) = \sin \frac{\pi x}{\sqrt{2}} + \cos \frac{\pi x}{\sqrt{3}}$$

$$(iii) \quad f(x) = \sin \frac{\pi x}{\sqrt{3}} + \cos \frac{\pi x}{2\sqrt{3}}$$

Sol.(i) Period of $e^{x - [x]} = 1$

$$\text{period of } \sin x = 2\pi$$

→ L.C.M. of rational and an irrational number does not exist.

∴ not periodic.

$$(ii) \quad \text{Period of } \sin \frac{\pi x}{\sqrt{2}} = \frac{2\pi}{\pi/\sqrt{2}} = 2\sqrt{2}$$

$$\text{Period of } \cos \frac{\pi x}{\sqrt{3}} = \frac{2\pi}{\pi/\sqrt{3}} = 2\sqrt{3}$$

→ L.C.M. of two different kinds of irrational number does not exist.

∴ not periodic.

(iii) Period of $\sin \frac{\pi x}{\sqrt{3}} = \frac{2\pi}{\pi/\sqrt{3}} = 2\sqrt{3}$


Period of $\cos \frac{\pi x}{2\sqrt{3}} = \frac{2\pi}{\pi/2\sqrt{3}} = 4\sqrt{3}$

→ L.C.M. of two similar irrational number exist.

∴ Periodic with period = $4\sqrt{3}$ **Ans.**

20.(i) Let $f(x) = x^2 + 2x$; $x \geq -1$. Draw graph of $f^{-1}(x)$ also find the number of solutions of the equation, $f(x) = f^{-1}(x)$

(ii) If $y = f(x) = x^2 - 3x + 1$, $x \geq 2$. Find the value of $g'(1)$ where g is inverse of f



$f(x) = f^{-1}(x)$ is equivalent to $f(x) = x$

$$\Rightarrow x^2 + 2x = x \Rightarrow x(x+1) = 0 \Rightarrow x = 0, -1$$

Hence two solution for $f(x) = f^{-1}(x)$

(iv) $y = 1$

$$\Rightarrow x^2 - 3x + 1 = 1$$

$$\Rightarrow x(x-3) = 0 \Rightarrow x = 0, 3$$

But $x \geq 2$ ∴ $x = 3$

Now $g(f(x)) = x$


Differentiating both sides w.r.t. x

$$\Rightarrow g'(f(x)) \cdot f'(x) = 1 \Rightarrow g'(f(x)) = \frac{1}{f'(x)}$$

$$\Rightarrow g'(f(3)) = \frac{1}{f'(3)} \Rightarrow g'(1) = \frac{1}{6-3} = \frac{1}{3} \quad (\text{As } f'(x) = 2x - 3)$$

Ex.21 Find $f(x) = \max \{1+x, 1-x, 2\}$.


Sol. From the graph it is clear that



$$f(x) = \begin{cases} 1-x & ; x < -1 \\ 2 & ; -1 \leq x \leq 1 \\ 1+x & ; x > 1 \end{cases}$$

Ex.22 Draw the graph of $y = |2 - |x - 1||$.

Sol.




Ex.23 Draw the graph of $f(x) = \cos x \cos(x+2) - \cos^2(x+1)$.

Sol. $f(x) = \cos x \cos(x+2) - \cos^2(x+1)$

$$= \frac{1}{2} [\cos(2x+2) + \cos 2] - \frac{1}{2} [\cos(2x+2) + 1]$$

$$= \frac{1}{2} \cos 2 - \frac{1}{2} < 0$$



Exercise # 1

[Single Correct Choice Type Questions]

- The domain of $f(x) = \sqrt{\frac{1-|x|}{2-|x|}}$, is -
 (A) $(-\infty, \infty) - [-2, 2]$ (B) $(-\infty, \infty) - [-1, 1]$
 (C) $[-1, 1] \cup (-\infty, -2) \cup (2, \infty)$ (D) none
- The domain of the function $f(x) = \sin^{-1}\left(\frac{1+x^3}{2x^{3/2}}\right) + \sqrt{\sin(\sin x)} + \log_{(3\{x\}+1)}(x^2+1)$, where $\{.\}$ represents fractional part function, is:
 (A) $x \in \{1\}$ (B) $x \in \mathbb{R} - \{1, -1\}$ (C) $x > 3, x \neq 1$ (D) none of these
- The domain of the function $f(x) = \frac{1}{\log_{10}(1-x)} + \sqrt{x+2}$, is -
 (A) $[-2, 0) \cup (0, 1)$ (B) $(-2, 0) \cup (0, 1]$ (C) $(-2, 0) \cup (0, 1]$ (D) $(-2, 0) \cup [0, 1]$
- If $q^2 - 4pr = 0, p > 0$, then the domain of the function $f(x) = \log(px^3 + (p+q)x^2 + (q+r)x + r)$ is:
 (A) $\mathbb{R} - \left\{-\frac{q}{2p}\right\}$ (B) $\mathbb{R} - \left[(-\infty, -1] \cup \left\{-\frac{q}{2p}\right\}\right]$
 (C) $\mathbb{R} - \left[(-\infty, -1) \cap \left\{-\frac{q}{2p}\right\}\right]$ (D) none of these
- If $f(x)$ is a polynomial function satisfying the condition $f(x) \cdot f(1/x) = f(x) + f(1/x)$ and $f(2) = 9$ then -
 (A) $2f(4) = 3f(6)$ (B) $14f(1) = f(3)$ (C) $9f(3) = f(5)$ (D) $f(10) = f(11)$
- Domain to function $\sqrt{\log\{(5x-x^2)/6\}}$ is -
 (A) $(2, 3)$ (B) $[2, 3]$ (C) $[1, 2]$ (D) $[1, 3]$
- Domain and range of $f(x) = \sqrt{x-1} + 2\sqrt{3-x}$ is
 (A) $D : [1, 3]; R : [\sqrt{2}, \sqrt{10}]$ (B) $D : [1, 5]; R : [\sqrt{2}, \sqrt{10}]$
 (C) $D : (-\infty, 1] \cup [3, \infty); R : [1, \sqrt{3}]$ (D) $D : [1, 5]; R : [1, \sqrt{3}]$
- If $A = \{-2, -1, 0, 1, 2\}$ & $f : A \rightarrow \mathbb{Z}; f(x) = x^2 + 1$, then the range of f is
 (A) $\{0, 1, 2, 5\}$ (B) $\{1, 2, 5\}$ (C) $\{-5, -2, 1, 2, 3\}$ (D) A

9. The greatest value of the function $f(x) = (\sin^{-1} x)^3 + (\cos^{-1} x)^3$ is:
- (A) $\frac{\pi^3}{32}$ (B) $\frac{\pi^3}{8}$ (C) $\frac{3\pi^3}{8}$ (D) $\frac{7\pi^3}{8}$
10. The range of the function $f(x) = e^x - e^{-x}$, is -
- (A) $[0, \infty)$ (B) $(-\infty, 0)$ (C) $(-\infty, \infty)$ (D) none
11. The range of the function $f(x) = {}^{7-x}P_{x-3}$, is -
- (A) $\{1, 2, 3\}$ (B) $\{1, 2, 3, 4, 5, 6\}$ (C) $\{1, 2, 3, 4\}$ (D) $\{1, 2, 3, 4, 5\}$
12. If $f(x) = 2[x] + \cos x$, then $f: \mathbb{R} \rightarrow \mathbb{R}$ is: (where $[\]$ denotes greatest integer function)
- (A) one-one and onto (B) one-one and into
(C) many-one and into (D) many-one and onto
13. $f: [-1, 1] \rightarrow [-1, 2]$, $f(x) = x + |x|$, is -
- (A) one-one onto (B) one-one into (C) many one onto (D) many one into
14. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $f(0) = 1$ and for any $x, y \in \mathbb{R}$, $f(xy + 1) = f(x)f(y) - f(y) - x + 2$. Then f is
- (A) one-one and onto (B) one-one but not onto (C) many one but onto (D) many one and into
15. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by $f(x) = \frac{x^2 - 3x + 4}{x^2 + 3x + 4}$ then f is -
- (A) one - one but not onto (B) onto but not one - one
(C) onto as well as one - one (D) neither onto nor one - one
16. Which one of the following pair of functions are identical ?
- (A) $e^{(\bullet nx)/2}$ and \sqrt{x}
(B) $\tan^{-1}(\tan x)$ and $\cot^{-1}(\cot x)$
(C) $\cos^2 x + \sin^4 x$ and $\sin^2 x + \cos^4 x$
(D) $a \frac{|x|}{x}$ and $\text{sgn}(x)$, where $\text{sgn}(x)$ stands for signum function.
17. If $f(x) = \cos \left[\frac{1}{2} \pi^2 \right] x + \sin x \left[\frac{1}{2} \pi^2 \right]$, $[x]$ denoting the greatest integer function, then -
- (A) $f(0) = 0$ (B) $f\left(\frac{\pi}{3}\right) = \frac{1}{4}$ (C) $f\left(\frac{\pi}{2}\right) = 1$ (D) $f(\pi) = 0$
18. If $f(x) = \cos(\log x)$, then $f(x)f(y) - \frac{1}{2}[f(x/y) + f(xy)]$ is equal to -
- (A) -1 (B) 1/2 (C) -2 (D) 0
19. The value of b and c for which the identity $f(x+1) - f(x) = 8x + 3$ is satisfied, where $f(x) = bx^2 + cx + d$, are -
- (A) $b = 2, c = 1$ (B) $b = 4, c = -1$ (C) $b = -1, c = 4$ (D) $b = -1, c = 1$

MATHS FOR JEE MAIN & ADVANCED

20. If $f(x) = \frac{4a-7}{3}x^3 + (a-3)x^2 + x + 5$ is a one-one function, then
 (A) $2 \leq a \leq 8$ (B) $1 \leq a \leq 2$ (C) $0 \leq a \leq 1$ (D) None of these
21. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by then -
 (A) f is a bijection (B) f is an injection only
 (C) f is a surjection (D) f is neither injection nor a surjection
22. If $f(x) = \{x\} + \{x+1\} + \{x+2\} \dots \{x+99\}$, then the value of $[f(\sqrt{2})]$ is, where $\{.\}$ denotes fractional part function & $[.]$ denotes the greatest integer function
 (A) 5050 (B) 4950 (C) 41 (D) 14
23. The minimum value of $f(x) = |3-x| + |2+x| + |5-x|$ is -
 (A) 0 (B) 7 (C) 8 (D) 10
24. If the function $f : \mathbb{R} \rightarrow A$ given by $f(x) = \frac{x^2}{x^2+1}$ is a surjection, then $A =$
 (A) \mathbb{R} (B) $[0, 1]$ (C) $(0, 1]$ (D) $[0, 1)$
25. The fundamental period of function $f(x) = [x] + \left[x + \frac{1}{3}\right] + \left[x + \frac{2}{3}\right] - 3x + 15$, where $[.]$ denotes greatest integer function, is :
 (A) $\frac{1}{3}$ (B) $\frac{2}{3}$ (C) 1 (D) non-periodic
26. $f(x) = |x-1|$, $f: \mathbb{R}^+ \rightarrow \mathbb{R}$, $g(x) = e^x$, $g: [-1, \infty) \rightarrow \mathbb{R}$. If the function $\text{fog}(x)$ is defined, then its domain and range respectively are:
 (A) $(0, \infty)$ and $[0, \infty)$ (B) $[-1, \infty)$ and $[0, \infty)$
 (C) $[-1, \infty)$ and $\left[1 - \frac{1}{e}, \infty\right)$ (D) $[-1, \infty)$ and $\left[\frac{1}{e} - 1, \infty\right)$
27. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by $f(x) = \frac{e^{|x|} - e^{-x}}{e^x + e^{-x}}$ then -
 (A) f is a bijection (B) f is an injection only
 (C) f is a surjection (D) f is neither injection nor a surjection
28. Let $f: (2, 4) \rightarrow (1, 3)$ be a function defined by $f(x) = x - \left[\frac{x}{2}\right]$ (where $[.]$ denotes the greatest integer function), then $f^{-1}(x)$ is equal to :
 (A) $2x$ (B) $x + \left[\frac{x}{2}\right]$ (C) $x+1$ (D) $x-1$


29. The mapping $f: \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x^3 + ax^2 + bx + c$ is a bijection if
 (A) $b^2 \leq 3a$ (B) $a^2 \leq 3b$ (C) $a^2 \geq 3b$ (D) $b^2 \geq 3a$
30. The period of the function $f(x) = \sin\left(\cos\frac{x}{2}\right) + \cos(\sin x)$ equal -
 (A) $\frac{\pi}{2}$ (B) 2π (C) π (D) 4π
31. Let $f(x) = \sin\sqrt{[a]} x$ (where $[]$ denotes the greatest integer function). If f is periodic with fundamental period π , then a belongs to -
 (A) $[2, 3)$ (B) $\{4, 5\}$ (C) $[4, 5]$ (D) $[4, 5)$
32. Which of the following function has a period of 2π ?
 (A) $f(x) = \sin\left(2\pi x + \frac{\pi}{3}\right) + 2\sin\left(3\pi x + \frac{\pi}{4}\right) + 3\sin 5\pi x$ (B) $f(x) = \sin \frac{\pi x}{3} + \sin \frac{\pi x}{4}$
 (C) $f(x) = \sin x + \cos 2x$ (D) none
33. A function whose graph is symmetrical about the origin is given by -
 (A) $f(x) = e^x + e^{-x}$ (B) $f(x) = \sin(\sin(\cos(\sin x)))$
 (C) $f(x+y) = f(x) + f(y)$ (D) $\sin x + \sin|x|$
34. If $f: \mathbb{R} \rightarrow \mathbb{R}$ is a function satisfying the property $f(x+1) + f(x+3) =$ then the period of $f(x)$ is -
 (A) 4 (B) K (C) 1 (D) π
35. If $f(x) = 3x - 5$, then $f^{-1}(x)$ -
 (A) is given by $\frac{1}{3x-5}$ (B) is given by $\frac{x+5}{3}$
 (C) does not exist because f is not one-one (D) does not exist because f is not onto
36. If the function $f: [1, \infty) \rightarrow [1, \infty)$ is defined by $f(x) = 2^{x(x-1)}$, then $f^{-1}(x)$ is -
 (A) $\left(\frac{1}{2}\right)^{x(x-1)}$ (B) $\frac{1}{2}\left(1 + \sqrt{1 + 4\log_2 x}\right)$ (C) $\frac{1}{2}\left(1 - \sqrt{1 + 4\log_2 x}\right)$ (D) Not defined

Exercise # 2


Part # I

[Multiple Correct Choice Type Questions]


- Which of the functions defined below are NOT one-one function(s) ?
 (A) $f(x) = 5(x^2 + 4), (x \in \mathbb{R})$
 (B) $g(x) = 2x + (1/x)$
 (C) $h(x) = \ln(x^2 + x + 1), (x \in \mathbb{R})$
 (D) $f(x) = e^{-x}$
- Which of the following functions from \mathbb{Z} to itself are NOT bijections ?
 (A) $f(x) = x^3$
 (B) $f(x) = x + 2$
 (C) $f(x) = 2x + 1$
 (D) $f(x) = x^2 + x$
- If $f(x) = \sin^{-1} \left(\frac{\sqrt{4-x^2}}{1-x} \right)$, then
 (A) domain of $f(x)$ is $(-2, 1)$
 (B) domain of $f(x)$ is $[-1, 1]$
 (C) range of $f(x)$ is $[-1, 1]$
 (D) range of $f(x)$ is $[-1, 1)$
- The function $\cot(\sin x)$ -
 (A) is not defined for $x = (4n + 1) \frac{\pi}{2}$
 (B) is not defined for $x = n\pi$
 (C) lies between $-\cot 1$ and $\cot 1$
 (D) can't lie between $-\cot 1$ and $\cot 1$
- The graph of function $f(x)$ is as shown, adjacently. Then the graph of $\frac{1}{f(|x|)}$ is -




(A)




(B)



(C)



(D)



6. Which of the following function(s) is/are periodic ?
 (A) $f(x) = 3x - [3x]$ (B) $g(x) = \sin(1/x), x \neq 0 \text{ \& } g(0) = 0$
 (C) $h(x) = x \cos x$ (D) $w(x) = \sin(\sin(x))$
7. The fundamental period of $\frac{|\sin x| + |\cos x|}{|\sin x - \cos x| + |\sin x + \cos x|}$ is -
 (A) π (B) $\frac{\pi}{2}$ (C) 2π (D) $\frac{2\pi}{3}$
8. The range of the function $f(x) = \sin \left[\log \left(\frac{\sqrt{4-x^2}}{1-x} \right) \right]$ is -
 (A) $[-1, 1]$ (B) $(-1, 1)$ (C) $[-1, 1)$ (D) cannot be determined
9. If $F(x) = \frac{\sin \pi [x]}{\{x\}}$, then $F(x)$ is: (where $\{ \cdot \}$ denotes fractional part function and $[\cdot]$ denotes greatest integer function and $\text{sgn}(x)$ is a signum function)
 (A) periodic with fundamental period 1 (B) even
 (C) range is singleton (D) identical to $\text{sgn} \left(\frac{\{x\}}{\sqrt{\{x\}}} \right) - 1$
10. In the following functions defined from $[-1, 1]$ to $[-1, 1]$, then functions which are not bijective are
 (A) $\sin(\sin^{-1}x)$ (B) $\frac{2}{\pi} \sin^{-1}(\sin x)$ (C) $(\text{sgn } x) \bullet n e^x$ (D) $x^3 \text{sgn } x$
11. Let $f: [-1, 1] \rightarrow [0, 2]$ be a linear function which is onto, then $f(x)$ is/are
 (A) $1-x$ (B) $1+x$ (C) $x-1$ (D) $x+2$
12. Which of the following functions are not homogeneous ?
 (A) $x + y \cos \frac{y}{x}$ (B) $\frac{xy}{x+y^2}$ (C) $\frac{x-y \cos x}{y \sin x + y}$ (D) $\frac{x}{y} \bullet n \left(\frac{y}{x} \right) + \frac{y}{x} \bullet n \left(\frac{x}{y} \right)$
13. Given the function $f(x) 2f(x) + xf\left(\frac{1}{x}\right) - 2f\left(\sqrt{2} \sin \pi \left(x + \frac{1}{4}\right)\right) = 4 \cos^2 \frac{\pi x}{2} + x \cos \frac{\pi}{x}$ such that , then which one of the following is correct ?
 (A) $f(2) + f(1/2) = 1$ (B) $f(1) = -1$, but the values of $f(2), f(1/2)$ cannot be determined
 (C) $f(2) + f(1) = f(1/2)$ (D) $f(2) + f(1) = 0$
14. The function $f(x) = \sqrt{\log_{x^2}(x)}$ is defined for x belonging to -
 (A) $(-\infty, 0)$ (B) $(0, 1)$ (C) $(1, \infty)$ (D) $(0, \infty)$

MATHS FOR JEE MAIN & ADVANCED

15. If $f(x + ay, x - ay) = axy$ then $f(x, y)$ is equal to -
(A) $\frac{x^2 - y^2}{4}$ (B) $\frac{x^2 + y^2}{4}$ (C) $4xy$ (D) none
16. Which of following pairs of functions are identical.
(A) $f(x) = e^{\lambda n \sec^{-1} x}$ and $g(x) = \sec^{-1} x$ (B) $f(x) = \tan(\tan^{-1} x)$ and $g(x) = \cot(\cot^{-1} x)$
(C) $f(x) = \operatorname{sgn}(x)$ and $g(x) = \operatorname{sgn}(\operatorname{sgn}(x))$ (D) $f(x) = \cot^2 x \cdot \cos^2 x$ and $g(x) = \cot^2 x - \cos^2 x$
17. Let $f(x) = \left(\frac{1-x}{1+x}\right)$, $0 \leq x \leq 1$ and $g(x) = 4x(1-x)$, $0 \leq x \leq 1$, then
(A) $f \circ g = \frac{1-4x+4x^2}{1+4x-4x^2}$, $0 \leq x \leq 1$ (B) $f \circ g = \frac{1-4x-4x^2}{1+4x-4x^2}$, $\frac{1}{2} \leq x \leq 1$
(C) $g \circ f = \frac{8x(1-x)}{(1+x)^2}$, $0 \leq x \leq 1$ (D) $g \circ f = \frac{8x(1+x)}{(1+x)^2}$, $0 \leq x \leq 1$
18. Function $f(x) = \sin x + \tan x + \operatorname{sgn}(x^2 - 6x + 10)$ is
(A) periodic with period 2π (B) periodic with period π
(C) Non-periodic (D) periodic with period 4π
19. Which of the functions are even -
(A) $\log\left(\frac{1+x^2}{1-x^2}\right)$ (B) $\sin^2 x + \cos^2 x$ (C) $\log\left(\frac{1+x^3}{1-x^3}\right)$ (D) $\frac{(1+2^x)^2}{2^x}$
20. Let $D \equiv [-1, 1]$ is the domain of the following functions, state which of them are injective.
(A) $f(x) = x^2$ (B) $g(x) = x^3$ (C) $h(x) = \sin 2x$ (D) $k(x) = \sin(\pi x/2)$
21. The period of the function $f(x) = \sin^4 3x + \cos^4 3x$ is:
(A) $\pi/6$ (B) $\pi/3$ (C) $\pi/2$ (D) $\pi/12$
22. Which of the following functions are aperiodic (where $[.]$ denotes greatest integer function)
(A) $y = [x + 1]$ (B) $y = \sin x^2$ (C) $y = \sin^2 x$ (D) $y = \sin^{-1} x$
23. If $f: \mathbb{R} \rightarrow [-1, 1]$, where $f(x) = \sin\left(\frac{\pi}{2}[x]\right)$, (where $[.]$ denotes the greatest integer function), then
(A) $f(x)$ is onto (B) $f(x)$ is into (C) $f(x)$ is periodic (D) $f(x)$ is many one
24. Identify the statement(s) which is/are incorrect ?
(A) the function $f(x) = \cos(\cos^{-1} x)$ is neither odd nor even
(B) the fundamental period of $f(x) = \cos(\sin x) + \cos(\cos x)$ is π
(C) the range of the function $f(x) = \cos(3 \sin x)$ is $[-1, 1]$
(D) none of these

Part # II

[Assertion & Reason Type Questions]

These questions contains, Statement I (assertion) and Statement II (reason).

- (A) Statement-I is true, Statement-II is true ; Statement-II is correct explanation for Statement-I.
 (B) Statement-I is true, Statement-II is true ; Statement-II is NOT a correct explanation for statement-I.
 (C) Statement-I is true, Statement-II is false.
 (D) Statement-I is false, Statement-II is true.

- Statement-I :** Fundamental period of $\cos x + \cot x$ is 2π .
Statement-II : If the period of $f(x)$ is T_1 and the period of $g(x)$ is T_2 , then the fundamental period of $f(x) + g(x)$ is the L.C.M. of T_1 and T_2 .
- Statement - I** If $y = f(x)$ is increasing in $[\alpha, \beta]$, then its range is $[f(\alpha), f(\beta)]$
Statement - II Every increasing function need not to be continuous.
- Statement-I :** Function $f(x) = \sin(x + 3\sin x)$ is periodic.
Statement-II : If $g(x)$ is periodic, then $f(g(x))$ may or may not be periodic.
- Statement : I :** All points of intersection of $y = f(x)$ and $y = f^{-1}(x)$ lies on $y = x$ only.
Statement : II : If point $P(\alpha, \beta)$ lies on $y = f(x)$, then $Q(\beta, \alpha)$ lies on $y = f^{-1}(x)$.
- Let function $f : \mathbb{R} \rightarrow \mathbb{R}$ is such that $f(x)f(y) - f(xy) = x + y$ for all $x, y \in \mathbb{R}$
Statement-I : $f(x)$ is a Bijective function.
Statement-II : $f(x)$ is a linear function.

Exercise # 3

Part # I

[Matrix Match Type Questions]

Following questions contains statements given in two columns, which have to be matched. The statements in **Column-I** are labelled as A, B, C and D while the statements in **Column-II** are labelled as p, q, r and s. Any given statement in **Column-I** can have correct matching with one statement in **Column-II**.

1. Let $f(x) = \sin^{-1} x$, $g(x) = \cos^{-1} x$ and $h(x) = \tan^{-1} x$. For what complete interval of variation of x the following are true.

Column - I

(A) $f(\sqrt{x}) + g(\sqrt{x}) = \pi/2$

(B) $f(x) + g(\sqrt{1-x^2}) = 0$

(C) $g\left(\frac{1-x^2}{1+x^2}\right) = 2h(x)$

(D) $h(x) + h(1) = h\left(\frac{1+x}{1-x}\right)$

Column - II

(p) $[0, \infty)$

(q) $[0, 1]$

(r) $(-\infty, 1)$

(s) $[-1, 0]$

2. **Column - I**

(A) Total number of solution $x^2 - 4 - [x] = 0$ where $[]$ denotes greatest integer function.

(B) Minimum period of $e^{\cos^4 \pi x + \cos^2 \pi x + x - [x]}$

(C) If $A = \{(x, y); y = \frac{1}{x}, x \in \mathbb{R}_0\}$ and $B = \{(x, y) : y = x, x \in \mathbb{R}\}$ then number of elements in $A \cap B$ is (are)

(D) Number of integers in the domain of

$$\sqrt{2^x - 3^x} + \log_3 \log_{1/2} x$$

Column - II

(p) 0

(q) 1

(r) 2

(s) 3

3. **Column - I**

(A) The period of the function $y = \sin(2\pi t + \pi/3) + 2 \sin(3\pi t + \pi/4) + 3 \sin 5\pi t$ is

(B) $y = \{\sin(\pi x)\}$ is a many one function for $x \in (0, a)$, where $\{x\}$ denotes fractional part of x , then a may be

(C) The fundamental period of the function

$$y = \frac{1}{2} \left(\frac{|\sin(\pi/4)x|}{\cos(\pi/4)x} + \frac{\sin(\pi/4)x}{|\cos(\pi/4)x|} \right) \text{ is}$$

(D) If $f: [0, 2] \rightarrow [0, 2]$ is bijective function defined by $f(x) = ax^2 + bx + c$, where a, b, c are non-zero real numbers, then $f(2)$ is equal to

Column - II

(p) $1/2$

(q) 8

(r) 2

(s) 0

4.

Column - I

- (A) $f: \mathbb{R} \rightarrow \mathbb{R}$
 $f(x) = (x-1)(x-2)\dots(x-11)$
- (B) $f: \mathbb{R} - \{-4/3\} \rightarrow \mathbb{R}$
 $f(x) = \frac{2x+1}{3x+4}$
- (C) $f: \mathbb{R} \rightarrow \mathbb{R}$
 $f(x) = e^{\sin x} + e^{-\sin x}$
- (D) $f: \mathbb{R} \rightarrow \mathbb{R}$
 $f(x) = \log(x^2 + 2x + 3)$

Column - II

- (p) one one
- (q) onto
- (r) many one
- (s) into

Part # II

[Comprehension Type Questions]

Comprehension # 1

Given a function $f: A \rightarrow B$; where $A = \{1, 2, 3, 4, 5\}$ and $B = \{6, 7, 8\}$

- Find number of all such functions $y = f(x)$ which are one-one ?
 (A) 0 (B) 3^5 (C) 5P_3 (D) 5^3
- Find number of all such functions $y = f(x)$ which are onto
 (A) 243 (B) 93 (C) 150 (D) none of these
- The number of mappings of $g(x): B \rightarrow A$ such that $g(i) \leq g(j)$ whenever $i < j$ is
 (A) 60 (B) 140 (C) 10 (D) 35

Comprehension # 2

If $f(x) = \begin{cases} x+1, & \text{if } x \leq 1 \\ 5-x^2, & \text{if } x > 1 \end{cases}$ & $g(x) = \begin{cases} x, & \text{if } x \leq 1 \\ 2-x, & \text{if } x > 1 \end{cases}$

On the basis of above information, answer the following questions :

- The range of $f(x)$ is -
 (A) $(-\infty, 4)$ (B) $(-\infty, 5)$ (C) \mathbb{R} (D) $(-\infty, 4]$
- If $x \in (1, 2)$, then $g(f(x))$ is equal to -
 (A) $x^2 + 3$ (B) $x^2 - 3$ (C) $5 - x^2$ (D) $1 - x$
- Number of negative integral solutions of $g(f(x)) + 2 = 0$ are -
 (A) 0 (B) 3 (C) 1 (D) 2

Comprehension # 3

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ is a function satisfying $f(2-x) = f(2+x)$ and $f(20-x) = f(x)$, $\forall x \in \mathbb{R}$.

On the basis of above information, answer the following questions :

- If $f(0) = 5$, then minimum possible number of values of x satisfying $f(x) = 5$, for $x \in [0, 170]$ is-
 (A) 21 (B) 12 (C) 11 (D) 22
- Graph of $y = f(x)$ is -
 (A) symmetrical about $x = 18$ (B) symmetrical about $x = 5$
 (C) symmetrical about $x = 8$ (D) symmetrical about $x = 20$
- If $f(2) \neq f(6)$, then
 (A) fundamental period of $f(x)$ is 1 (B) fundamental period of $f(x)$ may be 1
 (C) period of $f(x)$ can't be 1 (D) fundamental period of $f(x)$ is 8

Comprehension # 4

Let $f(x) = \frac{x^3}{3} + \frac{x^2}{2} + ax + b \quad \forall x \in \mathbb{R}$

- Least value of 'a' for which $f(x)$ is injective function, is
 (A) $\frac{1}{4}$ (B) 1 (C) $\frac{1}{2}$ (D) $\frac{1}{8}$
- If $a = -1$, then $f(x)$ is
 (A) bijective (B) many-one and onto (C) one-one and into (D) many-one and into
- $f(x)$ is invertible iff
 (A) $a \in \left[\frac{1}{4}, \infty\right)$, $b \in \mathbb{R}$ (B) $a \in \left[\frac{1}{8}, \infty\right)$, $b \in \mathbb{R}$
 (C) $a \in \left(-\infty, \frac{1}{4}\right]$, $b \in \mathbb{R}$ (D) $a \in \left(-\infty, \frac{1}{4}\right]$, $b \in \mathbb{R}$

Exercise # 4

[Subjective Type Questions]

1. Find the domain of definitions of the following functions :

(i) $f(x) = \sqrt{3 - 2^x - 2^{1-x}}$

(ii) $f(x) = (x^2 + x + 1)^{-3/2}$

(iii) $f(x) = \sqrt{\tan x - \tan^2 x}$

(iv) $f(x) = \bullet \log_{10}(1 - \bullet \log_{10}(x^2 - 5x + 16))$

(v) If $f(x) = \sqrt{x^2 - 5x + 4}$ & $g(x) = x + 3$, then find the domain of $\frac{f}{g}(x)$

(vi) $f(x) = \frac{1}{[x]} + \log_{1-\{x\}}(x^2 - 3x + 10) + \frac{1}{\sqrt{2-|x|}} + \frac{1}{\sqrt{\sec(\sin x)}}$

2. Find the range of the following functions :

(i) $f(x) = 1 - |x - 2|$

(ii) $f(x) = \frac{1}{\sqrt{x-5}}$

(iii) $f(x) = \frac{1}{2 - \cos 3x}$

(iv) $f(x) = \frac{x+2}{x^2 - 8x - 4}$

(v) $f(x) = \frac{x^2 - 2x + 4}{x^2 + 2x + 4}$

(vi) $f(x) = 3 \sin \sqrt{\frac{\pi^2}{16} - x^2}$

(vii) $f(x) = x^4 - 2x^2 + 5$

(viii) $f(x) = x^3 - 12x$, where $x \in [-3, 1]$

(ix) $f(x) = \sin^2 x + \cos^4 x$

3. Let f be a function such that $f(3) = 1$ and $f(3x) = x + f(3x - 3)$ for all x . Then find the value of $f(300)$.

4. Let $f(x) = \frac{9^x}{9^x + 3}$ then find the value of the sum $f\left(\frac{1}{2008}\right) + f\left(\frac{2}{2008}\right) + f\left(\frac{3}{2008}\right) + \dots + f\left(\frac{2007}{2008}\right)$

5. Examine whether the following functions are even or odd or neither even nor odd, where $[]$ denotes greatest integer function.

(i) $f(x) = \frac{(1 + 2^x)^7}{2^x}$

(ii) $f(x) = \frac{\sec x + x^2 - 9}{x \sin x}$

(iii) $f(x) = \sqrt{1+x+x^2} - \sqrt{1-x+x^2}$

(iv) $f(x) = \begin{cases} x|x|, & x \leq -1 \\ [1+x] + [1-x], & -1 < x < 1 \\ -x|x|, & x \geq 1 \end{cases}$

(v) $f(x) = \frac{2x(\sin x + \tan x)}{2\left[\frac{x+2\pi}{\pi}\right] - 3}$

6. Find the fundamental period of the following functions :

(i) $f(x) = 1 - \frac{\sin^2 x}{1 + \cot x} - \frac{\cos^2 x}{1 + \tan x}$

(ii) $f(x) = \tan \frac{\pi}{2} [x]$, where $[.]$ denotes greatest integer function.

(iii) $f(x) = \bullet \log (2 + \cos 3x)$

(iv) $f(x) = e^{\ln \sin x + \tan^3 x - \operatorname{cosec} (3x - 5)}$

(v) $f(x) = \sin x + \tan \frac{x}{2} + \sin \frac{x}{2^2} + \tan \frac{x}{2^3} + \dots + \sin \frac{x}{2^{n-1}} + \tan \frac{x}{2^n}$

(vi) $f(x) = \frac{\sin x + \sin 3x}{\cos x + \cos 3x}$

7. Let $f(x) = \begin{cases} 1+x, & 0 \leq x \leq 2 \\ 3-x, & 2 < x \leq 3 \end{cases}$, then find $(f \circ f)(x)$.

8. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ is a function satisfying $f(10-x) = f(x)$ and $f(2-x) = f(2+x) \forall x \in \mathbb{R}$. If $f(0) = 101$, then the minimum possible number of values of x satisfying $f(x) = 101 \forall x \in [0, 25]$ is

9. Show if $f(x) = \sqrt[n]{a-x^n}$, $x > 0$, $n \geq 2$, $n \in \mathbb{N}$, then $(f \circ f)(x) = x$. Find also the inverse of $f(x)$.

10. Let $f: \mathbb{N} \rightarrow \mathbb{N}$, where $f(x) = x + (-1)^{x-1}$, then find the inverse of f .

Exercise # 5

Part # I

[Previous Year Questions] [AIEEE/JEE-MAIN]

1. Which of the following is not a periodic function- [AIEEE 2002]
 (1) $\sin 2x + \cos x$ (2) $\cos \sqrt{x}$ (3) $\tan 4x$ (4) $\log \cos 2x$
2. The period of $\sin^2 x$ is- [AIEEE 2002]
 (1) $\pi/2$ (2) π (3) $3\pi/2$ (4) 2π
3. The function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = \sin x$ is- [AIEEE 2002]
 (1) into (2) onto (3) one-one (4) many-one
4. The range of the function $f(x) = \frac{2+x}{2-x}$, $x \neq 2$ is- [AIEEE 2002]
 (1) \mathbb{R} (2) $\mathbb{R} - \{-1\}$ (3) $\mathbb{R} - \{1\}$ (4) $\mathbb{R} - \{2\}$
5. The domain of $\sin^{-1} \left[\log_3 \left(\frac{x}{3} \right) \right]$ is- [AIEEE 2002]
 (1) $[1, 9]$ (2) $[-1, 9]$ (3) $[-9, 1]$ (4) $[-9, -1]$
6. The function $f(x) = \log(x + \sqrt{x^2 + 1})$, is- [AIEEE 2003]
 (1) neither an even nor an odd function (2) an even function
 (3) an odd function (4) a periodic function
7. Domain of definition of the function $f(x) = \frac{3}{4-x^2} + \log_{10}(x^3 - x)$, is- [AIEEE 2003]
 (1) $(-1, 0) \cup (1, 2) \cup (2, \infty)$ (2) $(1, 2)$
 (3) $(-1, 0) \cup (1, 2)$ (4) $(1, 2) \cup (2, \infty)$
8. If $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfies $f(x+y) = f(x) + f(y)$, for all $x, y \in \mathbb{R}$ and $f(1) = 7$, then $\sum_{r=1}^n f(r)$ is - [AIEEE 2003]
 (1) $\frac{7n(n+1)}{2}$ (2) $\frac{7n}{2}$ (3) $\frac{7(n+1)}{2}$ (4) $7n(n+1)$
9. A function f from the set of natural numbers to integers defined by $f(n) = \begin{cases} \frac{n-1}{2}, & \text{when } n \text{ is odd} \\ -\frac{n}{2}, & \text{when } n \text{ is even} \end{cases}$ is - [AIEEE 2003]
 (1) neither one-one nor onto (2) one-one but not onto
 (3) onto but not one-one (4) one-one and onto both

10. The domain of the function $f(x) = \frac{\sin^{-1}(x-3)}{\sqrt{9-x^2}}$ is [AIEEE 2004]
 (1) $[1, 2]$ (2) $[2, 3]$ (3) $[1, 2]$ (4) $[2, 3]$
11. The range of the function $f(x) = {}^{7-x}P_{x-3}$ is- [AIEEE 2004]
 (1) $\{1, 2, 3, 4, 5\}$ (2) $\{1, 2, 3, 4, 5, 6\}$ (3) $\{1, 2, 3\}$ (4) $\{1, 2, 3, 4\}$
12. If $f : \mathbb{R} \rightarrow \mathbb{S}$ defined by $f(x) = \sin x - \sqrt{3} \cos x + 1$ is onto, then the interval of \mathbb{S} is- [AIEEE 2004]
 (1) $[-1, 3]$ (2) $[-1, 1]$ (3) $[0, 1]$ (4) $[0, -1]$
13. Let $f : (-1, 1) \rightarrow \mathbb{B}$, be a function defined by $f(x) = \tan^{-1} \frac{2x}{1-x^2}$, then f is both one-one and onto when \mathbb{B} is the interval- [AIEEE 2005]
 (1) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ (2) $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (3) $\left(0, \frac{\pi}{2}\right)$ (4) $\left[0, \frac{\pi}{2}\right]$
14. A real valued function $f(x)$ satisfies the function equation $f(x-y) = f(x)f(y) - f(a-x)f(a+y)$ where a is a given constant and $f(0) = 1$, $f(2a-x)$ is equal to [AIEEE 2005]
 (1) $f(1) + f(a-x)$ (2) $f(-x)$ (3) $-f(x)$ (4) $f(x)$
15. If x is real, the maximum value of $\frac{3x^2+9x+17}{3x^2+9x+7}$ is- [AIEEE 2006]
 (1) 41 (2) 1 (3) $\frac{17}{7}$ (4) $\frac{1}{4}$
16. The largest interval lying in $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ for which the function is defined, $\left[f(x) = 4^{-x^2} + \cos^{-1}\left(\frac{x}{2}-1\right) + \log(\cos x)\right]$ is [AIEEE 2007]
 (1) $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (2) $\left[-\frac{\pi}{4}, \frac{\pi}{2}\right)$ (3) $\left[0, \frac{\pi}{2}\right)$ (4) $[0, \pi]$
17. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by $f(x) = \text{Min} \{x+1, |x|+1\}$. Then which of the following is true ? [AIEEE 2007]
 (1) $f(x)$ is not differentiable at $x = 1$ (2) $f(x)$ is differentiable everywhere
 (3) $f(x)$ is not differentiable at $x = 0$ (4) $f(x) \geq 1$ for all $x \in \mathbb{R}$

18. Let $f: \mathbb{N} \rightarrow \mathbb{Y}$ be a function defined as $f(x) = 4x + 3$ where [AIEEE 2008]

$\mathbb{Y} = \{y \in \mathbb{N} : y = 4x + 3 \text{ for some } x \in \mathbb{N}\}$. So that f is invertible and its inverse is

(1) $g(y) = \frac{3y+4}{3}$ (2) $g(y) = 4 + \frac{y+3}{4}$ (3) $g(y) = \frac{y+3}{4}$ (4) $g(y) = \frac{y-3}{4}$

19. For real x , let $f(x) = x^3 + 5x + 1$, then :- [AIEEE 2009]

- (1) f is one-one and onto \mathbb{R} (2) f is neither one-one nor onto \mathbb{R}
(3) f is one-one but not onto \mathbb{R} (4) f is onto \mathbb{R} but not one-one

20. Let $f(x) = (x+1)^2 - 1, x \geq -1$. [AIEEE 2009]

Statement-1 : The set $\{x : f(x) = f^{-1}(x)\} = \{0, -1\}$.

Statement-2 : f is a bijection.

- (1) Statement-1 is true, Statement-2 is false.
(2) Statement-1 is false, Statement-2 is true.
(3) Statement-1 is true, Statement-2 is true ; Statement-2 is a correct explanation for Statement-1.
(4) Statement-1 is true, Statement-2 is true ; Statement-2 is not a correct explanation for statement-1.

21. The domain of the function $f(x) = \frac{1}{\sqrt{|x|} - x}$ is :- [AIEEE 2011]

- (1) $(-\infty, 0)$ (2) $(-\infty, \infty) - \{0\}$ (3) $(-\infty, \infty)$ (4) $(0, \infty)$

22. Let f be a function defined by $f(x) = (x-1)^2 + 1, (x \geq 1)$ [AIEEE 2011]

Statement - 1 : The set $\{x : f(x) = f^{-1}(x)\} = \{1, 2\}$

Statement - 2 : f is bijection and $f^{-1}(x) = 1 + \sqrt{x-1}, x \geq 1$.

- (1) Statement-1 is true, Statement-2 is false.
(2) Statement-1 is false, Statement-2 is true.
(3) Statement-1 is true, Statement-2 is true ; Statement-2 is a correct explanation for Statement-1.
(4) Statement-1 is true, Statement-2 is true ; Statement-2 is not a correct explanation for statement-1.

23. If $f: \mathbb{R} \rightarrow \mathbb{R}$ is a function defined by $f(x) = [x] \cos \pi \left(\frac{2x-1}{2} \right)$, where $[x]$ denotes the greatest integer function, then f is : [AIEEE 2012]
- (1) continuous only at $x = 0$.
 (2) continuous for every real x .
 (3) discontinuous only at $x = 0$.
 (4) discontinuous only at non-zero integral values of x .
24. If $X = \{4^n - 3n - 1 : n \in \mathbb{N}\}$ and $Y = \{9(n-1) : n \in \mathbb{N}\}$, where \mathbb{N} is the set of natural numbers, then $X \cup Y$ is equal to : [Main 2014]
- (1) \mathbb{N} (2) $Y - X$ (3) X (4) Y
25. If $f(x) + 2f\left(\frac{1}{x}\right) = 3x$, $x \neq 0$ and $S = \{x \in \mathbb{R} : f(x) = f(-x)\}$; then S : [Main 2016]
- (1) contains exactly one element. (2) contains exactly two elements.
 (3) contains more than two elements. (4) is an empty set.
26. For $x \in \mathbb{R}$, $f(x) = |\log 2 - \sin x|$ and $g(x) = f(f(x))$, then: [Main 2016]
- (1) $g'(0) = \cos(\log 2)$ (2) $g'(0) = -\cos(\log 2)$
 (3) g is differentiable at $x = 0$ and $g'(0) = -\sin(\log 2)$ (4) g is not differentiable at $x = 0$

Part # II

[Previous Year Questions][IIT-JEE ADVANCED]

1. The domain of definition of the function, $y(x)$ given by the equation, $2^x + 2^y = 2$ is : [JEE 2000]
- (A) $0 < x \leq 1$ (B) $0 \leq x \leq 1$ (C) $-\infty < x \leq 0$ (D) $-\infty < x < 1$
2. Given $x = \{1, 2, 3, 4\}$, find all one-one, onto mappings, $f: X \rightarrow X$ such that, $f(1) = 1$, $f(2) \neq 2$ and $f(4) \neq 4$. [JEE 2000]
3. Let $g(x) = 1 + x - [x]$ & $f(x) = \begin{cases} -1, & x < 0 \\ 0, & x = 0 \\ 1, & x > 0 \end{cases}$. Then for all x , $f(g(x))$ is equal to [JEE 2001]
- (A) x (B) 1 (C) $f(x)$ (D) $g(x)$
- where $[]$ denotes the greatest integer function.
4. If $f: [1, \infty) \rightarrow [2, \infty)$ is given by, $f(x) = x + \frac{1}{x}$, then $f^{-1}(x)$ equals : [JEE 2001]
- (A) $\frac{x + \sqrt{x^2 - 4}}{2}$ (B) $\frac{x}{1 + x^2}$ (C) $\frac{x - \sqrt{x^2 - 4}}{2}$ (D) $1 - \sqrt{x^2 - 4}$

5. The domain of definition of $f(x) = \frac{\log_2(x+3)}{x^2 + 3x + 2}$ is : [JEE 2001]
 (A) $\mathbb{R} \setminus \{-1, -2\}$ (B) $(-2, \infty)$ (C) $\mathbb{R} \setminus \{-1, -2, -3\}$ (D) $(-3, \infty) \setminus \{-1, -2\}$
6. Let $E = \{1, 2, 3, 4\}$ & $F = \{1, 2\}$. Then the number of onto functions from E to F is [JEE 2001]
 (A) 14 (B) 16 (C) 12 (D) 8
7. Let $f(x) = \frac{\alpha x}{x+1}$, $x \neq -1$. Then for what value of α is $f(f(x)) = x$? [JEE 2001]
 (A) $\sqrt{2}$ (B) $-\sqrt{2}$ (C) 1 (D) -1
8. Suppose $f(x) = (x+1)^2$ for $x \geq -1$. If $g(x)$ is the function whose graph is the reflection of the graph of $f(x)$ with respect to the line $y = x$, then $g(x)$ equals -
 (A) $-\sqrt{x} - 1, x \geq 0$ (B) $\frac{1}{(1+x)^2}, x \geq -1$ (C) $\sqrt{x+1}, x \geq -1$ (D) $\sqrt{x} - 1, x \geq 0$
9. Let function $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = 2x + \sin x$ for $x \in \mathbb{R}$. Then f is - [JEE 2002]
 (A) one to one and onto (B) one to one but not onto
 (C) onto but not one to one (D) neither one to one nor onto
10. Range of the function $f(x) = \frac{x^2 + x + 2}{x^2 + x + 1}$ is - [JEE 2003]
 (A) $[1, 2]$ (B) $[1, \infty)$ (C) $\left[2, \frac{7}{3}\right]$ (D) $\left(1, \frac{7}{3}\right]$
11. Let $f(x) = \frac{x}{1+x}$ defined from $(0, \infty) \rightarrow [0, \infty)$, then $f(x)$ is -
 (A) one-one but not onto (B) one-one and onto
 (C) Many one but not onto (D) Many one and onto
12. Let $f(x) = \sin x + \cos x$, $g(x) = x^2 - 1$. Thus $g(f(x))$ is invertible for $x \in$ [JEE 2004]
 (A) $\left[-\frac{\pi}{2}, 0\right]$ (B) $\left[-\frac{\pi}{2}, \pi\right]$ (C) $\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]$ (D) $\left[0, \frac{\pi}{2}\right]$
13. If functions $f(x)$ and $g(x)$ are defined on $\mathbb{R} \rightarrow \mathbb{R}$ such that $f(x) = \begin{cases} 0, & x \in \text{rational} \\ x, & x \in \text{irrational} \end{cases}$, $g(x) = \begin{cases} 0, & x \in \text{irrational} \\ x, & x \in \text{rational} \end{cases}$, then $(f-g)(x)$ is - [JEE 2005]
 (A) one-one and onto (B) neither one-one nor onto
 (C) one-one but not onto (D) onto but not one-one

14. Let $f(x) = x^2$ and $g(x) = \sin x$ for all $x \in \mathbb{R}$. Then the set of all x satisfying $(f \circ g \circ g \circ f)(x) = (g \circ g \circ f)(x)$, where $(f \circ g)(x) = f(g(x))$, is- [JEE 2011]
- (A) $\pm\sqrt{n\pi}, n \in \{0, 1, 2, \dots\}$ (B) $\pm\sqrt{n\pi}, n \in \{1, 2, \dots\}$
- (C) $\frac{\pi}{2} + 2n\pi, n \in \{\dots, -2, -1, 0, 1, 2, \dots\}$ (D) $2n\pi, n \in \{\dots, -2, -1, 0, 1, 2, \dots\}$
15. The function $f : [0, 3] \rightarrow [1, 29]$, defined by $f(x) = 2x^3 - 15x^2 + 36x + 1$, is: [JEE 2012]
- (A) one-one and onto (B) onto but not one-one
- (C) one-one but not onto (D) neither one-one nor onto
16. Let $f : (-1, 1) \rightarrow \mathbb{R}$ be such that $f(\cos 4\theta) = \frac{2}{2 - \sec^2 \theta}$ for $\theta \in \left(0, \frac{\pi}{4}\right) \cup \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$. Then the value(s) of $f\left(\frac{1}{3}\right)$ is (are)- [JEE 2012]
- (A) $1 - \sqrt{\frac{3}{2}}$ (B) $1 + \sqrt{\frac{3}{2}}$ (C) $1 - \sqrt{\frac{2}{3}}$ (D) $1 + \sqrt{\frac{2}{3}}$
17. For every pair of continuous functions $f, g : [0, 1] \rightarrow \mathbb{R}$ such that $\max \{f(x) : x \in [0, 1]\} = \max \{g(x) : x \in [0, 1]\}$, the correct statement(s) is (are): [JEE Ad. 2014]
- (A) $(f(c))^2 + 3f(c) = (g(c))^2 + 3g(c)$ for some $c \in [0, 1]$
- (B) $(f(c))^2 + f(c) = (g(c))^2 + 3g(c)$ for some $c \in [0, 1]$
- (C) $(f(c))^2 + 3f(c) = (g(c))^2 + g(c)$ for some $c \in [0, 1]$
- (D) $(f(c))^2 = (g(c))^2$ for some $c \in [0, 1]$
18. Let $f : \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \rightarrow \mathbb{R}$ be given by [JEE Ad. 2014]
- $f(x) = (\log(\sec x + \tan x))^3$
- Then
- (A) $f(x)$ is an odd function (B) $f(x)$ is a one-one function
- (C) $f(x)$ is an onto function (D) $f(x)$ is an even function
19. Let $f_1 : \mathbb{R} \rightarrow \mathbb{R}, f_2 : [0, \infty) \rightarrow \mathbb{R}, f_3 : \mathbb{R} \rightarrow \mathbb{R}$ and $f_4 : \mathbb{R} \rightarrow [0, \infty)$ be defined by [JEE Ad. 2014]
- $f_1(x) = \begin{cases} |x| & \text{if } x < 0 \\ e^x & \text{if } x \geq 0 \end{cases}; f_2(x) = x^2; f_3(x) = \begin{cases} \sin x & \text{if } x < 0 \\ x & \text{if } x \geq 0 \end{cases}; f_4(x) = \begin{cases} f_2(f_1(x)) & \text{if } x < 0 \\ f_2(f_1(x)) - 1 & \text{if } x \geq 0 \end{cases}$
- List - I**
- (p) f_4 is
- (q) f_3 is
- (r) $f_2 \circ f_1$ is
- (s) f_2 is
- List - II**
- (1) onto but not one-one
- (2) neither continuous nor one-one
- (3) differentiable but not one-one
- (4) continuous and one-one

Codes :

	p	q	r	s
(A)	3	1	4	2
(B)	1	3	4	2
(C)	3	1	2	4
(D)	1	3	2	4

20. Let $f(x) = \sin\left(\frac{\pi}{6} \sin\left(\frac{\pi}{2} \sin x\right)\right)$ for all $x \in \mathbb{R}$ and $g(x) = \frac{\pi}{2} \sin x$ for all $x \in \mathbb{R}$. Let $(f \circ g)(x)$ denote $f(g(x))$ and $(g \circ f)(x)$ denote $g(f(x))$. Then which of the following is (are) true ? [JEE Ad. 2015]

(A) Range of f is $\left[-\frac{1}{2}, \frac{1}{2}\right]$

(B) Range of $f \circ g$ is $\left[-\frac{1}{2}, \frac{1}{2}\right]$

(C) $\lim_{x \rightarrow 0} \frac{f(x)}{g(x)} = \frac{\pi}{6}$

(D) There is an $x \in \mathbb{R}$ such that $(g \circ f)(x) = 1$

21. Let $f: \mathbb{R} \rightarrow \mathbb{R}$, $g: \mathbb{R} \rightarrow \mathbb{R}$ and $h: \mathbb{R} \rightarrow \mathbb{R}$ be differentiable functions such that $f(x) = x^3 + 3x + 2$, $(12\alpha + 20)\frac{K^2}{2} = K^3$, $g(f(x)) = x$ and $h(g(g(x))) = x$ for all $x \in \mathbb{R}$. Then [JEE Ad. 2016]

(A) $g'(2) = \frac{1}{15}$

(B) $h'(1) = 666$

(C) $h(0) = 16$

(D) $h(g(3)) = 36$

MOCK TEST

SECTION - I : STRAIGHT OBJECTIVE TYPE

1. If $f(x) \cdot f(y) = f(x) + f(y) + f(xy) - 2 \quad \forall x, y \in \mathbb{R}$ and if $f(x)$ is not a constant function, then the value of $f(1)$ is equal to
 (A) 1 (B) 2 (C) 0 (D) -1
2. The domain of the function $f(x) = \sqrt{-\log_{\frac{x+4}{2}} \left(\log_2 \frac{2x-1}{3+x} \right)}$ is
 (A) $(-4, -3) \cup (4, \infty)$ (B) $(-\infty, -3) \cup (4, \infty)$ (C) $(-\infty, -4) \cup (3, \infty)$ (D) None
3. Let $f(x) = ax^2 + bx + c$, where a, b, c are rational and $f : \mathbb{Z} \rightarrow \mathbb{Z}$, where \mathbb{Z} is the set of integers. Then $a + b$ is :
 (A) a negative integer (B) an integer
 (C) non-integral rational number (D) none of these
4. If $f(x) = \frac{\sin^2 x + 4 \sin x + 5}{2 \sin^2 x + 8 \sin x + 8}$, then range of $f(x)$ is
 (A) $\left(\frac{1}{2}, \infty\right)$ (B) $\left(\frac{5}{9}, 1\right)$ (C) $\left[\frac{5}{9}, 1\right]$ (D) $\left[\frac{5}{9}, \infty\right)$
5. If $f(x) = x + \tan x$ and $g(x)$ is the inverse of $f(x)$ then $g'(x)$ is equal to
 (A) $\frac{1}{1 + (g(x) - x)^2}$ (B) $\frac{1}{2 + (g(x) - x)^2}$ (C) $\frac{1}{2 + (g(x) - x)^2}$ (D) none of these
6. Let $f(x) = \tan x$, $g(f(x)) = f\left(x - \frac{\pi}{4}\right)$, where $f(x)$ and $g(x)$ are real valued functions. For all possible values of x , $f(g(x)) =$
 (A) $\tan\left(\frac{x-1}{x+1}\right)$ (B) $\tan(x-1) - \tan(x+1)$ (C) $\frac{f(x)+1}{f(x)-1}$ (D) $\frac{x - \pi/4}{x + \pi/4}$
7. The range of the function $f(x) = \sin^{-1}\left[x^2 + \frac{1}{2}\right] + \cos^{-1}\left[x^2 - \frac{1}{2}\right]$, where $[]$ is the greatest integer function, is:
 (A) $\left\{\frac{\pi}{2}, \pi\right\}$ (B) $\left\{0, \frac{\pi}{2}\right\}$ (C) $\{\pi\}$ (D) $\left(0, \frac{\pi}{2}\right)$

8. It is given that $f(x)$ is a function defined on \mathbb{R} , satisfying $f(1) = 1$ and for any $x \in \mathbb{R}$
 $f(x+5) \geq f(x) + 5$
 and $f(x+1) \leq f(x) + 1$
 If $g(x) = f(x) + 1 - x$, then $g(2013)$ equals
 (A) 2014 (B) 2013 (C) 1 (D) 0
9. The image of the interval $[-1, 3]$ under the mapping specified by the function $f(x) = 4x^3 - 12x$ is :
 (A) $[f(+1), f(-1)]$ (B) $[f(-1), f(3)]$ (C) $[-8, 16]$ (D) $[-8, 72]$
10. Let $f(x) = x(2-x)$, $0 \leq x \leq 2$. If the definition of ' f ' is extended over the set,
 $\mathbb{R} - [0, 2]$ by $f(x-2) = f(x)$, then ' f ' is a :
 (A) periodic function of period 1 (B) non-periodic function
 (C) periodic function of period 2 (D) periodic function of period $1/2$

SECTION - II : MULTIPLE CORRECT ANSWER TYPE

11. Suppose $f(x) = ax + b$ and $g(x) = bx + a$, where a and b are positive integers.
 If $f(g(50)) - g(f(50)) = 28$ then the product (ab) can have the value equal to
 (A) 12 (B) 48 (C) 180 (D) 210
12. Let $f(x) = \begin{cases} 0 & \text{for } x = 0 \\ x^2 \sin\left(\frac{\pi}{x}\right) & \text{for } -1 < x < 1 \ (x \neq 0) \\ x |x| & \text{for } x > 1 \text{ or } x < -1 \end{cases}$, then:
 (A) $f(x)$ is an odd function (B) $f(x)$ is an even function
 (C) $f(x)$ is neither odd nor even (D) $f'(x)$ is an even function
13. Which of the functions defined below are one-one function(s) ?
 (A) $f(x) = (x+1)$, $(x \geq -1)$ (B) $g(x) = x + (1/x)$ $(x > 0)$
 (C) $h(x) = x^2 + 4x - 5$, $(x > 0)$ (D) $f(x) = e^{-x}$, $(x \geq 0)$
14. If the function $f(x) = ax + b$ has its own inverse then the ordered pair (a, b) can be
 (A) $(1, 0)$ (B) $(-1, 0)$ (C) $(-1, 1)$ (D) $(1, 1)$
15. A continuous function $f(x)$ on $\mathbb{R} \rightarrow \mathbb{R}$ satisfies the relation
 $f(x) + f(2x+y) + 5xy = f(3x-y) + 2x^2 + 1$ for $\forall x, y \in \mathbb{R}$
 then which of the following hold(s) good ?
 (A) f is many one (B) f has no minima
 (C) f is neither odd nor even (D) f is bounded

SECTION - III : ASSERTION AND REASON TYPE

16. Let $g : \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) = \{e^x\}$, where $\{x\}$ denotes fractional part function.
Statement-I : $g(x)$ is periodic function.
Statement-II : $\{x\}$ is periodic function.
 (A) Statement-I is true, statement-II is true and statement-II is correct explanation for statement-I.
 (B) Statement-I is true, statement-II is true and statement-II is NOT the correct explanation for statement-I.
 (C) Statement-I is true, statement-II is false.
 (D) Statement-I is false, statement-II is true
17. **Statement-I :** Fundamental period of $\sin x + \tan x$ is 2π
Statement-II : If the period of $f(x)$ is T_1 and the period of $g(x)$ is T_2 , then the fundamental period of $f(x) + g(x)$ is the L.C.M. of T_1 and T_2
 (A) Statement-I is true, statement-II is true and statement-II is correct explanation for statement-I.
 (B) Statement-I is true, statement-II is true and statement-II is NOT the correct explanation for statement-I.
 (C) Statement-I is true, statement-II is false.
 (D) Statement-I is false, statement-II is true
18. **Statement-I :** If a function $y = f(x)$ is symmetric about $y = x$, then $f(f(x)) = x$
Statement-II : If $f(x) = \begin{cases} x & : x \text{ is rational} \\ 1-x & : x \text{ is irrational} \end{cases}$, then $f(f(x)) = x$
 (A) Statement-I is true, statement-II is true and statement-II is correct explanation for statement-I.
 (B) Statement-I is true, statement-II is true and statement-II is NOT the correct explanation for statement-I.
 (C) Statement-I is true, statement-II is false.
 (D) Statement-I is false, statement-II is true
19. **Statement-1 :** f is an even function, g and h are odd functions, all 3 are polynomials. Given $f(1) = 0$, $f(2) = 1$, $f(3) = -5$, $g(1) = 1$, $g(-3) = 2$, $g(5) = 3$, $h(1) = 3$, $h(3) = 5$ and $h(5) = 1$.
 The value of $f(g(h(1))) + g(h(f(3))) + h(f(g(-1)))$ is equal to zero.
Statement-2 : If a polynomial function $P(x)$ is odd then $P(0) = 0$.
 (A) Statement-I is true, statement-II is true and statement-II is correct explanation for statement-I.
 (B) Statement-I is true, statement-II is true and statement-II is NOT the correct explanation for statement-I.
 (C) Statement-I is true, statement-II is false.
 (D) Statement-I is false, statement-II is true
20. **Statement -1 :** e^x can not be expressed as the sum of even and odd function.
Statement -2 : e^x is neither even nor odd function
 (A) Statement-I is true, statement-II is true and statement-II is correct explanation for statement-I.
 (B) Statement-I is true, statement-II is true and statement-II is NOT the correct explanation for statement-I.
 (C) Statement-I is true, statement-II is false.
 (D) Statement-I is false, statement-II is true

SECTION - IV : MATRIX - MATCH TYPE

21. Column-I

Column-II

- | | |
|---|-------------------------|
| (A) Function $f: \left[0, \frac{\pi}{3}\right] \rightarrow [0, 1]$ defined by $f(x) = \sqrt{\sin x}$ is | (p) one to one function |
| (B) Function $f: (1, \infty) \rightarrow (1, \infty)$ defined by $f(x) = \frac{x+3}{x-1}$ is | (q) many – one function |
| (C) Function $f: \left[-\frac{\pi}{2}, \frac{4\pi}{3}\right] \rightarrow [-1, 1]$ defined by $f(x) = \sin x$ is | (r) into function |
| (D) Function $f: (2, \infty) \rightarrow [8, \infty)$ defined by $f(x) = \frac{x^2}{x-2}$ is | (s) onto function |

22. Let $f(x) = x + \frac{1}{x}$ and $g(x) = \frac{x+1}{x+2}$.

Match the composite function given in Column-I with their respective domains given in Column-II.

Column-I

Column-II

- | | |
|---------|---------------------------------|
| (A) fog | (p) $\mathbb{R} - \{-2, -5/3\}$ |
| (B) gof | (q) $\mathbb{R} - \{-1, 0\}$ |
| (C) fof | (r) $\mathbb{R} - \{0\}$ |
| (D) gog | (s) $\mathbb{R} - \{-2, -1\}$ |

SECTION - V : COMPREHENSION TYPE

23. Read the following comprehension carefully and answer the questions.

Let $f(x) = x^2 - 2x - 1 \quad \forall x \in \mathbb{R}$. Let $f: (-\infty, a] \rightarrow [b, \infty)$, where 'a' is the largest real number for which $f(x)$ is bijective.

1. The value of $(a + b)$ is equal to

- (A) -2 (B) -1 (C) 0 (D) 1

2. Let $f: \mathbb{R} \rightarrow \mathbb{R}$, $g(x) = f(x) + 3x - 1$, then the least value of function $y = g(|x|)$ is

- (A) -9/4 (B) -5/4 (C) -2 (D) -1

3. Let $f: [a, \infty) \rightarrow [b, \infty)$, then $f^{-1}(x)$ is given by

- (A) $1 + \sqrt{x+2}$ (B) $1 - \sqrt{x+3}$ (C) $1 - \sqrt{x+2}$ (D) $1 + \sqrt{x+3}$

4. Let $f: \mathbb{R} \rightarrow \mathbb{R}$, then range of values of k for which equation $f(|x|) = k$ has 4 distinct real roots is

- (A) $(-2, -1)$ (B) $(-2, 0)$ (C) $(-1, 0)$ (D) $(0, 1)$

24. Read the following comprehension carefully and answer the questions.

$$\text{Let } f(x) = \begin{cases} 2x + a & : x \geq -1 \\ bx^2 + 3 & : x < -1 \end{cases}$$

$$\text{and } g(x) = \begin{cases} x + 4 & : 0 \leq x \leq 4 \\ -3x - 2 & : -2 < x < 0 \end{cases}$$

1. $g(f(x))$ is not defined if
 (A) $a \in (6, \infty), b \in (5, \infty)$ (B) $a \in (4, 6), b \in (5, \infty)$ (C) $a \in (6, \infty), b \in (0, 1)$ (D) $a \in (4, 6), b \in (1, 5)$
2. If domain of $g(f(x))$ is $[-1, 2]$, then
 (A) $a = 1, b > 5$ (B) $a = 2, b > 7$ (C) $a = 2, b > 10$ (D) $a = 0, b \in \mathbb{R}$
3. If $a = 2$ and $b = 3$ then range of $g(f(x))$ is
 (A) $(-2, 8]$ (B) $(0, 8]$ (C) $[4, 8]$ (D) $[-1, 8]$

25. Read the following comprehension carefully and answer the questions.

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ is a function satisfying $f(2-x) = f(2+x)$ and $f(20-x) = f(x), \forall x \in \mathbb{R}$. For this function f answer the following.

1. If $f(0) = 5$, then minimum possible number of values of x satisfying $f(x) = 5$, for $x \in [0, 170]$, is
 (A) 21 (B) 12 (C) 11 (D) 22
2. Graph of $y = f(x)$ is
 (A) symmetrical about $x = 18$ (B) symmetrical about $x = 5$
 (C) symmetrical about $x = 8$ (D) symmetrical about $x = 20$
3. If $f(2) \neq f(6)$, then
 (A) fundamental period of $f(x)$ is 1 (B) fundamental period of $f(x)$ may be 1
 (C) period of $f(x)$ can't be 1 (D) fundamental period of $f(x)$ is 8

SECTION - VI : INTEGER TYPE

26. If $f(x) + f(y) + f(xy) = 2 + f(x) \cdot f(y)$, for all real values of x and y and $f(x)$ is a polynomial function with $f(4) = 17$ and $f(1) \neq 1$, then find the value of $f(5)$.
27. If $f(x) + f(y) + f(xy) = 2 + f(x) \cdot f(y)$, for all real values of x & y and $f(x)$ is a polynomial function with $f(4) = 17$, then find the value of $f(5)/14$, where $f(1) \neq 1$.
28. If f is a function satisfying the condition $f(x) + f(y) = f(x\sqrt{1-y^2} + y\sqrt{1-x^2})$ for all x and y in domain of f , then find value of $f(4x^3 - 3x) + 3f(x)$.
29. If domain of $f(x) = \frac{\sin^{-1}(\sin x)}{\sqrt{-\log\left(\frac{x+4}{2}\right) \log_2\left(\frac{2x-1}{3+x}\right)}}$ is $(a, b) \cup (c, \infty)$, then find the value of $a + b + 3c$.
30. The functional relation $f(x) + f\left(\frac{1}{1-x}\right) = \frac{2(1-2x)}{x(1-x)}$ is satisfying by the function $f(x) = \frac{x+1}{\lambda(x-1)}$, then find value of λ .

ANSWER KEY

EXERCISE - 1

1. C 2. D 3. A 4. B 5. B 6. B 7. A 8. B 9. D 10. C 11. A 12. C 13. D
 14. A 15. D 16. C 17. C 18. D 19. B 20. A 21. D 22. C 23. B 24. D 25. A 26. B
 27. D 28. C 29. B 30. D 31. D 32. C 33. C 34. A 35. B 36. B

EXERCISE - 2 : PART # I

1. ABC 2. ACD 3. AC 4. BD 5. AD 6. AD 7. B 8. A 9. ABCD
 10. BCD 11. AB 12. BC 13. ACD 14. BC 15. B 16. BCD 17. AC 18. AD
 19. ABD 20. BD 21. ABC 22. ABD 23. BCD 24. ABC

PART - II

1. C 2. D 3. C 4. D 5. A

EXERCISE - 3 : PART # I

1. $A \rightarrow q$ $B \rightarrow s$ $C \rightarrow p$ $D \rightarrow r$ 2. $A \rightarrow q$ $B \rightarrow r$ $C \rightarrow p$ $D \rightarrow s$
 3. $A \rightarrow q, r$ $B \rightarrow q, r$ $C \rightarrow q$ $D \rightarrow s$ 4. $A \rightarrow r$ $B \rightarrow p$ $C \rightarrow s$ $D \rightarrow q$

PART - II

Comprehension #1: 1. A 2. C 3. D

Comprehension #2: 1. A 2. B 3. C

Comprehension #3: 1. D 2. A 3. C

Comprehension #4: 1. A 2. B 3. A

EXERCISE - 5 : PART # I

1. 2 2. 2 3. 1,4 4. 2 5. 1 6. 3 7. 1 8. 1 9. 4 10. 2 11. 3 12. 1 13. 2
 14. 3 15. 1 16. 3 17. 2 18. 4 19. 1 20. 4 21. 1 22. 2 23. 2 24. 4 25. 2 26. 1

PART - II

1. D 2. $\{(1, 1), (2, 3), (3, 4), (4, 2)\}; \{(1, 1), (2, 4), (3, 2), (4, 3)\}$ and $\{(1, 1), (2, 4), (3, 3), (4, 2)\}$ 3. B 4. A
 5. D 6. A 7. D 8. D 9. A 10. D 11. A 12. C 13. A 14. A 15. B
 16. (zero marks to all) 17. AD 18. ABC 19. D 20. ABC 21. BC

MOCK TEST

1. B 2. A 3. B 4. C 5. C 6. A 7. C 8. C 9. D 10. C 11. A, D 12. A, D
13. A, C, D 14. A, B, C 15. A, B 16. D 17. C 18. A 19. A 20. D
21. $A \rightarrow p, r$ $B \rightarrow p, s$ $C \rightarrow q, s$ $D \rightarrow q, s$ 22. $A \rightarrow s$ $B \rightarrow q$ $C \rightarrow r$ $D \rightarrow p$
23. 1. B 2. C 3. A 4. A 24. 1. A 2. A 3. C 25. 1. A 2. A 3. C
26. 8 27. 9 28. 0 29. 5 30. 1