

SOLVED EXAMPLES

Ex. 1 Evaluate $\int \frac{\cos^3 x}{\sin^2 x + \sin x} dx$

Sol. $I = \int \frac{(1 - \sin^2 x) \cos x}{\sin x(1 + \sin x)} dx = \int \frac{1 - \sin x}{\sin x} \cos x dx$

Put $\sin x = t \Rightarrow \cos x dx = dt$

$$\Rightarrow I = \int \frac{1-t}{t} dt = \ln|t| - t + c = \ln|\sin x| - \sin x + c$$

Ex. 2 Evaluate $\int \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}} \cdot \frac{1}{x} dx$

Sol. Put $x = \cos^2 \theta$

$$\Rightarrow dx = -2 \sin \theta \cos \theta d\theta$$

$$\Rightarrow I = \int \sqrt{\frac{1-\cos \theta}{1+\cos \theta}} \cdot \frac{1}{\cos^2 \theta} (-2 \sin \theta \cos \theta) d\theta = -\int 2 \tan \frac{\theta}{2} \tan \theta d\theta$$

$$= -4 \int \frac{\sin^2(\theta/2)}{\cos \theta} d\theta = -2 \int \frac{1-\cos \theta}{\cos \theta} d\theta = -2 \ln|\sec \theta - \tan \theta| + 2\theta + c$$

$$= -2 \ln \left| \frac{1+\sqrt{1-x}}{x} \right| + 2 \cos^{-1} \sqrt{x} + c$$

Ex. 3 Evaluate : $\int x \ln(1+x) dx$

Sol. Let $I = \int x \ln(1+x) dx$

$$= \bullet n(x+1) \cdot \frac{x^2}{2} - \int \frac{1}{x+1} \cdot \frac{x^2}{2} dx$$

$$= \frac{x^2}{2} \bullet n(x+1) - \frac{1}{2} \int \frac{x^2}{x+1} dx = \frac{x^2}{2} \bullet n(x+1) - \frac{1}{2} \int \frac{x^2-1+1}{x+1} dx$$

$$= \frac{x^2}{2} \bullet n(x+1) - \frac{1}{2} \int \left(\frac{x^2-1}{x+1} + \frac{1}{x+1} \right) dx$$

$$= \frac{x^2}{2} \bullet n(x+1) - \frac{1}{2} \int \left((x-1) + \frac{1}{x+1} \right) dx$$

$$= \frac{x^2}{2} \bullet n(x+1) - \frac{1}{2} \left[\frac{x^2}{2} - x + \ln|x+1| \right] + C$$

Ex. 4 Evaluate $\int \frac{(x^2 - 1) dx}{(x^4 + 3x^2 + 1) \tan^{-1} \left(x + \frac{1}{x} \right)}$

Sol. The given integral can be written as

$$I = \int \frac{\left(1 - \frac{1}{x^2} \right) dx}{\left[\left(x + \frac{1}{x} \right)^2 + 1 \right] \tan^{-1} \left(x + \frac{1}{x} \right)}$$

Let $\left(x + \frac{1}{x} \right) = t$. Differentiating we get $\left(1 - \frac{1}{x^2} \right) dx = dt$

Hence $I = \int \frac{dt}{(t^2 + 1) \tan^{-1} t}$

Now make one more substitution $\tan^{-1} t = u$. Then $\frac{dt}{t^2 + 1} = du$ and $I = \int \frac{du}{u} = \ln |u| + c$

Returning to t , and then to x , we have

$$I = \ln | \tan^{-1} t | + c = \ln \left| \tan^{-1} \left(x + \frac{1}{x} \right) \right| + c$$

Ex. 5 Evaluate : $\int \frac{x}{1 + \sin x} dx$

Sol. **Let** $I = \int \frac{x}{1 + \sin x} dx = \int \frac{x(1 - \sin x)}{(1 + \sin x)(1 - \sin x)} dx$

$$= \int \frac{x(1 - \sin x)}{1 - \sin^2 x} dx = \int \frac{x(1 - \sin x)}{\cos^2 x} dx = \int x \sec^2 x dx - \int x \sec x \tan x dx$$

$$= \left[x \int \sec^2 x dx - \int \left\{ \frac{dx}{dx} \int \sec^2 x dx \right\} dx \right] - \left[x \int \sec x \tan x dx - \int \left\{ \frac{dx}{dx} \int \sec x \tan x dx \right\} dx \right]$$

$$= \left[x \tan x - \int \tan x dx \right] - \left[x \sec x - \int \sec x dx \right]$$

$$= [x \tan x - \ln | \sec x |] - [x \sec x - \ln | \sec x + \tan x |] + c$$

$$= x(\tan x - \sec x) + \ln \left| \frac{(\sec x + \tan x)}{\sec x} \right| + c = \frac{-x(1 - \sin x)}{\cos x} + \ln |1 + \sin x| + c$$

Ex. 6 Evaluate : $\int e^x \left(\frac{1 - \sin x}{1 - \cos x} \right) dx$

Sol. Given integral $= \int e^x \left(\frac{1 - 2 \sin \frac{x}{2} \cos \frac{x}{2}}{2 \sin^2 \frac{x}{2}} \right) dx$

$$= \int e^x \left(\frac{1}{2} \operatorname{cosec}^2 \frac{x}{2} - \cot \frac{x}{2} \right) dx = -e^x \cot \frac{x}{2} + C$$

Ex. 7 The value of $\int e^x \left(\frac{x^4 + 2}{(1 + x^2)^{5/2}} \right) dx$ is equal to -

Sol. Let $I = \int e^x \left(\frac{x^4 + 2}{(1 + x^2)^{5/2}} \right) dx = \int e^x \left(\frac{1}{(1 + x^2)^{1/2}} + \frac{1 - 2x^2}{(1 + x^2)^{5/2}} \right) dx$

$$= \int e^x \left(\frac{1}{(1 + x^2)^{1/2}} - \frac{x}{(1 + x^2)^{3/2}} + \frac{x}{(1 + x^2)^{3/2}} + \frac{1 - 2x^2}{(1 + x^2)^{5/2}} \right) dx$$

$$= \frac{e^x}{(1 + x^2)^{1/2}} + \frac{xe^x}{(1 + x^2)^{3/2}} + c = \frac{e^x \{1 + x^2 + x\}}{(1 + x^2)^{3/2}} + c$$

Ex. 8 Evaluate $\int \frac{x^4}{(x+2)(x^2+1)} dx$

Sol. $\frac{x^4}{(x+2)(x^2+1)} = (x-2) + \frac{3x^2+4}{(x+2)(x^2+1)}$

Now, $\frac{3x^2+4}{(x+2)(x^2+1)} = \frac{16}{5(x+2)} + \frac{-\frac{1}{5}x + \frac{2}{5}}{x^2+1}$

So, $\frac{x^4}{(x+2)(x^2+1)} = x-2 + \frac{16}{5(x+2)} + \frac{-\frac{1}{5}x + \frac{2}{5}}{x^2+1}$

Now, $\int \left((x-2) + \frac{16}{5(x+2)} + \frac{-\frac{1}{5}x + \frac{2}{5}}{x^2+1} \right) dx$

$$= \frac{x^2}{2} - 2x + \frac{2}{5} \tan^{-1} x + \frac{16}{5} \ln|x+2| - \frac{1}{10} \ln(x^2+1) + c$$

Ex. 9 Evaluate : $\int \frac{1}{x^4 + 5x^2 + 1} dx$

Sol. $I = \frac{1}{2} \int \frac{2}{x^4 + 5x^2 + 1} dx$

$$\Rightarrow I = \frac{1}{2} \int \frac{1+x^2}{x^4 + 5x^2 + 1} dx + \frac{1}{2} \int \frac{1-x^2}{x^4 + 5x^2 + 1} dx = \frac{1}{2} \int \frac{1+1/x^2}{x^2+5+1/x^2} dx - \frac{1}{2} \int \frac{1-1/x^2}{x^2+5+1/x^2} dx$$

{dividing N^r and D^r by x^2 }

$$= \frac{1}{2} \int \frac{(1+1/x^2)}{(x-1/x)^2+7} dx - \frac{1}{2} \int \frac{(1-1/x^2)}{(x+1/x)^2+3} dx = \frac{1}{2} \int \frac{dt}{t^2+(\sqrt{7})^2} - \frac{1}{2} \int \frac{du}{u^2+(\sqrt{3})^2}$$

where $t = x - \frac{1}{x}$ and $u = x + \frac{1}{x}$

$$I = \frac{1}{2} \cdot \frac{1}{\sqrt{7}} \left(\tan^{-1} \frac{t}{\sqrt{7}} \right) - \frac{1}{2} \cdot \frac{1}{\sqrt{3}} \left(\tan^{-1} \frac{u}{\sqrt{3}} \right) + c \Rightarrow \frac{1}{2} \left[\frac{1}{\sqrt{7}} \tan^{-1} \left(\frac{x-1/x}{\sqrt{7}} \right) - \frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{x+1/x}{\sqrt{3}} \right) \right] + c$$

Ex. 10 Evaluate : $\int \left[\ln(\ln x) + \frac{1}{(\ln x)^2} \right] dx$

Sol. Let $I = \int \left[\ln(\ln x) + \frac{1}{(\ln x)^2} \right] dx$ {put $x = e^t \Rightarrow dx = e^t dt$ }

$$\begin{aligned} \therefore I &= \int e^t \left(\ln t + \frac{1}{t^2} \right) dt = \int e^t \left(\ln t - \frac{1}{t} + \frac{1}{t} + \frac{1}{t^2} \right) dt \\ &= e^t \left(\ln t - \frac{1}{t} \right) + C = x \left[\ln(\ln x) - \frac{1}{\ln x} \right] + C \end{aligned}$$

Ex. 11 Evaluate $\int \sin^2 x \cos^4 x dx$

Sol.
$$\begin{aligned} \int \sin^2 x \cos^4 x dx &= \int \left(\frac{1 - \cos 2x}{2} \right) \left(\frac{\cos 2x + 1}{2} \right)^2 dx = \int \frac{1}{8} (1 - \cos 2x) (\cos^2 2x + 2 \cos 2x + 1) dx \\ &= \frac{1}{8} \int (\cos^2 2x + 2 \cos 2x + 1 - \cos^3 2x - 2 \cos^2 2x - \cos 2x) dx \\ &= \frac{1}{8} \int (-\cos^3 2x - \cos^2 2x + \cos 2x + 1) dx = -\frac{1}{8} \int \left(\frac{\cos 6x + 3 \cos 2x}{4} + \frac{1 + \cos 4x}{2} - \cos 2x - 1 \right) dx \\ &= -\frac{1}{32} \left[\frac{\sin 6x}{6} + \frac{3 \sin 2x}{2} \right] - \frac{1}{16} x - \frac{\sin 4x}{64} + \frac{\sin 2x}{16} + \frac{x}{8} + c \\ &= -\frac{\sin 6x}{192} - \frac{\sin 4x}{64} + \frac{1}{64} \sin 2x + \frac{x}{16} + c \end{aligned}$$

Ex. 12 Evaluate : $\int \frac{dx}{2 + \sin^2 x}$

Sol. Divide numerator and denominator by $\cos^2 x$

$$I = \int \frac{\sec^2 x dx}{2 \sec^2 x + \tan^2 x} = \int \frac{\sec^2 x dx}{2 + 3 \tan^2 x}$$

Let $\sqrt{3} \tan x = t \quad \therefore \sqrt{3} \sec^2 x dx = dt$

So $I = \frac{1}{\sqrt{3}} \int \frac{dt}{2 + t^2} = \frac{1}{\sqrt{3}} \cdot \frac{1}{\sqrt{2}} \tan^{-1} \frac{t}{\sqrt{2}} + c = \frac{1}{\sqrt{6}} \tan^{-1} \left(\frac{\sqrt{3} \tan x}{\sqrt{2}} \right) + c$

Ex. 13 Evaluate : $\int \frac{1}{x^2 - x + 1} dx$

Sol.
$$\begin{aligned} \int \frac{1}{x^2 - x + 1} dx &= \int \frac{1}{x^2 - x + \frac{1}{4} - \frac{1}{4} + 1} dx = \int \frac{1}{(x - 1/2)^2 + 3/4} dx \\ &= \int \frac{1}{(x - 1/2)^2 + (\sqrt{3}/2)^2} dx = \frac{1}{\sqrt{3}/2} \tan^{-1} \left(\frac{x - 1/2}{\sqrt{3}/2} \right) + C \\ &= \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{2x - 1}{\sqrt{3}} \right) + C. \end{aligned}$$

Ex. 14 Evaluate $\int \frac{\sqrt{\sin x}}{\cos^{9/2} x} dx$

Sol. Let $I = \int \frac{\sin^{1/2} x}{\cos^{9/2} x} dx = \int \frac{dx}{\sin^{-1/2} x \cos^{9/2} x}$

Here $m + n = \frac{1}{2} - \frac{9}{2} = -4$ (negative even integer).

Divide Numerator & Denominator by $\cos^4 x$.

$$\begin{aligned} I &= \int \sqrt{\tan x} \sec^4 x dx = \int \sqrt{\tan x} (1 + \tan^2 x) \sec^2 x dx \\ &= \int \sqrt{t} (1 + t^2) dt \quad (\text{using } \tan x = t) \\ &= \frac{2}{3} t^{3/2} + \frac{2}{7} t^{7/2} + c = \frac{2}{3} \tan^{3/2} x + \frac{2}{7} \tan^{7/2} x + c \end{aligned}$$

Ex. 15 Evaluate $\int \frac{\cos^4 x dx}{\sin^3 x \{\sin^5 x + \cos^5 x\}^{3/5}}$

Sol. $I = \int \frac{\cos^4 x}{\sin^3 x \{\sin^5 x + \cos^5 x\}^{3/5}} dx = \int \frac{\cos^4 x}{\sin^6 x \{1 + \cot^5 x\}^{3/5}} dx = \int \frac{\cot^4 x \operatorname{cosec}^2 x dx}{(1 + \cot^5 x)^{3/5}}$

Put $1 + \cot^5 x = t$

$5 \cot^4 x \operatorname{cosec}^2 x dx = -dt$

$$= -\frac{1}{5} \int \frac{dt}{t^{3/5}} = -\frac{1}{2} t^{2/5} + c = -\frac{1}{2} (1 + \cot^5 x)^{2/5} + c$$

Ex. 16 Evaluate : $\int (\sin x)^{1/3} (\cos x)^{-7/3} dx$

Sol. Here $m + n = \frac{1}{3} - \frac{7}{3} = -2$ (a negative integer)

$$\therefore \int (\sin x)^{1/3} (\cos x)^{-7/3} dx = \int (\tan x)^{1/3} \frac{1}{\cos^2 x} dx \quad \{\text{put } \tan x = t \Rightarrow \sec^2 x dx = dt\}$$

$$= \int t^{1/3} dt = \frac{3}{4} t^{4/3} + C = \frac{3}{4} (\tan x)^{4/3} + C$$

Ex. 17 Evaluate $\int \frac{dx}{(1+x^2)\sqrt{1-x^2}}$

Sol. Let, $I = \int \frac{dx}{(1+x^2)\sqrt{1-x^2}}$

Put $x = \frac{1}{t}$, So that $dx = \frac{-1}{t^2} dt$

$$\therefore I = \int \frac{-1/t^2 dt}{(1+1/t^2)\sqrt{1-1/t^2}} = -\int \frac{tdt}{(t^2+1)\sqrt{t^2-1}}$$

again let, $t^2 = u$. So that $2t \, dt = du$.

$$= \frac{-1}{2} \int \frac{du}{(u+1)\sqrt{u-1}} \text{ which reduces to the form } \int \frac{dx}{P\sqrt{Q}} \text{ where both P and Q are linear so that}$$

we put $u-1 = z^2$ so that $du = 2z \, dz$

$$\therefore I = -\frac{1}{2} \int \frac{2z \, dz}{(z^2+1+1)\sqrt{z^2}} = -\int \frac{dz}{(z^2+2)}$$

$$I = -\frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{z}{\sqrt{2}} \right) + c$$

$$I = -\frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{u-1}}{\sqrt{2}} \right) + c = -\frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{t^2-1}}{\sqrt{2}} \right) + c = -\frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{1-x^2}}{\sqrt{2}x} \right) + c$$

Ex. 18 $\int \frac{dx}{\cos^6 x + \sin^6 x}$ is equal to -

Sol. Let $I = \int \frac{dx}{\cos^6 x + \sin^6 x} = \int \frac{\sec^6 x}{1 + \tan^6 x} dx = \int \frac{(1 + \tan^2 x)^2 \sec^2 x \, dx}{1 + \tan^6 x}$

If $\tan x = p$, then $\sec^2 x \, dx = dp$

$$\begin{aligned} \Rightarrow I &= \int \frac{(1+p^2)^2 dp}{1+p^6} = \int \frac{(1+p^2)}{p^4-p^2+1} dp = \int \frac{p^2 \left(1 + \frac{1}{p^2} \right)}{p^2 \left(p^2 + \frac{1}{p^2} - 1 \right)} dp \\ &= \int \frac{dk}{k^2+1} = \tan^{-1}(k) + c \quad \left(\text{where } p - \frac{1}{p} = k, \left(1 + \frac{1}{p^2} \right) dp = dk \right) \\ &= \tan^{-1} \left(p - \frac{1}{p} \right) + c = \tan^{-1} (\tan x - \cot x) + c = \tan^{-1} (-2 \cot 2x) + c \end{aligned}$$

Ex. 19 Evaluate: $\int \frac{1}{x^4+1} dx$

Sol. We have,

$$\begin{aligned} I &= \int \frac{1}{x^4+1} dx = \int \frac{\frac{1}{x^2}}{x^2 + \frac{1}{x^2}} dx = \frac{1}{2} \int \frac{\frac{2}{x^2}}{x^2 + \frac{1}{x^2}} dx \\ &= \frac{1}{2} \int \frac{1 + \frac{1}{x^2}}{x^2 + \frac{1}{x^2}} - \frac{1 - \frac{1}{x^2}}{x^2 + \frac{1}{x^2}} dx = \frac{1}{2} \int \frac{1 + \frac{1}{x^2}}{x^2 + \frac{1}{x^2}} dx - \frac{1}{2} \int \frac{1 - \frac{1}{x^2}}{x^2 + \frac{1}{x^2}} dx \end{aligned}$$

$$= \frac{1}{2} \int \frac{1 + \frac{1}{x^2}}{\left(x - \frac{1}{x}\right)^2 + 2} dx - \frac{1}{2} \int \frac{1 - \frac{1}{x^2}}{\left(x + \frac{1}{x}\right)^2 - 2} dx$$

Putting $x - \frac{1}{x} = u$ in 1st integral and $x + \frac{1}{x} = v$ in 2nd integral, we get

$$\begin{aligned} I &= \frac{1}{2} \int \frac{du}{u^2 + (\sqrt{2})^2} - \frac{1}{2} \int \frac{dv}{v^2 - (\sqrt{2})^2} \\ &= \frac{1}{2\sqrt{2}} \tan^{-1} \left(\frac{u}{\sqrt{2}} \right) - \frac{1}{2} \cdot \frac{1}{2\sqrt{2}} \bullet n \left| \frac{v - \sqrt{2}}{v + \sqrt{2}} \right| + C \\ &= \frac{1}{2\sqrt{2}} \tan^{-1} \left(\frac{x - 1/x}{\sqrt{2}} \right) - \frac{1}{4\sqrt{2}} \bullet n \left| \frac{x + 1/x - \sqrt{2}}{x + 1/x + \sqrt{2}} \right| + C \\ &= \frac{1}{2\sqrt{2}} \tan^{-1} \left(\frac{x^2 - 1}{\sqrt{2}x} \right) - \frac{1}{4\sqrt{2}} \bullet n \left| \frac{x^2 - \sqrt{2}x + 1}{x^2 + x\sqrt{2} + 1} \right| + C \end{aligned}$$

Ex. 20 Evaluate : $\int \frac{2 \sin 2x - \cos x}{6 - \cos^2 x - 4 \sin x} dx$

Sol. $I = \int \frac{2 \sin 2x - \cos x}{6 - \cos^2 x - 4 \sin x} dx = \int \frac{(4 \sin x - 1) \cos x}{6 - (1 - \sin^2 x) - 4 \sin x} dx = \int \frac{(4 \sin x - 1) \cos x}{\sin^2 x - 4 \sin x + 5} dx$

Put $\sin x = t$, so that $\cos x dx = dt$.

$$\therefore I = \int \frac{(4t - 1)dt}{t^2 - 4t + 5} \quad \dots\dots (i)$$

Now, let $(4t - 1) = \lambda(2t - 4) + \mu$

Comparing coefficients of like powers of t , we get

$$\begin{aligned} 2\lambda &= 4, -4\lambda + \mu = -1 \\ \lambda &= 2, \mu = 7 \end{aligned} \quad \dots\dots (ii)$$

$$\begin{aligned} \therefore I &= \int \frac{2(2t - 4) + 7}{t^2 - 4t + 5} dt \quad \{\text{using (i) and (ii)}\} \\ &= 2 \int \frac{2t - 4}{t^2 - 4t + 5} dt + 7 \int \frac{dt}{t^2 - 4t + 5} = 2 \log |t^2 - 4t + 5| + 7 \int \frac{dt}{t^2 - 4t + 4 - 4 + 5} \\ &= 2 \log |t^2 - 4t + 5| + 7 \int \frac{dt}{(t - 2)^2 + (1)^2} = 2 \log |t^2 - 4t + 5| + 7 \cdot \tan^{-1} (t - 2) + c \\ &= 2 \log |\sin^2 x - 4 \sin x + 5| + 7 \tan^{-1} (\sin x - 2) + c. \end{aligned}$$

Ex. 21 The value of $\int \sqrt{\frac{3-x}{3+x}} \cdot \sin^{-1} \left(\frac{1}{\sqrt{6}} \sqrt{3-x} \right) dx$, is equal to -

Sol. Here, $I = \int \sqrt{\frac{3-x}{3+x}} \cdot \sin^{-1} \left(\frac{1}{\sqrt{6}} \sqrt{3-x} \right) dx$

Put $x = 3\cos 2\theta \Rightarrow dx = -6\sin 2\theta d\theta$

$$= \int \sqrt{\frac{3-3\cos 2\theta}{3+3\cos 2\theta}} \cdot \sin^{-1} \left(\frac{1}{\sqrt{6}} \sqrt{3-3\cos 2\theta} \right) (-6 \sin 2\theta) d\theta$$

$$= \int \frac{\sin \theta}{\cos \theta} \cdot \sin^{-1} (\sin \theta) \cdot (-6 \sin 2\theta) d\theta = -6 \int \theta \cdot (2 \sin^2 \theta) d\theta$$

$$= -6 \int \theta (1 - \cos 2\theta) d\theta = -6 \left\{ \frac{\theta^2}{2} - \int \theta \cos 2\theta d\theta \right\}$$

$$= -6 \left\{ \frac{\theta^2}{2} - \left(\theta \frac{\sin 2\theta}{2} - \int 1 \cdot \left(\frac{\sin 2\theta}{2} \right) d\theta \right) \right\} = -3\theta^2 + 6 \left\{ \theta \frac{\sin 2\theta}{2} + \frac{\cos 2\theta}{4} \right\} + c$$

$$= \frac{1}{4} \left\{ -3 \left(\cos^{-1} \left(\frac{x}{3} \right) \right)^2 + 2\sqrt{9-x^2} \cdot \cos^{-1} \left(\frac{x}{3} \right) + 2x \right\} + c$$

Ex. 22 Evaluate : $\int \frac{\tan \left(\frac{\pi}{4} - x \right)}{\cos^2 x \sqrt{\tan^3 x + \tan^2 x + \tan x}} dx$

Sol. $I = \int \frac{\tan \left(\frac{\pi}{4} - x \right)}{\cos^2 x \sqrt{\tan^3 x + \tan^2 x + \tan x}} dx = \int \frac{(1 - \tan^2 x) dx}{(1 + \tan x)^2 \cos^2 x \sqrt{\tan^3 x + \tan^2 x + \tan x}}$

$$I = \int \frac{-\left(1 - \frac{1}{\tan^2 x}\right) \sec^2 x dx}{\left(\tan x + 2 + \frac{1}{\tan x}\right) \sqrt{\tan x + 1 + \frac{1}{\tan x}}}$$

let, $y = \sqrt{\tan x + 1 + \frac{1}{\tan x}} \Rightarrow 2y dy = \left(\sec^2 x - \frac{1}{\tan^2 x} \cdot \sec^2 x \right) dx$

$\therefore I = \int \frac{-2y dy}{(y^2 + 1)y} = -2 \int \frac{dy}{1 + y^2}$

$$= -2 \tan^{-1} y + c = -2 \tan^{-1} \left(\sqrt{\tan x + 1 + \frac{1}{\tan x}} \right) + c$$

Exercise # 1

[Single Correct Choice Type Questions]

- $\int 4 \sin x \cos \frac{x}{2} \cos \frac{3x}{2} dx$ is equal to -

(A) $\cos x - \frac{1}{2} \cos 2x + \frac{1}{3} \cos 3x + c$ (B) $\cos x - \frac{1}{2} \cos 2x - \frac{1}{3} \cos 3x + c$

(C) $\cos x + \frac{1}{2} \cos 2x + \frac{1}{3} \cos 3x + c$ (D) $\cos x + \frac{1}{2} \cos 2x - \frac{1}{3} \cos 3x + c$
- The value of $\int \frac{dx}{\sin x \cdot \sin(x+\alpha)}$ is equal to

(A) $\operatorname{cosec} \alpha \bullet n \left| \frac{\sin x}{\sin(x+\alpha)} \right| + C$ (B) $\operatorname{cosec} \alpha \bullet n \left| \frac{\sin(x+\alpha)}{\sin x} \right| + C$

(C) $\operatorname{cosec} \alpha \bullet n \left| \frac{\sec(x+\alpha)}{\sec x} \right| + C$ (D) $\operatorname{cosec} \alpha \bullet n \left| \frac{\sec x}{\sec(x+\alpha)} \right| + C$
- Which one of the following is TRUE.

(A) $x \cdot \int \frac{dx}{x} = x \ln |x| + C$ (B) $x \cdot \int \frac{dx}{x} = x \ln |x| + Cx$

(C) $\frac{1}{\cos x} \cdot \int \cos x \, dx = \tan x + C$ (D) $\frac{1}{\cos x} \cdot \int \cos x \, dx = x + C$
- $\int \frac{8x+13}{\sqrt{4x+7}} dx$ is equal to -

(A) $\frac{1}{6} (8x+11) \sqrt{4x+7} + c$ (B) $\frac{1}{6} (8x+13) \sqrt{4x+7} + c$

(C) $\frac{1}{6} (8x+9) \sqrt{4x+7} + c$ (D) $\frac{1}{6} (8x+15) \sqrt{4x+7} + c$
- $\int \frac{(2x+1)}{(x^2+4x+1)^{3/2}} dx$

(A) $\frac{x^3}{(x^2+4x+1)^{1/2}} + C$ (B) $\frac{x}{(x^2+4x+1)^{1/2}} + C$ (C) $\frac{x^2}{(x^2+4x+1)^{1/2}} + C$ (D) $\frac{1}{(x^2+4x+1)^{1/2}} + C$
- If $I = \int \frac{\sin x + \sin^3 x}{\cos 2x} dx = A \cos x + B \bullet n |f(x)| + C$, then

(A) $A = \frac{1}{4}, B = \frac{-1}{\sqrt{2}}, f(x) = \frac{\sqrt{2} \cos x - 1}{\sqrt{2} \cos x + 1}$ (B) $A = -\frac{1}{2}, B = \frac{-3}{4\sqrt{2}}, f(x) = \frac{\sqrt{2} \cos x - 1}{\sqrt{2} \cos x + 1}$

(C) $A = -\frac{1}{2}, B = \frac{3}{\sqrt{2}}, f(x) = \frac{\sqrt{2} \cos x + 1}{\sqrt{2} \cos x - 1}$ (D) $A = \frac{1}{2}, B = \frac{-3}{4\sqrt{2}}, f(x) = \frac{\sqrt{2} \cos x - 1}{\sqrt{2} \cos x + 1}$

7. $\int \frac{1-x^7}{x(1+x^7)} dx$ equals :
- (A) $\ln x + \frac{2}{7} \ln(1+x^7) + c$ (B) $\ln x - \frac{2}{7} \ln(1-x^7) + c$
 (C) $\ln x - \frac{2}{7} \ln(1+x^7) + c$ (D) $\ln x + \frac{2}{7} \ln(1-x^7) + c$
8. $\int \left(\frac{\cos^8 x - \sin^8 x}{1 - 2 \sin^2 x \cos^2 x} \right) dx$ equals -
- (A) $-\frac{\sin 2x}{2} + c$ (B) $\frac{\sin 2x}{2} + c$ (C) $\frac{\cos 2x}{2} + c$ (D) $-\frac{\cos 2x}{2} + c$
9. The value of $\int \left\{ \ln(1 + \sin x) + x \tan \left(\frac{\pi}{4} - \frac{x}{2} \right) \right\} dx$ is equal to:
- (A) $x \bullet \ln(1 + \sin x) + C$ (B) $\bullet \ln(1 + \sin x) + C$ (C) $-x \bullet \ln(1 + \sin x) + C$ (D) $\bullet \ln(1 - \sin x) + C$
10. Suppose $A = \int \frac{dx}{x^2 + 6x + 25}$ and $B = \int \frac{dx}{x^2 - 6x - 27}$.
 If $12(A + B) = \lambda \cdot \tan^{-1} \left(\frac{x+3}{4} \right) + \mu \cdot \ln \left| \frac{x-9}{x+3} \right| + C$, then the value of $(\lambda + \mu)$ is
- (A) 3 (B) 4 (C) 5 (D) 6
11. The value of $\int \sqrt{\frac{1 - \cos x}{\cos \alpha - \cos x}} dx$, where $0 < \alpha < x < \pi$, is equal to
- (A) $2 \bullet \left(\cos \frac{\alpha}{2} - \cos \frac{x}{2} \right) + C$ (B) $\sqrt{2} \bullet \left(\cos \frac{\alpha}{2} - \cos \frac{x}{2} \right) + C$
 (C) $2\sqrt{2} \bullet \left(\cos \frac{\alpha}{2} - \cos \frac{x}{2} \right) + C$ (D) $-2 \sin^{-1} \left(\frac{\cos \frac{x}{2}}{\cos \frac{\alpha}{2}} \right) + C$
12. Primitive of $\sqrt[3]{\frac{x}{(x^4 - 1)^4}}$ w.r.t. x is -
- (A) $\frac{3}{4} \left(1 + \frac{1}{x^4 - 1} \right)^{\frac{1}{3}} + c$ (B) $-\frac{3}{4} \left(1 + \frac{1}{x^4 - 1} \right)^{\frac{1}{3}} + c$ (C) $\frac{4}{3} \left(1 + \frac{1}{x^4 - 1} \right)^{\frac{1}{3}} + c$ (D) $-\frac{4}{3} \left(1 + \frac{1}{x^4 - 1} \right)^{\frac{1}{3}} + c$
13. The value of $\int [1 + \tan x \cdot \tan(x + \alpha)] dx$ is equal to
- (A) $\cos \alpha \cdot \bullet \left| \frac{\sin x}{\sin(x + \alpha)} \right| + C$ (B) $\tan \alpha \cdot \bullet \left| \frac{\sin x}{\sin(x + \alpha)} \right| + C$
 (C) $\cot \alpha \cdot \bullet \left| \frac{\sec(x + \alpha)}{\sec x} \right| + C$ (D) $\cot \alpha \cdot \bullet \left| \frac{\cos(x + \alpha)}{\cos x} \right| + C$

14. $\int \frac{x dx}{\sqrt{1+x^2} + \sqrt{(1+x^2)^3}}$ is equal to :
 (A) $\frac{1}{2} \ln(1 + \sqrt{1+x^2}) + c$ (B) $2\sqrt{1+\sqrt{1+x^2}} + c$ (C) $2(1 + \sqrt{1+x^2}) + c$ (D) none of these
15. If $f(x) = \int \frac{2\sin x - \sin 2x}{x^3} dx$, where $x \neq 0$, then $\lim_{x \rightarrow 0} f'(x)$ has the value
 (A) 0 (B) 1 (C) 2 (D) not defined
16. If $I_n = \int (\sin x)^n dx$ $n \in \mathbb{N}$
 Then $5I_4 - 6I_6$ is equal to
 (A) $\sin x \cdot (\cos x)^5 + C$ (B) $\sin 2x \cdot \cos 2x + C$
 (C) $\frac{\sin 2x}{8} [\cos^2 2x + 1 - 2 \cos 2x] + C$ (D) $\frac{\sin 2x}{8} [\cos^2 2x + 1 + 2 \cos 2x] + C$
17. $\int \frac{x dx}{\sqrt{1+x^2} + \sqrt{(1+x^2)^3}}$ is equal to -
 (A) $\frac{1}{2} \ln(1 + \sqrt{1+x^2}) + c$ (B) $2\sqrt{1+\sqrt{1+x^2}} + c$
 (C) $2(1 + \sqrt{1+x^2}) + c$ (D) none of these
18. The value of $\int \sqrt{\sec x - 1} dx$ is equal to
 (A) $2 \bullet n \left(\cos \frac{x}{2} + \sqrt{\cos^2 \frac{x}{2} - \frac{1}{2}} \right) + C$ (B) $\bullet n \left(\cos \frac{x}{2} + \sqrt{\cos^2 \frac{x}{2} - \frac{1}{2}} \right) + C$
 (C) $-2 \bullet n \left(\cos \frac{x}{2} + \sqrt{\cos^2 \frac{x}{2} - \frac{1}{2}} \right) + C$ (D) none of these
19. If $\int \frac{\cos x - \sin x + 1 - x}{e^x + \sin x + x} dx = \ln(f(x)) + g(x) + C$ where C is the constant of integration and $f(x)$ is positive, then $f(x) + g(x)$ has the value equal to
 (A) $e^x + \sin x + 2x$ (B) $e^x + \sin x$ (C) $e^x - \sin x$ (D) $e^x + \sin x + x$
20. $\int \frac{1 \ln|x|}{x\sqrt{1+\ln|x|}} dx$ equals -
 (A) $\frac{2}{3} \sqrt{1+\ln|x|} (\ln|x| - 2) + c$ (B) $\frac{2}{3} \sqrt{1+\ln|x|} (\ln|x| + 2) + c$
 (C) $\frac{1}{3} \sqrt{1+\ln|x|} (\ln|x| - 2) + c$ (D) $2\sqrt{1+\ln|x|} (3\ln|x| - 2) + c$

21. The value of $\int \frac{dx}{\cos^3 x \sqrt{\sin 2x}}$ is equal to

(A) $\sqrt{2} \left(\sqrt{\cos x} + \frac{1}{5} \tan^{5/2} x \right) + C$

(B) $\sqrt{2} \left(\sqrt{\tan x} + \frac{1}{5} \tan^{5/2} x \right) + C$

(C) $\sqrt{2} \left(\sqrt{\tan x} - \frac{1}{5} \tan^{5/2} x \right) + C$

(D) none of these

22. $\int x \cdot \frac{\ln(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx$ equals :

(A) $\sqrt{1+x^2} \ln(x + \sqrt{1+x^2}) - x + c$

(B) $\frac{x}{2} \cdot \ln^2(x + \sqrt{1+x^2}) - \frac{x}{\sqrt{1+x^2}} + c$

(C) $\frac{x}{2} \cdot \ln^2(x + \sqrt{1+x^2}) + \frac{x}{\sqrt{1+x^2}} + c$

(D) $\sqrt{1+x^2} \ln(x + \sqrt{1+x^2}) + x + c$

23. If $\int \frac{x^4 + 1}{x(x^2 + 1)^2} dx = A \ln|x| + \frac{B}{1+x^2} + c$, where c is the constant of integration then :

(A) $A = 1; B = -1$

(B) $A = -1; B = 1$

(C) $A = 1; B = 1$

(D) $A = -1; B = -1$

24. $\int \left(\frac{x}{1+x^5} \right)^{3/2} dx$ equals -

(A) $\frac{2}{5} \sqrt{\frac{x^5}{1+x^5}} + c$

(B) $\frac{2}{5} \sqrt{\frac{x}{1+x^5}} + c$

(C) $\frac{2}{5} \frac{1}{\sqrt{1+x^5}} + c$

(D) none of these

25. $\int \frac{(x^2 - 1) dx}{(x^4 + 3x^2 + 1) \tan^{-1}\left(\frac{x^2 + 1}{x}\right)} = \ln|f(x)| + C$ then $f(x)$ is

(A) $\ln\left(x + \frac{1}{x}\right)$

(B) $\tan^{-1}\left(x + \frac{1}{x}\right)$

(C) $\cot^{-1}\left(x + \frac{1}{x}\right)$

(D) $\ln\left(\tan^{-1}\left(x + \frac{1}{x}\right)\right)$

26. The value of $\int \sqrt{\frac{e^x - 1}{e^x + 1}} dx$ is equal to

(A) $\bullet_n \left(e^x + \sqrt{e^{2x} - 1} \right) - \sec^{-1}(e^x) + C$

(B) $\bullet_n \left(e^x + \sqrt{e^{2x} - 1} \right) + \sec^{-1}(e^x) + C$

(C) $\bullet_n \left(e^x - \sqrt{e^{2x} - 1} \right) - \sec^{-1}(e^x) + C$

(D) none of these

27. If $I_n = \int \cot^n x \, dx$, then $I_0 + I_1 + 2(I_2 + I_3 + \dots + I_8) + I_9 + I_{10}$ equals to :
(where $u = \cot x$)
- (A) $u + \frac{u^2}{2} + \dots + \frac{u^9}{9}$ (B) $-\left(u + \frac{u^2}{2} + \dots + \frac{u^9}{9}\right)$
(C) $-\left(u + \frac{u^2}{2!} + \dots + \frac{u^9}{9!}\right)$ (D) $\frac{u}{2} + \frac{2u^2}{3} + \dots + \frac{9u^9}{10}$
28. The value of $\int \frac{1}{\cos^6 x + \sin^6 x} \, dx$ is equal to
- (A) $\tan^{-1}(\tan x + \cot x) + C$ (B) $-\tan^{-1}(\tan x + \cot x) + C$
(C) $\tan^{-1}(\tan x - \cot x) + C$ (D) $-\tan^{-1}(\tan x - \cot x) + C$
29. Let $\int \frac{dx}{x^{2008} + x} = \frac{1}{p} \ln\left(\frac{x^q}{1+x^r}\right) + C$
where $p, q, r \in \mathbb{N}$ and need not be distinct, then the value of $(p + q + r)$ equals
- (A) 6024 (B) 6022 (C) 6021 (D) 6020
30. If $\int \frac{\sqrt{\cos^3 x}}{\sin^{11} x} \, dx = -2\left(A \tan^{\frac{-9}{2}} x + B \tan^{\frac{-5}{2}} x\right) + C$, then
- (A) $A = \frac{1}{9}, B = \frac{-1}{5}$ (B) $A = \frac{1}{9}, B = \frac{1}{5}$ (C) $A = -\frac{1}{9}, B = \frac{1}{5}$ (D) none of these
31. The integral $\int \sqrt{\cot x} e^{\sqrt{\sin x}} \sqrt{\cos x} \, dx$ equals
- (A) $\frac{\sqrt{\tan x} e^{\sqrt{\sin x}}}{\sqrt{\cos x}} + C$ (B) $2e^{\sqrt{\sin x}} + C$ (C) $-\frac{1}{2}e^{\sqrt{\sin x}} + C$ (D) $\frac{\sqrt{\cot x} e^{\sqrt{\sin x}}}{2\sqrt{\cos x}} + C$
32. $\int \frac{x^2 - 4}{x^4 + 24x^2 + 16} \, dx$ equals -
- (A) $\frac{1}{4} \tan^{-1}\left(\frac{(x^2 + 4)}{4x}\right) + c$ (B) $-\frac{1}{4} \cot^{-1}\left(\frac{(x^2 + 4)}{x}\right) + c$
(C) $-\frac{1}{4} \cot^{-1}\left(\frac{4(x^2 + 4)}{x}\right) + c$ (D) $\frac{1}{4} \cot^{-1}\left(\frac{(x^2 + 4)}{x}\right) + c$
33. $\int \frac{x^4 - 4}{x^2 \sqrt{4 + x^2 + x^4}} \, dx$ equals-
- (A) $\frac{\sqrt{4 + x^2 + x^4}}{x} + c$ (B) $\sqrt{4 + x^2 + x^4} + c$ (C) $\frac{\sqrt{4 + x^2 + x^4}}{2} + c$ (D) $\frac{\sqrt{4 + x^2 + x^4}}{2x} + c$

34. $\int \frac{x^2(1-\ln x)}{\ln^4 x - x^4} dx$ equals
- (A) $\frac{1}{2} \ln \left(\frac{x}{\ln x} \right) - \frac{1}{4} \ln (\ln^2 x - x^2) + C$ (B) $\frac{1}{4} \ln \left(\frac{\ln x - x}{\ln x + x} \right) - \frac{1}{2} \tan^{-1} \left(\frac{\ln x}{x} \right) + C$
- (C) $\frac{1}{4} \ln \left(\frac{\ln x + x}{\ln x - x} \right) + \frac{1}{2} \tan^{-1} \left(\frac{\ln x}{x} \right) + C$ (D) $\frac{1}{4} \left(\ln \left(\frac{\ln x - x}{\ln x + x} \right) + \tan^{-1} \left(\frac{\ln x}{x} \right) \right) + C$
35. If $\int \frac{4e^x + 6e^{-x}}{9e^x - 4e^{-x}} dx = Ax + B \bullet n |9e^{2x} - 4| + C$, then
- (A) $A = -\frac{3}{2}, B = \frac{35}{36}, C = 0$ (B) $A = \frac{35}{36}, B = -\frac{3}{2}, C \in \mathbb{R}$
- (C) $A = -\frac{3}{2}, B = \frac{35}{36}, C \in \mathbb{R}$ (D) $A = \frac{3}{2}, B = \frac{35}{36}, C \in \mathbb{R}$
36. If $\int \frac{(2x+3)dx}{x(x+1)(x+2)(x+3)+1} = C - \frac{1}{f(x)}$ where $f(x)$ is of the form of $ax^2 + bx + c$ then $(a+b+c)$ equals
- (A) 4 (B) 5 (C) 6 (D) none
37. $\int \frac{x^9}{(x^2+4)^6} dx$ is equal to -
- (A) $\frac{1}{5x} \left(4 + \frac{1}{x^2} \right)^{-5} + c$ (B) $\frac{1}{5} \left(4 + \frac{1}{x^2} \right)^{-5} + c$
- (C) $\frac{1}{10x} (1 + 4x^2)^{-5} + c$ (D) $\frac{1}{40} (1 + 4x^2)^{-5} + c$
38. $\int (\sin(101x) \cdot \sin^{99} x) dx$ equals
- (A) $\frac{\sin(100x)(\sin x)^{100}}{100} + C$ (B) $\frac{\cos(100x)(\sin x)^{100}}{100} + C$
- (C) $\frac{\cos(100x)(\cos x)^{100}}{100} + C$ (D) $\frac{\sin(100x)(\sin x)^{101}}{101} + C$
39. If $\int \frac{dx}{\sqrt{\sin^3 x \cos^5 x}} = a \sqrt{\cot x} + b \sqrt{\tan^3 x} + C$, where C is an arbitrary constant of integration, then the values of 'a' and 'b' are respectively:
- (A) -2 & $\frac{2}{3}$ (B) 2 & $-\frac{2}{3}$ (C) 2 & $\frac{2}{3}$ (D) none

Exercise # 2

Part # I

[Multiple Correct Choice Type Questions]

- Primitive of $\sqrt{1 + 2 \tan x (\sec x + \tan x)}$ w.r.t. x is -
 (A) $\ln|\sec x| - \ln|\sec x - \tan x| + c$ (B) $\ln|\sec x + \tan x| + \ln|\sec x| + c$
 (C) $2 \ln\left|\sec \frac{x}{2} + \tan \frac{x}{2}\right| + c$ (D) $\ln|1 + \tan x(\sec x + \tan x)| + c$
- If $\int e^{3x} \cos 4x \, dx = e^{3x} (A \sin 4x + B \cos 4x) + C$ then:
 (A) $4A = 3B$ (B) $2A = 3B$ (C) $3A = 4B$ (D) $4A + 3B = 1$
- Let $f(x) = 3x^2 \cdot \sin \frac{1}{x} - x \cos \frac{1}{x}$, $x \neq 0$, $f(0) = 0$, $f\left(\frac{1}{\pi}\right) = 0$, then which of the following is/are not correct.
 (A) $f(x)$ is continuous at $x = 0$ (B) $f(x)$ is non-differentiable at $x = 0$
 (C) $f'(x)$ is discontinuous at $x = 0$ (D) $f'(x)$ is differentiable at $x = 0$
- $\int \frac{\sin 2x}{\sin^4 x + \cos^4 x} dx$ is equal to -
 (A) $\cot^{-1}(\cot^2 x) + c$ (B) $-\cot^{-1}(\tan^2 x) + c$ (C) $\tan^{-1}(\tan^2 x) + c$ (D) $-\tan^{-1}(\cos 2x) + c$
- $\int \frac{\ln(\tan x)}{\sin x \cos x} dx$ equal:
 (A) $\frac{1}{2} \ln^2(\cot x) + c$ (B) $\frac{1}{2} \ln^2(\sec x) + c$
 (C) $\frac{1}{2} \ln^2(\sin x \sec x) + c$ (D) $\frac{1}{2} \ln^2(\cos x \operatorname{cosec} x) + c$
- The value of $\int 2^{mx} \cdot 3^{nx} dx$ (when $m, n \in \mathbb{N}$) is equal to :
 (A) $\frac{2^{mx} + 3^{nx}}{m \ln 2 + n \ln 3} + C$ (B) $\frac{e^{(m \ln 2 + n \ln 3)x}}{m \ln 2 + n \ln 3} + C$
 (C) $\frac{2^{mx} \cdot 3^{nx}}{\ln(2^m \cdot 3^n)} + C$ (D) $\frac{(mn) \cdot 2^x \cdot 3^x}{m \ln 2 + n \ln 3} + C$
- $\int \frac{dx}{x^3 \left(1 - \frac{1}{2x^2}\right)}$ equals-
 (A) $\ln|2x^2 - 1| + 2 \ln|x| + c$ (B) $\ln|2x^2 - 1| - 2 \ln|x| + c$
 (C) $\ln|2x^2 - 1| - \ln(x^2) - \ln 2 + c$ (D) $\ln\left|1 - \frac{1}{2x^2}\right| + c$

8. $\int \frac{1}{x^2-1} \ln \frac{x-1}{x+1} dx$ equals -
- (A) $\frac{1}{2} \ln^2 \left| \frac{x-1}{x+1} \right| + c$ (B) $\frac{1}{4} \ln^2 \left| \frac{x-1}{x+1} \right| + c$ (C) $\frac{1}{2} \ln^2 \left| \frac{x+1}{x-1} \right| + c$ (D) $\frac{1}{4} \ln^2 \left| \frac{x+1}{x-1} \right| + c$
9. If $\int e^u \cdot \sin 2x dx$ can be found in terms of known functions of x then u can be:
- (A) x (B) $\sin x$ (C) $\cos x$ (D) $\cos 2x$
10. If $\int \frac{dx}{5+4\cos x} = I \tan^{-1} \left(m \tan \frac{x}{2} \right) + C$ then :
- (A) $I = 2/3$ (B) $m = 1/3$ (C) $I = 1/3$ (D) $m = 2/3$
11. $\int \sin 2x dx$ equals -
- (A) $-\frac{\cos 2x}{2} + c$ (B) $\frac{\sin^2 x}{2} + c$ (C) $-\frac{\cos^2 x}{2} + c$ (D) $\frac{\cos 2x}{2} + c$
12. The value of $\int \frac{x^2 + \cos^2 x}{1+x^2} \operatorname{cosec}^2 x dx$ is equal to:
- (A) $\cot x - \cot^{-1} x + C$ (B) $C - \cot x + \cot^{-1} x$
- (C) $-\tan^{-1} x - \frac{\operatorname{cosec} x}{\sec x} + C$ (D) $-e^{\lambda \tan^{-1} x} - \cot x + C$
13. If $I_n = \int \cot^n x dx$ and $I_0 + I_1 + 2(I_2 + \dots + I_8) + I_9 + I_{10} = A \left(u + \frac{u^2}{2} + \dots + \frac{u^9}{9} \right) + C$, where $u = \cot x$ and C is an arbitrary constant, then
- (A) A is constant (B) $A = -1$ (C) $A = 1$ (D) A is dependent on x
14. Suppose $J = \int \frac{\sin^2 x + \sin x}{1 + \sin x + \cos x} dx$ and $K = \int \frac{\cos^2 x + \cos x}{1 + \sin x + \cos x} dx$. If C is an arbitrary constant of integration then which of the following is/are correct?
- (A) $J = \frac{1}{2} (x - \sin x + \cos x) + C$ (B) $J = K - (\sin x + \cos x) + C$
- (C) $J = x - K + C$ (D) $K = \frac{1}{2} (x - \sin x + \cos x) + C$
15. $\int \frac{dx}{\sqrt{x-x^2}}$ equals, where $x \in \left(\frac{1}{2}, 1 \right)$ -
- (A) $2 \sin^{-1} \sqrt{x} + c$ (B) $\sin^{-1} (2x-1) + c$ (C) $c - \cos^{-1} (2x-1)$ (D) $\cos^{-1} 2\sqrt{x-x^2} + c$

In each of the following questions, a statement of Assertion (A) is given followed by a corresponding statement of Reason (R) just below it. Of the statements mark the correct answer as

- (A) Statement-I is True, Statement-II is True ; Statement-II is a correct explanation for Statement-I
 (B) Statement-I is True, Statement-II is True ; Statement-II is NOT a correct explanation for Statement-I
 (C) Statement-I is True, Statement-II is False.
 (D) Statement-I is False, Statement-II is True.

1. If $D(x) = \begin{vmatrix} f_1(x) & f_2(x) & f_3(x) \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$, where f_1, f_2, f_3 are differentiable function and $a_2, b_2, c_2, a_3, b_3, c_3$ are constants.

Statement - I $\int D(x)dx = \begin{vmatrix} \int f_1(x)dx & \int f_2(x)dx & \int f_3(x)dx \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} + c$

Statement - II Integration of sum of several function is equal to sum of integration of individual functions.

2. **Statement - I** $\int \frac{\ln(e^x + 1)}{e^x} dx = x - \left(\frac{1 + e^x}{e^x} \right) \bullet n(e^x + 1) + C$

Statement - II $\int \frac{f'(x)}{f(x)} dx = \bullet n |f(x)| + C$

3. **Statement - I** If $a > 0$ and $b^2 - 4ac < 0$, then the values of integral $\int \frac{dx}{ax^2 + bx + c}$ will be of the type

$\mu \tan^{-1} \frac{x+A}{B} + c$, where A, B, C, μ are constants.

Statement - II If $a > 0$, $b^2 - 4ac < 0$, then $ax^2 + bx + c$ can be written as sum of two squares.

4. **Statement - I** The function $F(x) = \int \frac{x}{(x-1)(x^2+1)} dx$ is discontinuous at $x = 1$

Statement - II If $F(x) = \int f(x) dx$ and $f(x)$ is discontinuous at $x = a$ then $F(x)$ is also discontinuous at $x = a$.

5. **Statement - I** If $x > 0, x \neq 1$ then $\int (\log_x e - (\log_x e)^2) dx = x \log_x e + C$

Statement - II $\int e^x (f(x) + f'(x)) dx = e^x f(x) + C$ and $e^t = x$ iff $t = \bullet nx$

6. **Statement - I** $\int (\sin x)^5 \cos x dx = \frac{\sin^6 x}{6} + C$

Statement - II $\int (f(x))^n f'(x) dx = \frac{(f(x))^{n+1}}{n+1} + C, n \in I$

7. If y is a function of x such that $y(x - y)^2 = x$.

Statement - I : $\int \frac{dx}{x-3y} = \frac{1}{2} \log[(x-y)^2 - 1]$

Statement - II : $\int \frac{dx}{x-3y} = \log(x - 3y) + c$.

Exercise # 3

Part # I

[Matrix Match Type Questions]

Following question contains statements given in two columns, which have to be matched. The statements in **Column-I** are labelled as A, B, C and D while the statements in **Column-II** are labelled as p, q, r and s. Any given statement in **Column-I** can have correct matching with one or more statement(s) in **Column-II**.

1. The antiderivative of

Column-I

(A) $f(x) = \frac{1}{(a^2 + b^2) - (a^2 - b^2)\cos x}$ is

(B) $f(x) = \frac{1}{a^2 \sin^2 x + b^2 \cos^2 x}$ is

(C) $f(x) = \frac{1}{a \cos x + b \sin x}$ is

(D) $f(x) = \frac{1}{a^2 - b^2 \cos^2 x}$ is ; ($a^2 > b^2$)

Column-II

(p) $\frac{1}{ab} \tan^{-1} \left(\frac{a}{b} \tan \frac{x}{2} \right) + c$

(q) $\frac{1}{a^2 \sin \alpha} \tan^{-1} \left(\frac{\tan x}{\sin \alpha} \right) + c, \alpha = \cos^{-1} \frac{b}{a}$

(r) $\frac{1}{ab} \tan^{-1} \left(\frac{a}{b} \tan x \right) + c$

(s) $\frac{1}{\sqrt{a^2 + b^2}} \log \left| \tan \frac{1}{2} \left(x + \tan^{-1} \frac{a}{b} \right) \right| + c$

2. If $I = \int \frac{dx}{a + b \cos x}$, where $a, b > 0$ and $a + b = u, a - b = v$, then match the following column

Column-I

(A) $v = 0$

(B) $v > 0$

(C) $v < 0$

Column-II

(p) $I = \frac{1}{\sqrt{uv}} \bullet \frac{\left| \frac{\sqrt{u} + \sqrt{v} \tan \frac{x}{2}}{\sqrt{u} - \sqrt{v} \tan \frac{x}{2}} \right|}{\left| \frac{\sqrt{u} + \sqrt{v} \tan \frac{x}{2}}{\sqrt{u} - \sqrt{v} \tan \frac{x}{2}} \right|} + C$

(q) $I = \frac{2}{\sqrt{uv}} \tan^{-1} \left(\sqrt{\frac{v}{u}} \tan \frac{x}{2} \right) + C$

(r) $I = \frac{1}{\sqrt{-uv}} \bullet \frac{\left| \frac{\sqrt{u} + \sqrt{-v} \tan \frac{x}{2}}{\sqrt{u} - \sqrt{-v} \tan \frac{x}{2}} \right|}{\left| \frac{\sqrt{u} + \sqrt{-v} \tan \frac{x}{2}}{\sqrt{u} - \sqrt{-v} \tan \frac{x}{2}} \right|} + C$

(s) $\frac{2}{u} \tan \frac{x}{2} + C$

- 3.

Column-I

(A) Let $f(x) = \int x^{\sin x} (1 + x \cos x \cdot \ln x + \sin x) dx$ and $f\left(\frac{\pi}{2}\right) = \frac{\pi^2}{4}$ then the value of $f(\pi)$ is

(B) Let $g(x) = \int \frac{1 + 2 \cos x}{(\cos x + 2)^2} dx$ and $g(0) = 0$ then the value of $g\left(\frac{\pi}{2}\right)$ is

Column-II

(p) rational

(q) irrational

(r) integral

(C) If real numbers x and y satisfy $(x+5)^2 + (y-12)^2 = (14)^2$ then the minimum value of $\sqrt{(x^2 + y^2)}$ is

(s) prime

(D) Let $k(x) = \int \frac{(x^2+1)dx}{\sqrt[3]{x^3+3x+6}}$ and $k(-1) = \frac{1}{\sqrt[3]{2}}$ then the value of $k(-2)$ is

4. Column-I

(A) If $F(x) = \int \frac{x + \sin x}{1 + \cos x} dx$ and $F(0) = 0$, then the value of $F(\pi/2)$ is (p)

$\frac{\pi}{2}$

(B) Let $F(x) = \int e^{\sin^{-1} x} \left(1 - \frac{x}{\sqrt{1-x^2}} \right) dx$ and $F(0) = 1$,

(q) $\frac{\pi}{3}$

If $F(1/2) = \frac{k\sqrt{3}e^{\pi/6}}{\pi}$, then the value of k is

(C) Let $F(x) = \int \frac{dx}{(x^2+1)(x^2+9)}$ and $F(0) = 0$,

(r) $\frac{\pi}{4}$

if $F(\sqrt{3}) = \frac{5}{36} k$, then the value of k is

(D) Let $F(x) = \int \frac{\sqrt{\tan x}}{\sin x \cos x} dx$ and $F(0) = 0$

(s) π

if $F(\pi/4) = \frac{2k}{\pi}$, then the value of k is

5. $\int f(x)dx$ when

Column-I

(A) $f(x) = \frac{1}{(a^2 + x^2)^{3/2}}$

(B) $f(x) = \frac{x^2}{\sqrt{a^2 - x^2}}$

(C) $f(x) = \frac{1}{(x^2 - a^2)^{3/2}}$

(D) $f(x) = \frac{1}{x\sqrt{x^2 - a^2}}$

Column-II

(p) $c - \frac{1}{a} \sin^{-1} \frac{a}{|x|}$

(q) $\frac{a^2}{2} \sin^{-1} \frac{x}{a} - \frac{x}{2} \sqrt{a^2 - x^2} + c$

(r) $c - \frac{x}{a^2 \sqrt{x^2 - a^2}}$

(s) $\frac{x}{a^2 \sqrt{x^2 + a^2}} + c$

Comprehension # 1

In calculating a number of integrals we had to use the method of integration by parts several times in succession. The result could be obtained more rapidly and in a more concise form by using the so-called generalized formula for integration by parts

$$\int u(x)v(x)dx = u(x)v_1(x) - u'(x)v_2(x) + u''(x)v_3(x) - \dots + (-1)^{n-1}u^{(n-1)}(x)v_n(x) - (-1)^{n-1} \int u^{(n)}(x)v_n(x)dx$$

$$\text{where } v_1(x) = \int v(x)dx, v_2(x) = \int v_1(x)dx, \dots, v_n(x) = \int v_{n-1}(x)dx$$

Of course, we assume that all derivatives and integrals appearing in this formula exist. The use of the generalized formula for integration by parts is especially useful when calculating $\int P_n(x) Q(x) dx$, where $P_n(x)$, is polynomial of degree n and the factor $Q(x)$ is such that it can be integrated successively $n + 1$ times.

- If $\int (x^3 - 2x^2 + 3x - 1)\cos 2x dx = \frac{\sin 2x}{4} u(x) + \frac{\cos 2x}{8} v(x) + c$, then -

(A) $u(x) = x^3 - 4x^2 + 3x$ (B) $u(x) = 2x^3 - 4x^2 + 3x$
 (C) $v(x) = 3x^2 - 4x + 3$ (D) $v(x) = 6x^2 - 8x$
- If $\int e^{2x} \cdot x^4 dx = \frac{e^{2x}}{2} f(x) + C$ then $f(x)$ is equal to -

(A) $\left(x^4 - 2x^3 + 3x^2 - 3x + \frac{3}{2}\right) \frac{1}{2}$ (B) $x^4 - x^3 + 2x^2 - 3x + 2$
 (C) $x^4 - 2x^3 + 3x^2 - 3x + \frac{3}{2}$ (D) $x^4 - 2x^3 + 2x^2 - 3x + \frac{3}{2}$

Comprehension # 2

It is known that

$$\sqrt{\tan x} + \sqrt{\cot x} = \begin{cases} \frac{\sqrt{\sin x}}{\sqrt{\cos x}} + \frac{\sqrt{\cos x}}{\sqrt{\sin x}} & \text{if } 0 < x < \frac{\pi}{2} \\ \frac{\sqrt{-\sin x}}{\sqrt{-\cos x}} + \frac{\sqrt{-\cos x}}{\sqrt{-\sin x}} & \text{if } \pi < x < \frac{3\pi}{2} \end{cases},$$

$$\frac{d}{dx} (\sqrt{\tan x} - \sqrt{\cot x}) = \frac{1}{2} (\sqrt{\tan x} + \sqrt{\cot x}) (\tan x + \cot x), \forall x \in \left(0, \frac{\pi}{2}\right) \cup \left(\pi, \frac{3\pi}{2}\right)$$

and $\frac{d}{dx} (\sqrt{\tan x} + \sqrt{\cot x}) = \frac{1}{2} (\sqrt{\tan x} - \sqrt{\cot x}) (\tan x + \cot x), \forall x \in \left(0, \frac{\pi}{2}\right) \cup \left(\pi, \frac{3\pi}{2}\right).$

1. Value of integral $I = \int (\sqrt{\tan x} + \sqrt{\cot x}) dx$, where $x \in \left(0, \frac{\pi}{2}\right) \cup \left(\pi, \frac{3\pi}{2}\right)$ is
- (A) $\sqrt{2} \tan^{-1} \left(\frac{\sqrt{\tan x} - \sqrt{\cot x}}{\sqrt{2}} \right) + C$ (B) $\sqrt{2} \tan^{-1} \left(\frac{\sqrt{\tan x} + \sqrt{\cot x}}{\sqrt{2}} \right) + C$
- (C) $-\sqrt{2} \tan^{-1} \left(\frac{\sqrt{\tan x} - \sqrt{\cot x}}{\sqrt{2}} \right) + C$ (D) $-\sqrt{2} \tan^{-1} \left(\frac{\sqrt{\tan x} + \sqrt{\cot x}}{\sqrt{2}} \right) + C$
2. Value of the integral $I = \int (\sqrt{\tan x} + \sqrt{\cot x}) dx$, where $x \in \left(0, \frac{\pi}{2}\right)$, is
- (A) $\sqrt{2} \sin^{-1} (\cos x - \sin x) + C$ (B) $\sqrt{2} \sin^{-1} (\sin x - \cos x) + C$
- (C) $\sqrt{2} \sin^{-1} (\sin x + \cos x) + C$ (D) $-\sqrt{2} \sin^{-1} (\sin x + \cos x) + C$
3. Value of the integral $I = \int (\sqrt{\tan x} + \sqrt{\cot x}) dx$, where $x \in \left(\pi, \frac{3\pi}{2}\right)$, is
- (A) $\sqrt{2} \sin^{-1} (\cos x - \sin x) + C$ (B) $\sqrt{2} \sin^{-1} (\sin x - \cos x) + C$
- (C) $\sqrt{2} \sin^{-1} (\sin x + \cos x) + C$ (D) $-\sqrt{2} \sin^{-1} (\sin x + \cos x) + C$

Comprehension # 3

Let $I_{n,m} = \int \sin^n x \cos^m x dx$. Then we can relate $I_{n,m}$ with each of the following

- (i) $I_{n-2,m}$ (ii) $I_{n+2,m}$ (iii) $I_{n,m-2}$
- (iv) $I_{n,m+2}$ (v) $I_{n-2,m+2}$ (vi) $I_{n+2,m-2}$

Suppose we want to establish a relation between $I_{n,m}$ and $I_{n,m-2}$, then we set

$$P(x) = \sin^{n+1} x \cos^{m-1} x \quad \dots\dots\dots (1)$$

In $I_{n,m}$ and $I_{n,m-2}$ the exponent of $\cos x$ is m and $m-2$ respectively, the minimum of the two is $m-2$, adding 1 to the minimum we get $m-2+1 = m-1$. Now choose the exponent $m-1$ of $\cos x$ in $P(x)$. Similarly choose the exponent of $\sin x$ for $P(x)$

Now differentiating both sides of (1), we get

$$\begin{aligned} P'(x) &= (n+1) \sin^n x \cos^m x - (m-1) \sin^{n+2} x \cos^{m-2} x \\ &= (n+1) \sin^n x \cos^m x - (m-1) \sin^n x (1 - \cos^2 x) \cos^{m-2} x \\ &= (n+1) \sin^n x \cos^m x - (m-1) \sin^n x \cos^{m-2} x + (m-1) \sin^n x \cos^m x \\ &= (n+m) \sin^n x \cos^m x - (m-1) \sin^n x \cos^{m-2} x \end{aligned}$$

Now integrating both sides, we get

$$\sin^{n+1} x \cos^{m-1} x = (n+m) I_{n,m} - (m-1) I_{n,m-2}.$$

Similarly we can establish the other relations.

1. The relation between $I_{4,2}$ and $I_{2,2}$ is

(A) $I_{4,2} = \frac{1}{6} (-\sin^3 x \cos^3 x + 3I_{2,2})$

(B) $I_{4,2} = \frac{1}{6} (\sin^3 x \cos^3 x + 3I_{2,2})$

(C) $I_{4,2} = \frac{1}{6} (\sin^3 x \cos^3 x - 3I_{2,2})$

(D) $I_{4,2} = \frac{1}{4} (-\sin^3 x \cos^3 x + 2I_{2,2})$

2. The relation between $I_{4,2}$ and $I_{6,2}$ is

(A) $I_{4,2} = \frac{1}{5} (\sin^5 x \cos^3 x + 8I_{6,2})$

(B) $I_{4,2} = \frac{1}{5} (-\sin^5 x \cos^3 x + 8I_{6,2})$

(C) $I_{4,2} = \frac{1}{5} (\sin^5 x \cos^3 x - 8I_{6,2})$

(D) $I_{4,2} = \frac{1}{6} (\sin^5 x \cos^3 x + 8I_{6,2})$

3. The relation between $I_{4,2}$ and $I_{4,4}$ is

(A) $I_{4,2} = \frac{1}{3} (\sin^5 x \cos^3 x + 8I_{4,4})$

(B) $I_{4,2} = \frac{1}{3} (-\sin^5 x \cos^3 x + 8I_{4,4})$

(C) $I_{4,2} = \frac{1}{3} (\sin^5 x \cos^3 x - 8I_{4,4})$

(D) $I_{4,2} = \frac{1}{3} (\sin^5 x \cos^3 x + 6I_{4,4})$

Comprehension # 4

Integrals of class of functions following a definite pattern can be found by the method of reduction and recursion. Reduction formulas make it possible to reduce an integral dependent on the index $n > 0$, called the order of the integral, to an integral of the same type with a smaller index. Integration by parts helps us to derive reduction formulas. (Add a constant in each question)

1. If $I_n = \int \frac{dx}{(x^2 + a^2)^n}$ then $I_{n+1} + \frac{1-2n}{2n} \cdot \frac{1}{a^2} I_n$ is equal to -

(A) $\frac{x}{(x^2 + a^2)^n}$

(B) $\frac{1}{2na^2} \cdot \frac{1}{(x^2 + a^2)^{n-1}}$

(C) $\frac{1}{2na^2} \cdot \frac{x}{(x^2 + a^2)^n}$

(D) $\frac{1}{2na^2} \cdot \frac{1}{(x^2 + a^2)}$

2. If $I_{n,-m} = \int \frac{\sin^n x}{\cos^m x} dx$ then $I_{n,-m} + \frac{n-1}{m-1} I_{n-2,-m}$ is equal to-

(A) $\frac{\sin^{n-1} x}{\cos^{m-1} x}$

(B) $\frac{1}{(m-1)} \frac{\sin^{n-1} x}{\cos^{m-1} x}$

(C) $\frac{1}{(n-1)} \frac{\sin^{n-1} x}{\cos^{m-1} x}$

(D) $\frac{n-1}{m-1} \frac{\sin^{n-1} x}{\cos^{m-1} x}$

3. If $u_n = \int \frac{x^n}{\sqrt{ax^2 + 2bx + c}} dx$, then $(n+1)au_{n+1} + (2n+1)bu_n + ncu_{n-1}$ is equal to -

(A) $x^{n-1} \sqrt{ax^2 + 2bx + c}$

(B) $\frac{x^{n-2}}{\sqrt{ax^2 + 2bx + c}}$

(C) $\frac{x^n}{\sqrt{ax^2 + 2bx + c}}$

(D) $x^n \sqrt{ax^2 + 2bx + c}$

Exercise # 4

[Subjective Type Questions]

1. $\int \frac{dx}{\sin(x-a)\sin(x-b)}$
2. $\int \sqrt{x + \sqrt{x^2 + 2}} dx$
3. $\int \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}} dx$
4. $\int \frac{x \ln x}{(x^2 - 1)^{3/2}} dx$
5. $\int \frac{5x^4 + 4x^5}{(x^5 + x + 1)^2} dx$
6. $\int \sqrt{\frac{\sin(x-a)}{\sin(x+a)}} dx$
7. Find the value of $\int \frac{d(x^2+1)}{\sqrt{(x^2+2)}}$.
8. $\int \frac{\cot x dx}{(1 - \sin x)(\sec x + 1)}$
9. $\int \frac{x^3 + 3x + 2}{(x^2 + 1)^2(x+1)} dx$
10. $\int \frac{\sqrt{\cot x} - \sqrt{\tan x}}{1 + 3 \sin 2x} dx$
11. $\int \left[\frac{\sqrt{x^2 + 1} [\ln(x^2 + 1) - 2 \ln x]}{x^4} \right] dx$
12. $\int \frac{dx}{\sin x \sqrt{\sin(2x + \alpha)}}$
13. $\int \frac{x^2}{(x \sin x + \cos x)^2} dx$
14. $\int \frac{e^x (2 - x^2)}{(1 - x)\sqrt{1 - x^2}} dx$
15. $\int x \sqrt{\frac{a^2 - x^2}{a^2 + x^2}} dx$
16. Integrate $\frac{1}{2} f'(x)$ w.r.t. x^4 , where $f(x) = \tan^{-1} x + \ln \sqrt{1+x} - \ln \sqrt{1-x}$
17. $\int \frac{\ln(\cos x + \sqrt{\cos 2x})}{\sin^2 x} dx$
18. $\int \frac{dx}{x^2(x^4 + 1)^{3/4}}$
19. $\int \frac{(\cos 2x)^{1/2}}{\sin x} dx$
20. The antiderivative of $f(x) = \ln(\ln x) + (\ln x)^{-2}$ whose graph passes through (e, e) is $x \ln(\ln x) - x(\ln x)^{-1} + 2$.
21. $\int \sqrt{\frac{\operatorname{cosec} x - \cot x}{\operatorname{cosec} x + \cot x}} \cdot \frac{\sec x}{\sqrt{1 + 2 \sec x}} dx$
22. Let $\begin{bmatrix} 1 & 0 & 0 \\ 6 & 2 & 0 \\ 5 & 4 & 3 \end{bmatrix} \begin{bmatrix} x \\ x^2 \\ 1 \end{bmatrix} = \begin{bmatrix} x \\ 2\alpha x + \beta x^2 \\ 5x + \gamma x^2 + 3 \end{bmatrix}$, $\forall x \in \mathbb{R}$ and $f(x)$ is a differentiable function satisfying, $f(xy) = f(x) + x^2(y^2 - 1) + x(y - 1)$; $\forall x, y \in \mathbb{R}$ and $f(1) = 3$. Evaluate $\int \frac{\alpha x^2 + \beta x + \gamma}{f(x)} dx$

23. $\int \frac{dx}{\sin^2 x + \sin 2x}$
24. $\int \frac{(ax^2 - b) dx}{x\sqrt{c^2 x^2 - (ax^2 + b)^2}}$
25. $\int (\sin x)^{-11/3} (\cos x)^{-1/3} dx$
26. $\int \sin^{-1} \sqrt{\frac{x}{a+x}} dx$
27. $\int \frac{2 \sin 2\phi - \cos \phi}{6 - \cos^2 \phi - 4 \sin \phi} d\phi$
28. $\int \frac{1}{\sqrt[3]{x} + \sqrt[4]{x}} dx$
29. Let $f(x)$ is a quadratic function such that $f(0) = 1$ and $\int \frac{f(x)dx}{x^2(x+1)^3}$ is a rational function, find the value of $f'(0)$
30. $\int \cos 2\theta \ln \frac{\cos \theta + \sin \theta}{\cos \theta - \sin \theta} d\theta$
31. $\int \left[\left(\frac{x}{e} \right)^x + \left(\frac{e}{x} \right)^x \right] \ln x dx$
32. $\int \frac{x}{(7x - 10 - x^2)^{3/2}} dx$
33. $\int \frac{1 + x \cos x}{x(1 - x^2 e^{2 \sin x})} dx$
34. $\int \frac{e^{\cos x} (x \sin^3 x + \cos x)}{\sin^2 x} dx$
35. $\int (\sqrt{\tan x} + \sqrt{\cot x}) dx$

Exercise # 5

Part # I > [Previous Year Questions] [AIEEE/JEE-MAIN]

1. $\int \frac{\cos 2x - 1}{\cos 2x + 1} dx =$ [AIEEE-2002]

- (1) $\tan x - x + C$ (2) $x + \tan x + C$ (3) $x - \tan x + C$ (4) $-x - \cot x + C$

2. $\int \frac{(\log x)}{x^2} dx$ [AIEEE-2002]

- (1) $\frac{1}{2} (\log x + 1) + C$ (2) $-\frac{1}{x} (\log x + 1) + C$ (3) $\frac{1}{x} (\log x - 1) + C$ (4) $\log(x + 1) + C$

3. If $\int \frac{\sin x}{\sin(x - \alpha)} dx = Ax + B \log \sin(x - \alpha) + C$ then values of (A, B) is - [AIEEE-2004]

- (1) $(\sin \alpha, \cos \alpha)$ (2) $(\cos \alpha, \sin \alpha)$ (3) $(-\sin \alpha, \cos \alpha)$ (4) $(-\cos \alpha, \sin \alpha)$

4. $\int \frac{dx}{\cos x - \sin x}$ is equal to- [AIEEE-2004]

- (1) $\frac{1}{\sqrt{2}} \log \left| \tan \left(\frac{x}{2} - \frac{\pi}{8} \right) \right| + C$ (2) $\frac{1}{\sqrt{2}} \log \left| \cot \left(\frac{x}{2} \right) \right| + C$
(3) $\frac{1}{\sqrt{2}} \log \left| \tan \left(\frac{x}{2} - \frac{3\pi}{8} \right) \right| + C$ (4) $\frac{1}{\sqrt{2}} \log \left| \tan \left(\frac{x}{2} + \frac{3\pi}{8} \right) \right| + C$

5. $\int \left\{ \frac{(\log x - 1)}{1 + (\log x)^2} \right\}^2 dx$ is equals to - [AIEEE-2005]

- (1) $\frac{\log x}{(\log x)^2 + 1} + C$ (2) $\frac{x}{x^2 + 1} + C$ (3) $\frac{xe^x}{1 + x^2} + C$ (4) $\frac{x}{(\log x)^2 + 1} + C$

6. $\int \frac{dx}{\cos x + \sqrt{3} \sin x}$ equals- [AIEEE-2007]

- (1) $\frac{1}{2} \log \tan \left(\frac{x}{2} + \frac{\pi}{12} \right) + C$ (2) $\frac{1}{2} \log \tan \left(\frac{x}{2} - \frac{\pi}{12} \right) + C$
(3) $\log \tan \left(\frac{x}{2} + \frac{\pi}{12} \right) + C$ (4) $\log \tan \left(\frac{x}{2} - \frac{\pi}{12} \right) + C$

7. The value of $\sqrt{2} \int \frac{\sin x dx}{\sin \left(x - \frac{\pi}{4} \right)}$ is - [AIEEE-2008]

- (1) $x + \log \left| \cos \left(x - \frac{\pi}{4} \right) \right| + c$ (2) $x - \log \left| \sin \left(x - \frac{\pi}{4} \right) \right| + c$ (3) $x + \log \left| \sin \left(x - \frac{\pi}{4} \right) \right| + c$ (4) $x - \log \left| \cos \left(x - \frac{\pi}{4} \right) \right| + c$

8. If the integral $\int \frac{5 \tan x}{\tan x - 2} dx = x + a \ln|\sin x - 2 \cos x| + k$ then a is equal to : [AIEEE-2012]

- (1) 2 (2) -1 (3) -2 (4) 1

9. If $\int f(x) dx = \Psi(x)$, then $\int x^5 f(x^3) dx$ is equal to : [JEE Main 2013]

- (1) $\frac{1}{3} [x^3 \Psi(x^3) - \int x^2 \Psi(x^3) dx] + C$ (2) $\frac{1}{3} x^3 \Psi(x^3) - 3 \int x^3 \Psi(x^3) dx + C$
(3) $\frac{1}{3} x^3 \Psi(x^3) - \int x^2 \Psi(x^3) dx + C$ (4) $\frac{1}{3} [x^3 \Psi(x^3) - \int x^3 \Psi(x^3) dx] + C$

10. The integral $\int \left(1 + x - \frac{1}{x}\right) e^{x + \frac{1}{x}} dx$ is equal to : [JEE Main 2014]

- (1) $(x-1) e^{x + \frac{1}{x}} + c$ (2) $x e^{x + \frac{1}{x}} + c$ (3) $(x+1) e^{x + \frac{1}{x}} + c$ (4) $-x e^{x + \frac{1}{x}} + c$

11. The integral $\int \frac{dx}{x^2 (x^4 + 1)^{3/4}}$ equals : [JEE Main 2015]

- (1) $-(x^4 + 1)^{1/4} + c$ (2) $-\left(\frac{x^4 + 1}{x^4}\right)^{1/4} + c$ (3) $\left(\frac{x^4 + 1}{x^4}\right)^{1/4} + c$ (4) $(x^4 + 1)^{1/4} + c$

12. The integral $\int \frac{2x^{12} + 5x^9}{(x^5 + x^3 + 1)^3} dx$ is equal to : [JEE Main 2016]

- (1) $\frac{x^{10}}{2(x^5 + x^3 + 1)^2} + C$ (2) $\frac{x^5}{2(x^5 + x^3 + 1)^2} + C$ (3) $\frac{-x^{10}}{2(x^5 + x^3 + 1)^2} + C$ (4) $\frac{-x^5}{(x^5 + x^3 + 1)^2} + C$

Part # II

[Previous Year Questions][IIT-JEE ADVANCED]

1. Evaluate : $\int \sin^{-1} \left(\frac{2x+2}{\sqrt{4x^2+8x+13}} \right) dx$. [JEE 2001]

2. For any natural number m, evaluate $\int (x^{3m} + x^{2m} + x^m)(2x^{2m} + 3x^m + 6)^{\frac{1}{m}} dx$ where $x > 0$

[JEE 2002]

3. $\int \frac{x^2 - 1}{x^2 \sqrt{2x^4 - 2x^2 + 1}} dx$ is equal to - [JEE 2006]

- (A) $\frac{\sqrt{2x^4 - 2x^2 + 1}}{x^2} + c$ (B) $\frac{\sqrt{2x^4 - 2x^2 + 1}}{x^3} + c$
(C) $\frac{\sqrt{2x^4 - 2x^2 + 1}}{x} + c$ (D) $\frac{\sqrt{2x^4 - 2x^2 + 1}}{2x^2} + c$



4. Let $f(x) = \frac{x}{(1+x^n)^{1/n}}$ for $n \geq 2$ and $g(x) = \underset{\substack{\text{f occurs n times}}}{(f \circ f \circ \dots \circ f)}(x)$. Then $\int x^{n-2} g(x) dx$ equals. [JEE 2007]

- (A) $\frac{1}{n(n-1)}(1+nx^n)^{1-\frac{1}{n}} + K$ (B) $\frac{1}{n-1}(1+nx^n)^{1-\frac{1}{n}} + K$
 (C) $\frac{1}{n(n+1)}(1+nx^n)^{1+\frac{1}{n}} + K$ (D) $\frac{1}{n+1}(1+nx^n)^{1+\frac{1}{n}} + K$

5. Let $F(x)$ be an indefinite integral of $\sin^2 x$. [JEE 2007]

Statement-1 : The function $F(x)$ satisfies $F(x + \pi) = F(x)$ for all real x .

Statement-2 : $\sin^2(x + \pi) = \sin^2 x$ for all real x .

- (A) Statement-1 is True, Statement-2 is True ; Statement-2 is a correct explanation for Statement-1.
 (B) Statement-1 is True, Statement-2 is True ; Statement-2 is NOT a correct explanation for Statement-1.
 (C) Statement-1 is True, Statement-2 is False.
 (D) Statement-1 is False, Statement-2 is True.

6. Let $I = \int \frac{e^x}{e^{4x} + e^{2x} + 1} dx$, $J = \int \frac{e^{-x}}{e^{-4x} + e^{-2x} + 1} dx$. [JEE 2008]

Then, for an arbitrary constant c , the value of $J - I$ equals

- (A) $\frac{1}{2} \log \left(\frac{e^{4x} - e^{2x} + 1}{e^{4x} + e^{2x} + 1} \right) + c$ (B) $\frac{1}{2} \log \left(\frac{e^{4x} + e^{2x} + 1}{e^{2x} - e^{2x} + 1} \right) + c$
 (C) $\frac{1}{2} \log \left(\frac{e^{2x} - e^x + 1}{e^{2x} + e^x + 1} \right) + c$ (D) $\frac{1}{2} \log \left(\frac{e^{4x} + e^{2x} + 1}{e^{4x} - e^{2x} + 1} \right) + c$

7. The integral $\int \frac{\sec^2 x}{(\sec x + \tan x)^{9/2}} dx$ equals (for some arbitrary constant K) [JEE 2012]

- (A) $-\frac{1}{(\sec x + \tan x)^{11/2}} \left\{ \frac{1}{11} - \frac{1}{7} (\sec x + \tan x)^2 \right\} + K$
 (B) $\frac{1}{(\sec x + \tan x)^{11/2}} \left\{ \frac{1}{11} - \frac{1}{7} (\sec x + \tan x)^2 \right\} + K$
 (C) $-\frac{1}{(\sec x + \tan x)^{11/2}} \left\{ \frac{1}{11} + \frac{1}{7} (\sec x + \tan x)^2 \right\} + K$
 (D) $\frac{1}{(\sec x + \tan x)^{11/2}} \left\{ \frac{1}{11} + \frac{1}{7} (\sec x + \tan x)^2 \right\} + K$

MOCK TEST

SECTION - I : STRAIGHT OBJECTIVE TYPE

1. If $\int \frac{2 \cos x - \sin x + \lambda}{\cos x + \sin x - 2} dx = A \ln |\cos x + \sin x - 2| + Bx + C$.
Then the ordered triplet A, B, λ is
(A) $\left(\frac{1}{2}, \frac{3}{2}, -1\right)$ (B) $\left(\frac{3}{2}, \frac{1}{2}, -1\right)$ (C) $\left(\frac{1}{2}, -1, -\frac{3}{2}\right)$ (D) $\left(\frac{3}{2}, -1, \frac{1}{2}\right)$
2. The value of $\int x \cdot \frac{\ln(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx$ equals:
(A) $\sqrt{1+x^2} \ln(x + \sqrt{1+x^2}) - x + C$ (B) $\frac{x}{2} \ln(x + \sqrt{1+x^2}) - \frac{x}{\sqrt{1+x^2}} + C$
(C) $\frac{x}{2} \ln(x + \sqrt{1+x^2}) + \frac{x}{\sqrt{1+x^2}} + C$ (D) $\sqrt{1+x^2} \ln(x + \sqrt{1+x^2}) + x + C$
3. If $\int \frac{e^{x-1}}{(x^2 - 5x + 4)} 2x dx = A F(x-1) + B F(x-4) + C$ and $F(x) = \int \frac{e^x}{x} dx$, then A & B ordered set is
(A) $\left(-\frac{2}{3}, \frac{8}{3}\right)$ (B) $\left(-\frac{2}{3}, \frac{8e^3}{3}\right)$ (C) $\left(\frac{8}{3}, \frac{2}{3}\right)$ (D) $\left(-\frac{2}{3}, -\frac{8e^3}{3}\right)$
4. $I_n = \int (\log x)^n dx$, then $I_n + nI_{n-1} =$
(A) $n(\log x)^n$ (B) $(n \log x)^{n-1}$ (C) $(\log x)^{n-1}$ (D) $(n \log x)^n$
5. The value of $\int \frac{\sin^8 x - \cos^8 x}{1 - 2 \sin^2 x \cos^2 x} dx$ is :
(A) $\frac{1}{2} \sin 2x + C$ (B) $-\frac{1}{2} \sin 2x + C$ (C) $-\frac{1}{2} \sin x + C$ (D) $-\sin^2 x + C$
6. If A is square matrix and e^A is defined as $e^A = I + A + \frac{A^2}{2!} + \frac{A^3}{3!} + \dots = \frac{1}{2} \begin{bmatrix} f(x) & g(x) \\ g(x) & f(x) \end{bmatrix}$, where
 $A = \begin{bmatrix} x & x \\ x & x \end{bmatrix}$ and $0 < x < 1$, I is an identity matrix, then $\int \frac{g(x)}{f(x)} dx$ is equal to
(A) $\log(e^x + e^{-x}) + C$ (B) $\log(e^x - e^{-x}) + C$ (C) $\log(e^{2x} - 1) + C$ (D) None of these

7. The value of $\int \{1 + 2 \tan x (\tan x + \sec x)\}^{1/2} dx$ is equal to
 (A) $\bullet_n |\sec x (\sec x - \tan x)| + C$ (B) $\bullet_n |\operatorname{cosec} x (\sec x + \tan x)| + C$
 (C) $\bullet_n |\sec x (\sec x + \tan x)| + C$ (D) $\bullet_n |(\sec x + \tan x)| + C$
8. The value of $\int e^{(x \sin x + \cos x)} \left(\frac{x^4 \cos^3 x - x \sin x + \cos x}{x^2 \cos^2 x} \right) dx$, is equal to
 (A) $e^{x \sin x + \cos x} \cdot \left(x + \frac{1}{x \cos x} \right) + C$ (B) $e^{x \sin x + \cos x} \cdot \left(x \cos x + \frac{1}{x} \right) + C$
 (C) $e^{x \sin x + \cos x} \cdot \left(x - \frac{1}{x \cos x} \right) + C$ (D) none of these
9. The value of $2 \int \sin x \cdot \operatorname{cosec} 4x dx$ is equal to
 (A) $\frac{1}{2\sqrt{2}} \bullet_n \left| \frac{1 + \sqrt{2} \sin x}{1 - \sqrt{2} \sin x} \right| - \frac{1}{4} \bullet_n \left| \frac{1 + \sin x}{1 - \sin x} \right| + C$ (B) $\frac{1}{2\sqrt{2}} \bullet_n \left| \frac{1 + \sqrt{2} \sin x}{1 - \sqrt{2} \sin x} \right| + \frac{1}{4} \bullet_n \left| \frac{1 + \sin x}{1 - \sin x} \right| + C$
 (C) $\frac{1}{2\sqrt{2}} \bullet_n \left| \frac{1 - \sqrt{2} \sin x}{1 + \sqrt{2} \sin x} \right| - \frac{1}{4} \bullet_n \left| \frac{1 + \sin x}{1 - \sin x} \right| + C$ (D) none of these
10. If $\int \frac{dx}{a + \cos x} = f\left(\tan \frac{x}{2}\right) + C$, then –
 S_1 : f is a log function for $a = 0$
 S_2 : f is a inverse trigonometric function for $|a| < 1$
 S_3 : f is a polynomial function for $a = 1$
 S_4 : f is rational function but not polynomial for $a = 1$
 (A) T T T F (B) T F T F (C) F T T F (D) T F F T

SECTION - II : MULTIPLE CORRECT ANSWER TYPE

11. If $\int \frac{3 \cot 3x - \cot x}{\tan x - 3 \tan 3x} dx = p f(x) + q g(x) + C$, where 'C' is a constant of integration, then
 (A) $p = 1; q = \frac{1}{\sqrt{3}}; f(x) = x; g(x) = \bullet_n \left| \frac{\sqrt{3} - \tan x}{\sqrt{3} + \tan x} \right|$ (B) $p = 1; q = -\frac{1}{\sqrt{3}}; f(x) = x; g(x) = \bullet_n \left| \frac{\sqrt{3} - \tan x}{\sqrt{3} + \tan x} \right|$
 (C) $p = 1; q = -\frac{2}{\sqrt{3}}; f(x) = x; g(x) = \bullet_n \left| \frac{\sqrt{3} + \tan x}{\sqrt{3} - \tan x} \right|$ (D) $p = 1; q = -\frac{1}{\sqrt{3}}; f(x) = x; g(x) = \bullet_n \left| \frac{\sqrt{3} + \tan x}{\sqrt{3} - \tan x} \right|$
12. If $\int \frac{(x-1) dx}{x^2 \sqrt{2x^2 - 2x + 1}} = \frac{\sqrt{f(x)}}{g(x)} + C$, then
 (A) $f(x) = 2x^2 - 2x + 1$ (B) $g(x) = x + 1$ (C) $g(x) = x$ (D) $f(x) = \sqrt{2x^2 - 2x}$

13. If $I_n = \int \cot^n x \, dx$ and $I_0 + I_1 + 2(I_2 + \dots + I_8) + I_9 + I_{10} = A \left(u + \frac{u^2}{2} + \dots + \frac{u^9}{9} \right) + C$, where $u = \cot x$ and C is an arbitrary constant, then
 (A) A is constant (B) $A = -1$ (C) $A = 1$ (D) A is dependent on x
14. If $\int e^{3x} \cos 4x \, dx = e^{3x} (A \sin 4x + B \cos 4x) + C$ then:
 (A) $4A = 3B$ (B) $2A = 3B$ (C) $3A = 4B$ (D) $4A + 3B = 1$
15. The value of $\int 2^{mx} \cdot 3^{nx} \, dx$ (when $m, n \in \mathbb{N}$) is equal to :
 (A) $\frac{2^{mx} + 3^{nx}}{m \ln 2 + n \ln 3} + C$ (B) $\frac{e^{(m \ln 2 + n \ln 3)x}}{m \ln 2 + n \ln 3} + C$ (C) $\frac{2^{mx} \cdot 3^{nx}}{\ln(2^m \cdot 3^n)} + C$ (D) $\frac{(mn) \cdot 2^x \cdot 3^x}{m \ln 2 + n \ln 3} + C$

SECTION - III : ASSERTION AND REASON TYPE

16. **Statement-I** : If $x > 0, x \neq 1$ then $\int (\log_x e - (\log_x e)^2) dx = x \log_x e + C$
Statement-II : $\int e^x (f(x) + f'(x)) dx = e^x f(x) + C$ and $e^t = x$ iff $t = \bullet nx$
 (A) Statement-I is True, Statement-II is True; Statement-II is a correct explanation for Statement-I
 (B) Statement-I is True, Statement-II is True; Statement-II is NOT a correct explanation for Statement-I
 (C) Statement-I is True, Statement-II is False
 (D) Statement-I is False, Statement-II is True
17. **Statement-I** : If $y = \sin^{-1} x$, then $\int \sin^{-1} x \, dx = \int y \cos y \, dy + c$
Statement-II : If $y = f^{-1}(x)$, then $\int f^{-1}(x) \, dx = \int y f'(y) \, dy + c$
 (A) Statement-I is True, Statement-II is True; Statement-II is a correct explanation for Statement-I
 (B) Statement-I is True, Statement-II is True; Statement-II is NOT a correct explanation for Statement-I
 (C) Statement-I is True, Statement-II is False
 (D) Statement-I is False, Statement-II is True
18. **Statement-I** : $\int \frac{\ln(e^x + 1)}{e^x} dx = x - \left(\frac{1 + e^x}{e^x} \right) \bullet n(e^x + 1) + C$
Statement-II : $\int \frac{f'(x)}{f(x)} dx = \bullet n |f(x)| + C$
 (A) Statement-I is True, Statement-II is True; Statement-II is a correct explanation for Statement-I
 (B) Statement-I is True, Statement-II is True; Statement-II is NOT a correct explanation for Statement-I
 (C) Statement-I is True, Statement-II is False
 (D) Statement-I is False, Statement-II is True

19. **Statement-I :** $\int 2^{\tan^{-1}x} d(\cot^{-1}x) = \frac{2^{\tan^{-1}x}}{\ln 2} + c$

Statement-II : $\frac{d}{dx}(a^x + c) = a^x \bullet n a$

- (A) Statement-I is True, Statement-II is True; Statement-II is a correct explanation for Statement-I
 (B) Statement-I is True, Statement-II is True; Statement-II is NOT a correct explanation for Statement-I
 (C) Statement-I is True, Statement-II is False
 (D) Statement-I is False, Statement-II is True

20. **Statement-I :** $\int e^{\sin^{-1}x} \left(1 - \frac{x}{\sqrt{1-x^2}} \right) dx = e^{\sin^{-1}x} \cdot \sqrt{1-x^2} + c$

Statement-II : $\int e^{g(x)} (g'(x)f(x) + f'(x)) dx = e^{g(x)} f(x) + c$

- (A) Statement-I is True, Statement-II is True; Statement-II is a correct explanation for Statement-I
 (B) Statement-I is True, Statement-II is True; Statement-II is NOT a correct explanation for Statement-I
 (C) Statement-I is True, Statement-II is False
 (D) Statement-I is False, Statement-II is True

SECTION - IV : MATRIX - MATCH TYPE

21. **Column – I**

- (A) If $I = \int \frac{\sin x - \cos x}{|\sin x - \cos x|} dx$, where $\frac{\pi}{4} < x < \frac{3\pi}{8}$, then I equal to
 (B) If $\int \frac{x^2}{(x^3+1)(x^3+2)} dx = \frac{1}{3} f\left(\frac{x^3+1}{x^3+2}\right) + C$, then $f(x)$ is equal to
 (C) If $\int \sin^{-1}x \cdot \cos^{-1}x dx = f^{-1}(x) \left[\frac{\pi}{2}x - x f^{-1}(x) - 2\sqrt{1-x^2} \right] + \frac{\pi}{2}\sqrt{1-x^2} + 2x + C$, then $f(x)$ is equal to
 (D) If $\int \frac{dx}{xf(x)} = f(f(x)) + C$, then $f(x)$ is equal to

Column – II

- (p) $\sin x$
 (q) $x + c$
 (r) $\bullet n|x|$
 (s) $\sin^{-1}x$
 (t) $-x + c$

22. **Column – I**

- (A) If $F(x) = \int \frac{x + \sin x}{1 + \cos x} dx$ and $F(0) = 0$, then $F(\pi/2) =$
 (B) Let $F(x) = \int e^{\sin^{-1}x} \left(1 - \frac{x}{\sqrt{1-x^2}} \right) dx$ and $F(0) = 1$,
 If $F(1/2) = \frac{k\sqrt{3}e^{\pi/6}}{\pi}$, then $k =$

Column – II

- (p) $-\frac{\pi}{2}$
 (q) $\frac{\pi}{3}$



(C) Let $F(x) = \int \frac{dx}{(x^2+1)(x^2+9)}$ and $F(0)=0$,

(r) $\frac{\pi}{4}$

if $F(\sqrt{3}) = \frac{5}{36} k$, then $k =$

(D) Let $F(x) = \int \frac{\sqrt{\tan x}}{\sin x \cos x} dx$ and $F(0)=0$

(s) π

if $F(\pi/4) = \frac{2k}{\pi}$, then $k =$

(t) $\frac{\pi}{2}$

SECTION - V : COMPREHENSION TYPE

23. Read the following comprehension carefully and answer the questions.

Integrals of the form $\int \frac{p_m(x)}{\sqrt{ax^2+bx+c}} dx$, where $p_m(x)$ is a polynomial of degree m , are calculated by the reduction formula.

$$\int \frac{p_m(x)}{\sqrt{ax^2+bx+c}} dx = p_{m-1}(x) \sqrt{ax^2+bx+c} + \lambda \int \frac{dx}{\sqrt{ax^2+bx+c}}$$

where $p_{m-1}(x)$ is a polynomial of degree $(m-1)$ and λ is some constant number.

e.g. $I = \int \frac{x^3-x-1}{\sqrt{x^2+2x+2}} dx$ then applying the above formula, we can write

$$\int \frac{x^3-x-1}{\sqrt{x^2+2x+2}} dx = (Ax^2+Bx+C) \sqrt{x^2+2x+2} + \lambda \int \frac{dx}{\sqrt{x^2+2x+2}}$$

differentiate both sides, we get

$$\frac{x^3-x-1}{\sqrt{x^2+2x+2}} = (Ax^2+Bx+C) \cdot \frac{2(x+1)}{2\sqrt{x^2+2x+2}} + (2Ax+B) \sqrt{x^2+2x+2} + \frac{\lambda}{\sqrt{x^2+2x+2}}$$

$$x^3-x-1 = (Ax^2+Bx+C)(x+1) + (2Ax+B)(x^2+2x+2) + \lambda$$

On comparing coefficients of like powers of x we obtain the values of A , B , C and λ .

1. If $\int \frac{x^3-6x^2+11x-6}{\sqrt{x^2+4x+3}} dx = (Ax^2+Bx+C) \sqrt{x^2+4x+3} + \lambda \int \frac{dx}{\sqrt{x^2+4x+3}}$, then value of 'A' is

(A) $\frac{1}{3}$

(B) 1

(C) 3

(D) $-1/3$

2. In Q.No. 1 value of 'C' is

(A) -37

(B) $-\frac{14}{3}$

(C) $\frac{14}{3}$

(D) 37

3. In Q.No. 1 value of ' λ ', is

(A) 66

(B) -66

(C) $\frac{37}{3}$

(D) $-\frac{37}{3}$



24. Read the following comprehension carefully and answer the questions.

It is known that

$$\sqrt{\tan x} + \sqrt{\cot x} = \begin{cases} \frac{\sqrt{\sin x}}{\sqrt{\cos x}} + \frac{\sqrt{\cos x}}{\sqrt{\sin x}} & \text{if } 0 < x < \frac{\pi}{2} \\ \frac{\sqrt{-\sin x}}{\sqrt{-\cos x}} + \frac{\sqrt{-\cos x}}{\sqrt{-\sin x}} & \text{if } \pi < x < \frac{3\pi}{2} \end{cases},$$

$$\frac{d}{dx} (\sqrt{\tan x} - \sqrt{\cot x}) = \frac{1}{2} (\sqrt{\tan x} + \sqrt{\cot x}) (\tan x + \cot x), \forall x \in \left(0, \frac{\pi}{2}\right) \cup \left(\pi, \frac{3\pi}{2}\right)$$

and $\frac{d}{dx} (\sqrt{\tan x} + \sqrt{\cot x}) = \frac{1}{2} (\sqrt{\tan x} - \sqrt{\cot x}) (\tan x + \cot x), \forall x \in \left(0, \frac{\pi}{2}\right) \cup \left(\pi, \frac{3\pi}{2}\right).$

1. Value of integral $I = \int (\sqrt{\tan x} + \sqrt{\cot x}) dx$, where $x \in \left(0, \frac{\pi}{2}\right) \cup \left(\pi, \frac{3\pi}{2}\right)$ is

(A) $\sqrt{2} \tan^{-1} \left(\frac{\sqrt{\tan x} - \sqrt{\cot x}}{\sqrt{2}} \right) + C$

(B) $\sqrt{2} \tan^{-1} \left(\frac{\sqrt{\tan x} + \sqrt{\cot x}}{\sqrt{2}} \right) + C$

(C) $-\sqrt{2} \tan^{-1} \left(\frac{\sqrt{\tan x} - \sqrt{\cot x}}{\sqrt{2}} \right) + C$

(D) $-\sqrt{2} \tan^{-1} \left(\frac{\sqrt{\tan x} + \sqrt{\cot x}}{\sqrt{2}} \right) + C$

2. Value of the integral $I = \int (\sqrt{\tan x} + \sqrt{\cot x}) dx$, where $x \in \left(0, \frac{\pi}{2}\right)$, is

(A) $\sqrt{2} \sin^{-1} (\cos x - \sin x) + C$

(B) $\sqrt{2} \sin^{-1} (\sin x - \cos x) + C$

(C) $\sqrt{2} \sin^{-1} (\sin x + \cos x) + C$

(D) $-\sqrt{2} \sin^{-1} (\sin x + \cos x) + C$

3. Value of the integral $I = \int (\sqrt{\tan x} + \sqrt{\cot x}) dx$, where $x \in \left(\pi, \frac{3\pi}{2}\right)$, is

(A) $\sqrt{2} \sin^{-1} (\cos x - \sin x) + C$

(B) $\sqrt{2} \sin^{-1} (\sin x - \cos x) + C$

(C) $\sqrt{2} \sin^{-1} (\sin x + \cos x) + C$

(D) $-\sqrt{2} \sin^{-1} (\sin x + \cos x) + C$

25. Read the following comprehension carefully and answer the questions.

Let $I_{n,m} = \int \sin^n x \cos^m x dx$. Then we can relate $I_{n,m}$ with each of the following

(i) $I_{n-2,m}$

(ii) $I_{n+2,m}$

(iii) $I_{n,m-2}$

(iv) $I_{n,m+2}$

(v) $I_{n-2,m+2}$

(vi) $I_{n+2,m-2}$

Suppose we want to establish a relation between $I_{n,m}$ and $I_{n,m-2}$, then we set

$P(x) = \sin^{n+1} x \cos^{m-1} x$ (i)

In $I_{n,m}$ and $I_{n,m-2}$ the exponent of $\cos x$ is m and $m-2$ respectively, the minimum of the two is $m-2$, adding 1 to the minimum we get $m-2+1 = m-1$. Now choose the exponent $m-1$ of $\cos x$ in $P(x)$. Similarly choose the exponent of $\sin x$ for $P(x)$

Now differentiating both sides of (1), we get

$$\begin{aligned} P'(x) &= (n+1) \sin^n x \cos^m x - (m-1) \sin^{n+2} x \cos^{m-2} x \\ &= (n+1) \sin^n x \cos^m x - (m-1) \sin^n x (1 - \cos^2 x) \cos^{m-2} x \\ &= (n+1) \sin^n x \cos^m x - (m-1) \sin^n x \cos^{m-2} x + (m-1) \sin^n x \cos^m x \\ &= (n+m) \sin^n x \cos^m x - (m-1) \sin^n x \cos^{m-2} x \end{aligned}$$

Now integrating both sides, we get

$$\sin^{n+1} x \cos^{m-1} x = (n+m) I_{n,m} - (m-1) I_{n,m-2}.$$

Similarly we can establish the other relations.

1. The relation between $I_{4,2}$ and $I_{2,2}$ is

$$\begin{aligned} \text{(A)} \quad I_{4,2} &= \frac{1}{6} (-\sin^3 x \cos^3 x + 3I_{2,2}) & \text{(B)} \quad I_{4,2} &= \frac{1}{6} (\sin^3 x \cos^3 x + 3I_{2,2}) \\ \text{(C)} \quad I_{4,2} &= \frac{1}{6} (\sin^3 x \cos^3 x - 3I_{2,2}) & \text{(D)} \quad I_{4,2} &= \frac{1}{4} (-\sin^3 x \cos^3 x + 2I_{2,2}) \end{aligned}$$

2. The relation between $I_{4,2}$ and $I_{6,2}$ is

$$\begin{aligned} \text{(A)} \quad I_{4,2} &= \frac{1}{5} (\sin^5 x \cos^3 x + 8I_{6,2}) & \text{(B)} \quad I_{4,2} &= \frac{1}{5} (-\sin^5 x \cos^3 x + 8I_{6,2}) \\ \text{(C)} \quad I_{4,2} &= \frac{1}{5} (\sin^5 x \cos^3 x - 8I_{6,2}) & \text{(D)} \quad I_{4,2} &= \frac{1}{6} (\sin^5 x \cos^3 x + 8I_{6,2}) \end{aligned}$$

3. The relation between $I_{4,2}$ and $I_{4,4}$ is

$$\begin{aligned} \text{(A)} \quad I_{4,2} &= \frac{1}{3} (\sin^5 x \cos^3 x + 8I_{4,4}) & \text{(B)} \quad I_{4,2} &= \frac{1}{3} (-\sin^5 x \cos^3 x + 8I_{4,4}) \\ \text{(C)} \quad I_{4,2} &= \frac{1}{3} (\sin^5 x \cos^3 x - 8I_{4,4}) & \text{(D)} \quad I_{4,2} &= \frac{1}{3} (\sin^5 x \cos^3 x + 6I_{4,4}) \end{aligned}$$

SECTION - VI : INTEGER TYPE

26. Evaluate : $\int \cos 2x \cdot n(1 + \tan x) dx$

27. $\int \frac{x \cos \alpha + 1}{(x^2 + 2x \cos \alpha + 1)^{3/2}} dx = \frac{\lambda x}{\sqrt{x^2 + 2x \cos \alpha + 1}} + c$, then find λ

28. Evaluate : $\int \frac{\operatorname{cosec} x - \cot x}{\operatorname{cosec} x + \cot x} \cdot \frac{\sec x}{\sqrt{1 + 2 \sec x}} dx$

29. $\int \left(\frac{x-1}{x+1} \right) \frac{dx}{\sqrt{x^3 + x^2 + x}} = \lambda \tan^{-1} \sqrt{\left(x + \frac{1}{x} + 1 \right)} + c$, then find λ

30. Evaluate : $\int e^x \frac{1 + nx^{n-1} - x^{2n}}{(1 - x^n) \sqrt{1 - x^{2n}}} dx$

ANSWER KEY

EXERCISE - 1

1. B 2. A 3. B 4. A 5. B 6. D 7. C 8. B 9. A 10. B 11. D 12. B 13. C
14. B 15. B 16. C 17. B 18. C 19. B 20. A 21. B 22. A 23. C 24. A 25. B 26. A
27. B 28. C 29. C 30. B 31. B 32. A 33. A 34. B 35. C 36. B 37. D 38. A 39. A

EXERCISE - 2 : PART # I

1. ABD 2. CD 3. BCD 4. ABCD 5. ACD 6. BC 7. BCD 8. BD 9. ABCD
10. AB 11. ABC 12. BCD 13. AB 14. BC 15. ABCD

PART - II

1. A 2. A 3. A 4. C 5. A 6. C 7. C

EXERCISE - 3 : PART # I

1. $A \rightarrow p$ $B \rightarrow r$ $C \rightarrow s$ $D \rightarrow q$ 2. $A \rightarrow s$ $B \rightarrow q$ $C \rightarrow r$ 3. $A \rightarrow q$ $B \rightarrow p$ $C \rightarrow p, r$ $D \rightarrow p, r, s$
4. $A \rightarrow p$ $B \rightarrow p$ $C \rightarrow r$ $D \rightarrow s$ 5. $A \rightarrow s$ $B \rightarrow q$ $C \rightarrow r$ $D \rightarrow p$

PART - II

- Comprehension #1: 1. B 2. C Comprehension #2: 1. A 2. B 3. A
Comprehension #3: 1. A 2. A 3. B Comprehension #4: 1. C 2. B 3. D

EXERCISE - 5 : PART # I

1. 3 2. 2 3. 2 4. 4 5. 4 6. 1 7. 3 8. 1 9. 3 10. 2 11. 2 12. 1

PART - II

1. $(x+1)\tan^{-1}\frac{2(x+1)}{3} - \frac{3}{4}\ln(4x^2+8x+13)+c$ 2. $\frac{(2x^{3m}+3x^{2m}+6x^m)^{\frac{m+1}{m}}}{6(m+1)} + C$
3. D 4. A 5. D 6. C 7. C

MOCK TEST

1. B 2. A 3. B 4. A 5. B 6. A 7. C 8. C 9. A
 10. B 11. AD 12. AC 13. AB 14. CD 15. BC 16. A 17. A 18. A
 19. D 20. A
 21. $A \rightarrow q \quad B \rightarrow r \quad C \rightarrow p \quad D \rightarrow r$ 22. $A \rightarrow t \quad B \rightarrow t \quad C \rightarrow r \quad D \rightarrow s$
 23. 1. A 2. D 3. B 24. 1. A 2. B 3. A 25. 1. A 2. A 3. B
 26. $\frac{1}{2} [\sin 2x \cdot n(1 + \tan x) - x + n |\sin x + \cos x|] + C$ 27. 1 28. $\sin^{-1} \left(\frac{1}{2} \sec^2 \frac{x}{2} \right) + C$ 29. 2
 30. $e^x \sqrt{\frac{1+x^n}{1-x^n}} + C$