

## HINTS & SOLUTIONS

### EXERCISE - 1 Single Choice

2.  $\beta = \cos^{-1} \left( \cos \left( \frac{\pi}{2} - \cos^{-1} x \right) \right); \beta = \cos^{-1} \{ \cos(\sin^{-1} x) \}$

also  $\alpha = \sin^{-1} [\cos(\sin^{-1} x)]$

$\alpha + \beta = \pi/2 \Rightarrow \tan \alpha = \cot \beta$

3.  $(\sin^{-1} x + \sin^{-1} y)^2 = \pi^2$

$\Rightarrow \sin^{-1} x + \sin^{-1} y = \pm \pi$

$\Rightarrow \sin^{-1} x = \sin^{-1} y = \frac{\pi}{2}$

or  $\sin^{-1} x = \sin^{-1} y = -\frac{\pi}{2}$

$\Rightarrow x^2 + y^2 = 2.$

4.  $x = \sin 2\theta = 2 \sin \theta \cos \theta = \frac{4}{5}; y = \sin \frac{\phi}{2};$

$y > 0, \tan \phi = \frac{4}{3}$

$y^2 = \sin^2 \frac{\phi}{2} = \frac{1 - \cos \phi}{2} = \frac{1}{5} = 1 - \frac{4}{5} = 1 - x$

$\Rightarrow y^2 = 1 - x$

7.  $\tan \left[ \sin^{-1} \frac{3}{5} + \tan^{-1} \frac{2}{3} \right]$

$\Rightarrow \tan \left[ \tan^{-1} \frac{\frac{3}{5} + \frac{2}{3}}{1 - \frac{6}{15}} \right] = \tan \tan^{-1} \frac{17}{6} = \frac{17}{6}$

8.  $\sin^{-1} x - \cos^{-1} x = \cos^{-1} \frac{\sqrt{3}}{2}$

$\frac{\pi}{2} - 2\cos^{-1} x = \cos^{-1} \frac{\sqrt{3}}{2} = \frac{\pi}{6}$

$\Rightarrow \cos^{-1} x = \frac{\pi}{6}$

$x = \frac{\sqrt{3}}{2}$

9.  $\frac{1}{\sqrt{2}} \left( \cos \frac{7\pi}{5} - \sin \frac{2\pi}{5} \right)$

$= \cos \frac{\pi}{4} \cos \frac{7\pi}{5} - \sin \frac{\pi}{4} \sin \frac{2\pi}{5}$

$= \cos \frac{\pi}{4} \cos \frac{7\pi}{5} + \sin \frac{\pi}{4} \sin \frac{7\pi}{5}$

11.  $\sin^{-1} \left( -\sin \frac{50\pi}{9} \right) = -\sin^{-1} \sin \left( \frac{50\pi}{9} \right)$

$= -\sin^{-1} \sin \left( \frac{14\pi}{9} \right) = -\sin^{-1} \sin \left( 2\pi - \frac{4\pi}{9} \right)$

$= -\sin^{-1} \sin \left( -\frac{4\pi}{9} \right) = \frac{4\pi}{9}$

$\cos^{-1} \cos \left( -\frac{31\pi}{9} \right) = \cos^{-1} \cos \left( \frac{31\pi}{9} \right)$

$= \cos^{-1} \cos \left( 4\pi - \frac{5\pi}{9} \right) = \cos^{-1} \cos \frac{5\pi}{9} = \frac{5\pi}{9}$

Hence  $\sec \left( \frac{4\pi}{9} + \frac{5\pi}{9} \right) = \sec \pi = -1$

12.  $x = \frac{2\pi}{3} + \frac{\pi}{4} + \frac{\pi}{3} = \frac{5\pi}{4}$

$y = \cos \left( \frac{1}{2} \sin^{-1} \left( \sin \frac{5\pi}{8} \right) \right)$

$= \cos \left( \frac{1}{2} \left( \pi - \frac{5\pi}{8} \right) \right) = \cos \frac{3\pi}{16}.$

14.  $\tan^{-1} 2 + \tan^{-1} 3 = \operatorname{cosec}^{-1} x$

$\Rightarrow \pi + \tan^{-1}(-1) = \operatorname{cosec}^{-1} x$

$\Rightarrow \pi - \frac{\pi}{4} = \operatorname{cosec}^{-1} x \Rightarrow \frac{3\pi}{4} = \operatorname{cosec}^{-1} x$

$\Rightarrow \text{no solution. } \left\{ -\frac{\pi}{2} \leq \operatorname{cosec}^{-1} x \leq \frac{\pi}{2} \right\}$



15.  $\sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}$  and  
 $\sin^{-1}x - \cos^{-1}x = \sin^{-1}(3x - 2)$   
 $- \quad + \quad -$

$$2\cos^{-1}x = \cos^{-1}(3x - 2)$$

Also  $x \in [-1, 1]$

$$\begin{aligned} \cos^{-1}(2x^2 - 1) &= \cos^{-1}(3x - 2) \\ \text{and } (3x - 2) &\in [-1, 1] \text{ i.e. } -1 \leq 3x - 2 \leq 1 \\ 2x^2 - 1 &= 3x - 2 \end{aligned}$$

hence  $x \in \left[\frac{1}{3}, 1\right]$

$$2x^2 - 3x + 1 = 0 \Rightarrow x = 1 \text{ or } 1/2 \Rightarrow A$$

$$\begin{aligned} &= \cos\left(\frac{7\pi}{5} - \frac{\pi}{4}\right) = \cos\left(\frac{23\pi}{20}\right) = \cos\left(\pi + \frac{3\pi}{20}\right) \\ &= \cos\left(\pi - \frac{3\pi}{20}\right) = \cos\frac{17\pi}{20} \end{aligned}$$

$$\therefore \cos^{-1}\left\{\frac{1}{\sqrt{2}}\left(\cos\frac{7\pi}{5} - \sin\frac{2\pi}{5}\right)\right\} = \cos^{-1}\left(\cos\left(\frac{17\pi}{20}\right)\right)$$

$$= \frac{17\pi}{20} \text{ (since } \frac{17\pi}{20} \text{ lies between } 0 \text{ and } \pi)$$

16.  $\sin^{-1}\sqrt{1-x^2} + \cos^{-1}x = \cot^{-1}\frac{\sqrt{1-x^2}}{x} - \sin^{-1}x$

or  $\frac{\pi}{2} + \sin^{-1}\sqrt{1-x^2} = \cot^{-1}\frac{\sqrt{1-x^2}}{x}$

$$\tan^{-1}\frac{\sqrt{1-x^2}}{x} + \sin^{-1}\sqrt{1-x^2} = 0$$

$$\Rightarrow -1 \leq x < 0 \cup \{1\} \Rightarrow C$$

18.  $(\tan^{-1}x)^2 - 3\tan^{-1}x + 2 \geq 0$

$$(\tan^{-1}x - 1)(\tan^{-1}x - 2) \geq 0$$

we know that  $\tan^{-1}x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

so  $\tan^{-1}x \geq 2$  (not possible) or  $\tan^{-1}x \leq 1$

$$\Rightarrow x \in (-\infty, \tan 1]$$

19.  $\sin^{-1}x + \sin^{-1}y + \sin^{-1}z = \frac{3\pi}{2}$

$$\Rightarrow x = y = z = 1$$

$$\Rightarrow x^{100} + y^{100} + z^{100} - \frac{9}{x^{101} + y^{101} + z^{101}} = 0$$

20.  $\cot^{-1}\left\{\frac{\sqrt{1-\sin x} + \sqrt{1+\sin x}}{\sqrt{1-\sin x} - \sqrt{1+\sin x}}\right\} \quad \frac{\pi}{2} < x < \pi$

Rationalize the term in the bracket

$$= \cot^{-1}\left(\frac{2 + 2\sqrt{1-\sin^2 x}}{-2\sin x}\right) = \cot^{-1}\left(\frac{1-\cos x}{-\sin x}\right)$$

$$= \cot^{-1}\left(-\tan\frac{x}{2}\right) = \frac{\pi}{2} - \tan^{-1}\left(-\tan\frac{x}{2}\right)$$

$$= \frac{\pi}{2} + \tan^{-1}\tan\frac{x}{2} \quad \text{since } \frac{x}{2} \in \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$$

$$= \frac{\pi}{2} + \frac{x}{2}$$

21.  $\sin^{-1}\left(\frac{3\sin 2\theta}{5+4\cos 2\theta}\right) = \frac{\pi}{2}$

Taking sin on both side  $\frac{3\sin 2\theta}{5+4\cos 2\theta} = 1$

$$3\sin 2\theta = 5 + 4 \cos 2\theta$$

$$\frac{6\tan\theta}{1+\tan^2\theta} = 5 + 4\left(\frac{1-\tan^2\theta}{1+\tan^2\theta}\right)$$

$$\tan^2\theta - 6\tan\theta + 9 = 0$$

$$\tan\theta = 3$$

22. Consider  $\tan 65^\circ - 2 \tan 40^\circ$

$$\tan(45^\circ + 20^\circ) - 2 \tan 40^\circ$$

$$\frac{1 + \tan 20^\circ}{1 - \tan 20^\circ} - \frac{4 \tan 20^\circ}{1 - \tan^2 20^\circ}$$

$$\frac{(1 + \tan 20^\circ)^2 - 4 \tan 20^\circ}{(1 - \tan 20^\circ)(1 + \tan 20^\circ)}$$



$$= \frac{(1-\tan 20^\circ)(1-\tan 20^\circ)}{(1-\tan 20^\circ)(1+\tan 20^\circ)} = \frac{(1-\tan 20^\circ)}{(1+\tan 20^\circ)}$$

$$= \tan(45^\circ - 20^\circ) = \tan 25^\circ$$

$$\therefore \tan^{-1}(\tan 25^\circ) = 25^\circ$$

$$23. \cos^{-1}(2x) + \cos^{-1}(3x) = \pi - \cos^{-1}(x) = \cos^{-1}(-x)$$

$$\cos^{-1}[(2x)(3x) - \sqrt{1-4x^2}\sqrt{1-9x^2}] = \cos^{-1}(-x)$$

$$6x^2 - \sqrt{1-4x^2} \cdot \sqrt{1-9x^2} = -x$$

$$(6x^2 + x)^2 = (1-4x^2)(1-9x^2)$$

$$\Rightarrow x^2 + 12x^3 = 1 - 13x^2$$

$$\Rightarrow 12x^3 + 14x^2 - 1 = 0$$

$$\therefore a = 12; b = 14; c = 0$$

$$\Rightarrow a + b + c = 26$$

$$24. x = \frac{1}{2}; \quad y = \frac{1}{3} \quad (\text{from 2nd relation})$$


$$\cot^{-1} \frac{1}{2} + \cot^{-1} \frac{1}{3} = \tan^{-1} 2 + \tan^{-1} 3 = \frac{3\pi}{4} \text{ Ans. }]$$

$$25. \tan\left(\frac{\pi}{4} + \alpha\right) \quad \text{when } \alpha = \tan^{-1}\left(\frac{\frac{1}{4} + \frac{1}{5}}{1 - \frac{1}{20}}\right);$$

$$\alpha = \tan^{-1}\left(\frac{9}{19}\right) = \frac{1 + \frac{9}{19}}{1 - \frac{9}{19}} = \frac{28}{10} = \frac{14}{5} = \frac{a}{b}$$

$$\Rightarrow 14 + 5 = 19$$


26. Given equation is  $|\cos x| = \sin^{-1}(\sin x)$   $-\pi \leq x \leq \pi$



27. Let  $\tan^{-1}(x) = \theta$

$$\Rightarrow x = \tan \theta \\ \cos \theta = x \quad (\text{given})$$

$$\frac{1}{\sqrt{1+x^2}} = x$$



$$x^2(1+x^2) = 1$$

$$\Rightarrow x^2 = \frac{-1 \pm \sqrt{5}}{2} \quad (x^2 \text{ can not be -ve})$$

$$\Rightarrow x^2 = \frac{\sqrt{5}-1}{2} \Rightarrow \frac{x^2}{2} = \frac{\sqrt{5}-1}{4}$$

$$\cos^{-1}\left(\frac{\sqrt{5}-1}{4}\right) = \cos^{-1}\left(\sin\frac{\pi}{10}\right) = \cos^{-1}\left(\cos\frac{2\pi}{5}\right) = \frac{2\pi}{5}$$

28. Let  $x = \cos \theta, x \geq 0$

$$\text{LHS} = 20$$

$$\text{RHS} = \sin^{-1}(|2 \cos \theta| \sin \theta)$$

$$\Rightarrow \sin^2(\sin 2\theta)$$

$$\sin^{-1}(\sin 2\theta) \quad 0 \leq 2\theta \leq \pi/2$$

$$0 \leq \theta \leq \frac{\pi}{4} \Rightarrow \frac{1}{\sqrt{2}} \leq x \leq 1$$

29.  $x(x+1) \geq 0$  and  $0 \leq x^2 + x + 1 \leq 1$

$$\Rightarrow x \geq 0 \text{ or } x \leq -1 \quad \text{and} \quad x(x+1) \leq 0 \\ x \leq 0 \quad \text{or} \quad x \geq -1$$

Hence  $x = 0$  or  $x = -1$

## EXERCISE - 2

## Part # I : Multiple Choice

1. Let  $\tan^{-1}x = \alpha$  and  $\tan^{-1}x^3 = \beta$   
 $\tan \alpha = x$  and  $\tan \beta = x^3$

$$\therefore 2 \tan(\alpha + \beta) = \frac{2(\tan \alpha + \tan \beta)}{1 - \tan \alpha \tan \beta}$$

$$= 2 \left[ \frac{x + x^3}{1 - x^4} \right] = \frac{2x}{1 - x^2} \Rightarrow (\text{A})$$

$$\text{Also } \frac{2x}{1 - x^2} = \frac{2 \tan \alpha}{1 - \tan^2 \alpha} = \tan 2\alpha = \tan(2 \tan^{-1} x)$$

$\Rightarrow (\text{B})$

$$= \tan \left( 2 \left( \frac{\pi}{2} - \cot^{-1} x \right) \right) = \tan(\pi - \cot^{-1} x - \cot^{-1} x)$$

$$= \tan \left( \cot^{-1}(-x) - \cot^{-1}(x) \right)$$

$\Rightarrow (\text{C})$

2. Let  $\tan^{-1} \frac{a}{x} = \alpha \Rightarrow \tan \alpha = \frac{a}{x}$  etc.

$$\alpha + \beta + \gamma + \delta = \frac{\pi}{2}$$

$$\tan(\alpha + \beta + \gamma + \delta) = \tan \frac{\pi}{2}$$

$$\frac{S_2 - S_3}{1 - S_2 + S_4} = \infty$$

$$\Rightarrow 1 - S_2 + S_4 = 0$$

$$\Rightarrow S_4 - S_2 + 1 = 0$$

$$\text{How, } S_4 = \tan \alpha \cdot \tan \beta \cdot \tan \gamma \cdot \tan \delta = \frac{abcd}{x^4}$$

$$S_2 = \sum \tan \alpha \cdot \tan \beta = \frac{ab}{x^2}$$

$$\therefore \frac{abcd}{x^4} - \frac{ab}{x^2} + 1 = 0$$

$$x^4 - \sum ab x^2 + abcd = 0$$

$$\therefore x_1 + x_2 + x_3 + x_4 = 0 \quad \dots \text{(i)}$$

$$\sum x_1 x_2 x_3 = 0$$

..... (ii)

$$\sum x_1 x_2 x_3 = \underset{\text{non zero}}{1 \ 4 \ 2 \ 3 \ 4} \left[ \frac{1}{x_1} + \frac{1}{x_2} + \frac{1}{x_3} + \frac{1}{x_4} \right] = 0$$

$\Rightarrow (\text{B})$

$$x_1 x_2 x_3 x_4 = abcd$$

$\Rightarrow (\text{C})$

$$3. \cos \left( -\frac{14\pi}{5} \right) = \cos \frac{14\pi}{5} = \cos \frac{4\pi}{5}$$

$$\text{Hence } \cos \left( \frac{1}{2} \cos^{-1} \left( \cos \frac{4\pi}{5} \right) \right)$$

$$= \cos \frac{4\pi}{10} = \cos \frac{2\pi}{5}$$

$\Rightarrow \text{BCD}$

6. for (A) and (B)

$$\cos(\cos^{-1} 1) = 1$$

$$\Rightarrow \cos(\cos(\cos^{-1} 1)) = \cos 1$$

$$\sin^{-1}(\sin(\pi - 1)) = \pi - (\pi - 1) = 1$$

$$\Rightarrow \sin(\sin^{-1}(\sin(\pi - 1))) = \sin 1$$

$$\cos^{-1}(\cos(2\pi - 2)) = \cos^{-1}(\cos 2) = 2$$

$$\Rightarrow \sin(\cos^{-1}(\cos(2\pi - 2))) = \sin 2$$

$$(\tan(\cot^{-1}(\cot 1)) = \tan 1$$

It is easy to compare

$$\cos 1, \sin 1, \sin 2, \tan 1 \cos 1 < \sin 1 < \sin 2 < \tan 1$$

$\Rightarrow (\text{A})$  is correct

for (C)

$\cos^{-1} \cos x$  is periodic and even

$$\cos^{-1} \cos(2t\pi - 1) = \cos^{-1}(\cos 1) = 1 (t \in \mathbb{I})$$

$$\sum_{t=1}^{5000} \cos^{-1} \cos(2t\pi - 1) = 5000$$

$$\text{now } \cot^{-1} \cot(t\pi + 2) = 2$$

$[\cot^{-1} \cot x$  is periodic with period  $\pi$ ]



## MATHS FOR JEE MAIN & ADVANCED

7.  $2x = \tan(2\tan^{-1}a) + 2\tan(\tan^{-1}a + \tan^{-1}a^3)$

$$2x = \frac{2a}{1-a^2} + \frac{2(a+a^3)}{1-a^4}$$

(Using  $\tan 2\theta = \frac{2\tan\theta}{1-\tan^2\theta}$ )

$\therefore a \neq \pm 1$

$\Rightarrow$  (D)

$$x = \frac{a}{1-a^2} + \frac{a}{1-a^2} = \frac{2a}{1-a^2}$$

$$\Rightarrow x(1-a^2) = 2a$$

$$\Rightarrow a^2x + 2a = x$$

$\Rightarrow$  (A)

Hence B & C are invalid

10.  $\sum_{n=1}^{\infty} \tan^{-1} \frac{4n}{n^4 - 2n^2 + 2}$

$$= \lim_{k \rightarrow \infty} \sum_{n=1}^k \left\{ \tan^{-1}(n+1)^2 - \tan^{-1}(n-1)^2 \right\}$$

$$= \lim_{k \rightarrow \infty} \left\{ \tan^{-1}(k+1)^2 + \tan^{-1}k^2 - \tan^{-1}1 - \tan^{-1}0 \right\}$$

$$= \frac{\pi}{2} + \frac{\pi}{2} - \frac{\pi}{4} - 0 = \frac{3\pi}{4}$$

Also  $\tan^{-1}2 + \tan^{-1}3 = \pi + \tan^{-1}\left(\frac{3+2}{1-3 \cdot 2}\right)$

Since  $xy = 6 > 1$

$$= \frac{3\pi}{4} \text{ and } \sec^{-1}(-\sqrt{2}) = \frac{3\pi}{4}$$

12.  $\sum_{r=1}^{\infty} T_r = \cot^{-1}\left(r^2 + \frac{3}{4}\right) = \sum_{r=1}^{\infty} \tan^{-1}\left(\frac{1}{1+r^2 - \frac{1}{4}}\right)$

$$= \sum_{r=1}^{\infty} \tan^{-1}\left(\frac{\left(r+\frac{1}{2}\right) - \left(r-\frac{1}{2}\right)}{1 + \left(r-\frac{1}{2}\right)\left(r+\frac{1}{2}\right)}\right)$$

$$= \sum_{r=1}^{\infty} \tan^{-1}\left(r + \frac{1}{2}\right) - \sum_{r=1}^{\infty} \tan^{-1}\left(r - \frac{1}{2}\right)$$

Now it can be solved.

13. Let  $y = \frac{2(x^2 + 1) + 1}{(x^2 + 1)} = 2 + \frac{1}{(x^2 + 1)}$

$$\Rightarrow 2 < y \leq 3$$

Now  $\sin^{-1}\sin y \leq \pi - \frac{5}{2}$

$$\Rightarrow \pi - y \leq \pi - \frac{5}{2} \Rightarrow y \geq \frac{5}{2}$$

$$\Rightarrow \frac{2x^2 + 3}{x^2 + 1} \geq \frac{5}{2}$$

Now it can be solved

### Part # II : Assertion & Reason

2.  $x(x-2)(3x-7)=2$   
 $\Rightarrow 3x^3 - 13x^2 + 14x - 2 = 0$

$$s_1 = r + s + t = \frac{13}{3};$$

$$s_2 = \frac{14}{3}, s_3 = \frac{2}{3}$$

$$\tan^{-1}r + \tan^{-1}s + \tan^{-1}t = \pi + \tan^{-1}\left[\frac{s_1 - s_3}{1 - s_2}\right]$$

$$= \pi + \tan^{-1}[-1] = \frac{3\pi}{4}$$

Hence statement-I and statement-II both are true.

8. Using properties

$$\therefore \tan^{-1}x + \cot^{-1}x = \frac{\pi}{2}$$

$$\Rightarrow \frac{a}{x} = \frac{x}{b} \Rightarrow x = \sqrt{ab}$$

Statement-1 is true

$$\tan^{-1}\left(\frac{m}{n}\right) + \tan^{-1}\left(\frac{1 - \frac{m}{n}}{1 + \frac{m}{n}}\right)$$

$$= \tan^{-1}\frac{m}{n} + \tan^{-1}1 - \tan^{-1}\frac{m}{n} = \frac{\pi}{4}$$



## EXERCISE - 3

## Part # I : Matrix Match Type

4. (A)

$$\text{Let } x = \sqrt{\frac{a(a+b+c)}{bc}}, y = \sqrt{\frac{b(a+b+c)}{ac}},$$

$$z = , x, y, z > 0$$

$$\Rightarrow \theta = \tan^{-1}x + \tan^{-1}y + \tan^{-1}z$$

$$\text{Now } x + y + z =$$

$$\sqrt{\frac{a(a+b+c)}{bc}} + \sqrt{\frac{b(a+b+c)}{ac}} + \sqrt{\frac{c(a+b+c)}{ab}}$$

$$= \frac{(a+b+c)^{3/2}}{\sqrt{abc}}$$

$$\text{and } xyz = \frac{(a+b+c)^{3/2}}{\sqrt{abc}}$$

$$\Rightarrow x + y + z = xyz \Rightarrow \tan^{-1}x + \tan^{-1}y + \tan^{-1}z = \pi$$

$$\text{Hence } \theta = \pi$$

 (B) Let  $\alpha = \tan^{-1}(\cot A)$ 

$$\beta = \tan^{-1}(\cot^3 A)$$

$$\tan(\alpha + \beta) = \frac{\cot A + \cot^3 A}{1 - \cot^4 A}$$

$$\text{R.H.S. is negative} \Rightarrow \pi < \alpha + \beta < \frac{\pi}{2}$$

$$\tan(\alpha + \beta - \pi) = \frac{\cot A}{1 - \cot^2 A} = -\frac{\tan 2A}{2}$$

$$\Rightarrow \alpha + \beta = \pi - \tan^{-1}\left(\frac{\tan 2A}{2}\right)$$

$$\text{G.E.} = \pi \text{ independent of A.}$$

$$(C) x = \tan \theta \quad \theta < -\frac{\pi}{4} \quad \text{or} \quad \theta > \frac{\pi}{4}$$

$$\sin^{-1}\left(\frac{2x}{1+x^2}\right) = \sin^{-1}(\sin 2\theta) - \pi < 2\theta < -\frac{\pi}{2}$$

$$\text{or} \quad \frac{\pi}{2} < 2\theta$$

$$= \begin{cases} -\pi - 2\theta & ; \theta < -\frac{\pi}{4} \\ \pi - 2\theta & ; \theta > \frac{\pi}{4} \end{cases}$$

$$= \begin{cases} -\pi - 2\tan^{-1}x & ; x < -1 \\ \pi - 2\tan^{-1}x & ; x > 1 \end{cases}$$

$$\sin^{-1}\left(\frac{2x}{1+x^2}\right) + 2\tan^{-1}x = -\pi$$

$$(D) \sin^{-1}\left(\frac{3}{5}\right) - \cos^{-1}\left(\frac{12}{13}\right) + \cos^{-1}\left(\frac{16}{65}\right)$$

$$= \sin^{-1}\left(\frac{3}{5}\right) - \sin^{-1}\left(\frac{5}{13}\right) + \cos^{-1}\left(\frac{16}{65}\right)$$

$$= \sin^{-1}\left\{\frac{3}{5}\sqrt{1-\left(\frac{5}{13}\right)^2} - \frac{5}{13}\sqrt{1-\left(\frac{3}{5}\right)^2}\right\} + \cos^{-1}\left(\frac{16}{65}\right)$$

$$= \sin^{-1}\left\{\frac{3}{5}\cdot\frac{12}{13} - \frac{5}{13}\cdot\frac{4}{5}\right\} + \cos^{-1}\left(\frac{16}{65}\right)$$

$$= \sin^{-1}\left(\frac{16}{65}\right) + \cos^{-1}\left(\frac{16}{65}\right) = \frac{\pi}{2}$$

## Part # II : Comprehension

## Comprehension-1

$$1. (i) \sin\left(\frac{\cos^{-1} x}{y}\right) = 1$$

$$\Rightarrow \frac{\cos^{-1} x}{y} = 2n\pi + \frac{\pi}{2} \quad \& \quad y \neq 0$$

$$\Rightarrow \cos^{-1} x = (4n+1) \frac{\pi}{2} y$$



EXERCISE - 4

Subjective Type

when  $n = 0 \Rightarrow \cos^{-1} x = \frac{\pi}{2}$

when  $y = 1, x = 0 \quad \{0 < \frac{\pi}{2} y \leq \pi \text{ } y = 2, x = -1 \Rightarrow 0 < y \leq 2\}$

when

$n = 1 \text{ or } > 1 \quad \cos^{-1} x = \frac{5\pi}{2} \text{ } y \text{ or more (reject)}$

$n = -1 \text{ or } < -1 \quad \cos^{-1} x = \frac{-3\pi}{2} \text{ } y \text{ or more (reject)}$

(ii)  $\cos \left( \frac{\sin^{-1} x}{y} \right) = 0$

$\Rightarrow \frac{\sin^{-1} x}{y} = (2n + 1) \frac{\pi}{2} \quad \& \quad y \neq 0$

$n = 0 \quad \sin^{-1} x = \frac{\pi}{2} y$

$\{ \frac{-\pi}{2} \leq \frac{\pi}{2} y \leq \frac{\pi}{2} \Rightarrow -1 \leq y \leq 1 \}$

When  $y = 1, x = 1 \Rightarrow y = -1, x = -1$

$n = -1 \quad \sin^{-1} x = -\frac{\pi}{2} y$

When  $y = 1, x = -1 \Rightarrow y = -1, x = 1$

Other values of  $n$  &  $y$  are out of range.

1.  $(0, 1) \text{ & } (-1, 2)$
2.  $(1, 1), (1, -1), (-1, 1), (-1, -1)$
3. one one onto

1. (i) Let  $\tan^{-1} x = \theta \Rightarrow \tan \theta = x \cot \theta = \frac{1}{x} \forall x > 0$

$\theta = -\pi + \cot^{-1} \frac{1}{x} \forall x < 0$

$\sin \theta = \frac{x}{\sqrt{1+x^2}} \Rightarrow \theta = \sin^{-1} \frac{x}{\sqrt{1+x^2}}$

$\cos \theta = \frac{1}{\sqrt{1+x^2}} \quad x > 0$

$\theta = \cos^{-1} \frac{1}{\sqrt{1+x^2}} = \tan^{-1} x \quad x > 0$

$\Rightarrow \tan^{-1} x = -\pi + \cot^{-1} \frac{1}{x}$

$= \sin^{-1} \frac{x}{\sqrt{1+x^2}} = -\cos^{-1} \frac{x}{\sqrt{1+x^2}}$

where  $x < 0$

(ii) Let

$\theta = \cos^{-1} x \quad \text{given} \quad -1 < x < 0$

$\Rightarrow \cos \theta = x \quad \theta \in (\frac{\pi}{2}, \pi)$

$\sec \theta = \frac{1}{x} \quad \theta = \sec^{-1} \frac{1}{x}$

$\sin \theta = \sqrt{1-x^2} \Rightarrow \theta = \pi - \sin^{-1} \sqrt{1-x^2}$

$\tan \theta = \frac{\sqrt{1-x^2}}{x} \Rightarrow \theta = \pi + \tan^{-1} \frac{\sqrt{1-x^2}}{x}$

$\cot \theta = \frac{x}{\sqrt{1-x^2}} \Rightarrow \theta = \cot^{-1} \frac{x}{\sqrt{1-x^2}}$



5. (B)  $\sin(\sin^{-1}(\log_{1/2}x)) + 2|\cos(\sin^{-1}(x/2-1))| = 0$

$$-1 \leq \log_{1/2} x \leq 1 \Rightarrow \frac{1}{2} \leq x \leq 2 \quad \dots\dots \text{(i)}$$


and  $-1 \leq \frac{x}{2} - 1 \leq 1$

$$\Rightarrow 0 \leq \frac{x}{2} \leq 2 \Rightarrow 0 \leq x \leq 4 \quad \dots\dots \text{(ii)}$$

From (i) & (ii),  $\frac{1}{2} \leq x \leq 2$

Also  $\log_{1/2} x + 2\sqrt{x - \frac{x^2}{4}} = 0$

$$2\sqrt{x - \frac{x^2}{4}} = \log_2 x \quad \dots(1)$$



From graph it is clear that equation (1) does not have

any solution in  $\left[\frac{1}{2}, 2\right]$

8.  $0 \leq (\tan^{-1} x)^2 \leq \frac{\pi^2}{4}$   $\Rightarrow (\tan^{-1} x)^2 + (\cos^{-1} x)^2 \leq \frac{5\pi^2}{4}$   
 $0 \leq (\cos^{-1} y)^2 \leq \pi^2$

But  $(\tan^{-1} x)^2 + (\cos^{-1} x)^2 = \pi^2 k$

Hence  $k\pi^2 \leq \frac{5\pi^2}{4}$ ,  $k \leq \frac{5}{4}$   $\dots\dots \text{(i)}$

Now put  $\tan^{-1} x = \frac{\pi}{2} - \cos^{-1} y$

$$\left(\frac{\pi}{2} - \cos^{-1} y\right)^2 + (\cos^{-1} y)^2 = \pi^2 k$$

(where  $\cos^{-1} y = t$ )

$$2t^2 - \pi t + \left(\frac{\pi^2}{4} - k\pi^2\right) = 0$$

For real roots,  $D \geq 0$

$$\pi^2 - 8\left(\frac{\pi^2}{4} - k\pi^2\right) \geq 0$$

$$\Rightarrow 1 - 2 + 8k \geq 0, k \geq \frac{1}{8} \quad \dots\dots \text{(ii)}$$

From (i) and (ii),  $k = 1$

With

$$k = 1, t = \frac{\pi \pm \sqrt{8\pi^2 - \pi^2}}{4} = \frac{\pi + \sqrt{7}\pi}{4} = (1 \pm \sqrt{7}) \frac{\pi}{4}$$

or  $\cos^{-1} y = (\sqrt{7} + 1) \frac{\pi}{4}$  (as  $0 \leq \cos^{-1} y \leq \pi$ )

$$\therefore y = \cos((\sqrt{7} + 1) \frac{\pi}{4})$$

$$\therefore \tan^{-1} x = \frac{\pi}{2} - (\sqrt{7} + 1) \frac{\pi}{4} = \frac{\pi}{4} [(1 - \sqrt{7})]$$

$$\Rightarrow x = \tan((1 - \sqrt{7}) \frac{\pi}{4})$$



EXERCISE - 5

Part # I : AIEEE/JEE-MAIN

$$1. \text{ Now, } \tan^{-1} \frac{1}{4} + \tan^{-1} \frac{2}{9} = \tan^{-1} \left( \frac{\frac{1}{4} + \frac{2}{9}}{1 - \frac{1}{4} \times \frac{2}{9}} \right) \\ = \tan^{-1} \left( \frac{17}{34} \right) = \tan^{-1} \left( \frac{1}{2} \right)$$

$$2. \text{ Given that } \cot^{-1} (\sqrt{\cos \alpha}) - \tan^{-1} (\sqrt{\cos \alpha}) = x$$

$$\text{We know that, } \cot^{-1} (\sqrt{\cos \alpha}) + \tan^{-1} (\sqrt{\cos \alpha}) = \frac{\pi}{2}$$

.....(ii)

On adding equations (i) and (ii),

$$\text{We get } 2 \cot^{-1} (\sqrt{\cos \alpha}) = \frac{\pi}{2} + x$$

$$\Rightarrow \sqrt{\cos \alpha} = \cot \left( \frac{\pi}{4} + \frac{x}{2} \right) \Rightarrow \sqrt{\cos \alpha}$$

$$= \frac{\cot \frac{x}{2} - 1}{1 + \cot \frac{x}{2}} \Rightarrow \sqrt{\cos \alpha} = \frac{\cos \frac{x}{2} - \sin \frac{x}{2}}{\cos \frac{x}{2} + \sin \frac{x}{2}}$$

$$\Rightarrow \cos \alpha = \frac{1 - \sin x}{1 + \sin x}$$

$$\Rightarrow \frac{1 - \tan^2 \frac{\alpha}{2}}{1 + \tan^2 \frac{\alpha}{2}} = \frac{1 - \sin x}{1 + \sin x}$$

Applying componendo and dividendo rule,

$$\text{We get } \sin x = \tan^2 \left( \frac{\alpha}{2} \right)$$

$$3. \text{ Given that, } \sin^{-1} x = 2 \sin^{-1} \alpha$$

$$\text{Since, } -\frac{\pi}{2} \leq \sin^{-1} x \leq \frac{\pi}{2}$$

$$\Rightarrow -\frac{\pi}{2} \leq 2 \sin^{-1} \alpha \leq \frac{\pi}{2}$$

$$\Rightarrow -\frac{\pi}{4} \leq \sin^{-1} \alpha \leq \frac{\pi}{4}$$

$$\Rightarrow \sin \left( -\frac{\pi}{4} \right) \leq \alpha \leq \sin \left( \frac{\pi}{4} \right)$$

$$\Rightarrow -\frac{1}{\sqrt{2}} \leq \alpha \leq \frac{1}{\sqrt{2}} \Rightarrow |\alpha| \leq \frac{1}{\sqrt{2}}$$

$$4. \text{ Given that, } \cos^{-1} x - \cos^{-1} \frac{y}{2} = \alpha$$

$$\Rightarrow \cos^{-1} \left( \frac{xy}{2} + \sqrt{1-x^2} \sqrt{1-\frac{y^2}{4}} \right) = \alpha$$

$$\Rightarrow \frac{xy}{2} + \sqrt{1-x^2} \sqrt{1-\frac{y^2}{4}} = \cos \alpha$$

$$\Rightarrow 2 \sqrt{1-x^2} \sqrt{1-\frac{y^2}{4}} = 2 \cos \alpha - xy$$

On squaring both sides, we get

$$\frac{4(1-x^2)(4-y^2)}{4} = 4 \cos^2 \alpha + x^2 y^2 - 4xy \cos \alpha$$

$$\Rightarrow 4 - 4x^2 - y^2 + x^2 y^2 = 4 \cos^2 \alpha + x^2 y^2 - 4xy \cos \alpha$$

$$\Rightarrow 4x^2 - 4xy \cos \alpha + y^2 = 4 \sin^2 \alpha$$

$$5. \text{ Since, } \sin^{-1} \left( \frac{x}{5} \right) + \operatorname{cosec}^{-1} \left( \frac{5}{4} \right) = \frac{\pi}{2}$$

$$\Rightarrow \sin^{-1} \left( \frac{x}{5} \right) + \sin^{-1} \left( \frac{4}{5} \right) = \frac{\pi}{2}$$

$$\Rightarrow \sin^{-1} \left( \frac{x}{5} \right) = \frac{\pi}{2} - \sin^{-1} \left( \frac{4}{5} \right)$$

$$\Rightarrow \sin^{-1} \left( \frac{x}{5} \right) = \cos^{-1} \left( \frac{4}{5} \right) \Rightarrow \sin^{-1} \left( \frac{x}{5} \right) = \sin^{-1} \left( \frac{3}{5} \right)$$

$$\Rightarrow x = 3$$

$$6. \text{ Since, } \operatorname{cosec}^{-1} \left( \frac{5}{3} \right) = \tan^{-1} \left( \frac{3}{4} \right)$$

$$\cos \left( \tan^{-1} \frac{3}{4} + \tan^{-1} \frac{2}{3} \right) = \cot \tan^{-1} \left[ \frac{\frac{3}{4} + \frac{2}{3}}{1 - \frac{1}{2}} \right]$$

$$= \cot \tan^{-1} \left[ \frac{\left( \frac{17}{12} \right)}{\left( \frac{1}{2} \right)} \right] = \cos \left[ \tan^{-1} \left( \frac{17}{6} \right) \right] = \frac{6}{17}$$



7.  $2y = x + z$

$$2 \tan^{-1} y = \tan^{-1} x + \tan^{-1} (z)$$

$$\Rightarrow \tan^{-1} \left( \frac{2y}{1-y^2} \right) = \tan^{-1} \left( \frac{x+z}{1-xz} \right)$$

$$\Rightarrow \frac{x+z}{1-y^2} = \frac{x+z}{1-xz}$$

$$\Rightarrow y^2 = xz \quad \text{or} \quad x+z=0$$

$$\Rightarrow x=y=z$$

**Case-II:**  $x < 0$

$$\text{Let } \cot^{-1} x = \theta \quad \therefore \theta \in \left( \frac{\pi}{2}, \pi \right)$$

$$\Rightarrow \cot \theta = x$$

$$\therefore \sin \theta = \frac{1}{\sqrt{1+x^2}}$$

$$\Rightarrow \sin^{-1} \sin \theta = \sin^{-1} \frac{1}{\sqrt{1+x^2}}$$

$$\Rightarrow \pi - \theta = \sin^{-1} \frac{1}{\sqrt{1+x^2}} \Rightarrow \theta = \pi - \sin^{-1} \frac{1}{\sqrt{1+x^2}}$$

Therefore,

$$\begin{aligned} \text{LHS} &= \begin{cases} \cos \tan^{-1} \sin \sin^{-1} \frac{1}{\sqrt{1+x^2}}, & \text{if } x \geq 0 \\ \cos \tan^{-1} \sin \left( \pi - \sin^{-1} \frac{1}{\sqrt{1+x^2}} \right), & \text{if } x < 0 \end{cases} \\ &= \cos \tan^{-1} \sin \sin^{-1} \frac{1}{\sqrt{1+x^2}}; \quad x \in \mathbb{R} = \cos \tan^{-1} \frac{1}{\sqrt{1+x^2}} \end{aligned}$$

$$\text{Let } \phi = \tan^{-1} \frac{1}{\sqrt{1+x^2}}$$

$$\text{As } \frac{1}{\sqrt{1+x^2}} \in (0, 1] \quad \therefore \phi \in \left( 0, \frac{\pi}{4} \right]$$

$$\therefore \tan \phi = \frac{1}{\sqrt{1+x^2}} \quad \therefore \cos \phi = \sqrt{\frac{1+x^2}{2+x^2}}$$

$$\therefore \text{LHS} = \cos \cos^{-1} \sqrt{\frac{1+x^2}{2+x^2}} = \sqrt{\frac{1+x^2}{2+x^2}} = \text{RHS.}$$

4.  $\sin \cot^{-1} (1+x) = \cos (\tan^{-1} x)$

$$\text{If } \alpha = \cot^{-1} (1+x) \quad \text{and} \quad \beta = \tan^{-1} x$$

$$\text{Then} \quad \frac{1}{\sqrt{x^2 + 2x + 2}} = \frac{1}{\sqrt{1+x^2}}$$

$$\Rightarrow x = -1/2$$

5.  $\sin^{-1} (ax) + \cos^{-1} y + \cos^{-1} (bxy) = \frac{\pi}{2}$

(A)  $a = 1, \quad b = 0$

$$\Rightarrow \sin^{-1}(x) + \cos^{-1}(y) + \cos^{-1}(0) = \frac{\pi}{2}$$

$$\Rightarrow \sin^{-1}x + \cos^{-1}y = 0$$

3. **Case-I:**  $x \geq 0$

$$\text{Let } \cot^{-1} x = \theta \quad \therefore \theta \in \left[ 0, \frac{\pi}{2} \right]$$

$$\Rightarrow x = \cot \theta$$

$$\therefore \sin \theta = \frac{1}{\sqrt{1+x^2}}$$

$$\Rightarrow \sin^{-1} \sin \theta = \sin^{-1} \frac{1}{\sqrt{1+x^2}}$$

$$\Rightarrow \theta = \sin^{-1} \frac{1}{\sqrt{1+x^2}}$$



## MATHS FOR JEE MAIN & ADVANCED

$$\Rightarrow \cos^{-1}y = -\sin^{-1}x$$

$$\Rightarrow \cos^{-1}y = \cos^{-1}\sqrt{1-x^2}$$

$$\Rightarrow x^2 + y^2 = 1$$

(B)  $\sin^{-1}(x) + \cos^{-1}y + \cos^{-1}(xy) = \frac{\pi}{2}$

$$\Rightarrow \cos^{-1}(y) + \cos^{-1}(xy) = \cos^{-1}x.$$

$$\Rightarrow \cos^{-1}\left(xy^2 - \sqrt{(1-y^2)(1-x^2y^2)}\right) = \cos^{-1}x.$$

$$\Rightarrow xy^2 - \sqrt{(1-y^2)(1-x^2y^2)} = x$$

$$\Rightarrow 1 - x^2 - y^2 + x^2y^2 = 0$$

$$\Rightarrow (1-x^2)(1-y^2) = 0$$

(C)  $\sin^{-1}(x) + \cos^{-1}y + \cos^{-1}(2xy) = \frac{\pi}{2}$

$$\Rightarrow \cos^{-1}\left(2xy^2 - \sqrt{(1-y^2)(1-4x^2y^2)}\right) = \cos^{-1}x.$$

$$\Rightarrow 2xy^2 - \sqrt{(1-y^2)(1-4x^2y^2)} = x$$

$$\Rightarrow 2xy^2 - x = \sqrt{(1-y^2)(1-4x^2y^2)}$$

$$\Rightarrow 4x^2y^4 + x^2 - 4x^2y^2 = 1 - y^2 - 4x^2y^2 + 4x^2y^4$$

$$\Rightarrow x^2 + y^2 = 1.$$

(D)  $\sin^{-1}(2x) + \cos^{-1}y + \cos^{-1}(2xy) = \frac{\pi}{2}$

$$\Rightarrow \cos^{-1}\left(2y^2x - \sqrt{(1-y^2)(1-4x^2y^2)}\right) = \cos^{-1}(2x)$$

$$\Rightarrow 2y^2x - \sqrt{1-y^2 - 4x^2y^2 + 4x^2y^4} = 2x.$$

$$\Rightarrow 1 - 4x^2 - y^2 + 4x^2y^2 = 0$$

$$\Rightarrow (1-4x^2)(1-y^2) = 0.$$

6.  $\sqrt{1+x^2}$

$$\left[ \left\{ x \cos \cos^{-1} \left( \frac{x}{\sqrt{1+x^2}} \right) + \sin \left( \sin^{-1} \frac{1}{\sqrt{1+x^2}} \right) \right\}^2 - 1 \right]^{1/2}$$

$$= \sqrt{1+x^2} \left[ \left( \frac{x^2}{\sqrt{1+x^2}} + \frac{1}{\sqrt{1+x^2}} \right)^2 - 1 \right]^{\frac{1}{2}}$$

$= \sqrt{1+x^2} \cdot x$  Hence (C) is correct.

7.  $\tan^{-1}\left(\frac{\sin \theta}{\sqrt{\cos 2\theta}}\right) = \sin^{-1}\left(\frac{\sin \theta}{\cos \theta}\right)$

$$\therefore f(\theta) = \tan \theta$$

$$\therefore \frac{df}{d \tan \theta} = 1$$



## MOCK TEST

1. (D) Since  $\cos^{-1}\left(\frac{4}{5}\right) = \tan^{-1}\left(\frac{3}{4}\right)$


$$\therefore \tan\left[\cos^{-1}\left(\frac{4}{5}\right) + \tan^{-1}\left(\frac{2}{3}\right)\right]$$

$$= \tan\left[\tan^{-1}\frac{3}{4} + \tan^{-1}\frac{2}{3}\right] = \frac{\frac{3}{4} + \frac{2}{3}}{1 - \frac{3}{4} \cdot \frac{2}{3}} = \frac{17}{6}$$

2.  $-1 \leq \frac{x^2}{4} + \frac{y^2}{9} \leq 1$  represents interior and the boundary

of the ellipse  $\frac{x^2}{4} + \frac{y^2}{9} = 1$  .....(i)

Also  $-1 \leq \frac{x}{2\sqrt{2}} + \frac{y}{3\sqrt{2}} - 2 \leq 1$



i.e.  $\frac{x}{2\sqrt{2}} + \frac{y}{3\sqrt{2}} \geq 1$  and  $\frac{x}{2\sqrt{2}} + \frac{y}{3\sqrt{2}} \leq 3$

$\frac{x}{2\sqrt{2}} + \frac{y}{3\sqrt{2}} \geq 1$  represents the portion of xy plane which contains only one point viz :

$$\left(\sqrt{2}, \frac{3}{\sqrt{2}}\right) \text{ of } \frac{x^2}{4} + \frac{y^2}{9} \leq 1$$

$$\therefore \sin^{-1}\left(\frac{x^2}{4} + \frac{y^2}{9}\right) + \cos^{-1}\left(\frac{x}{2\sqrt{2}} + \frac{y}{3\sqrt{2}} - 2\right)$$

$$= \sin^{-1}\left(\frac{1}{2} + \frac{1}{2}\right) + \cos^{-1}\left(\frac{1}{2} + \frac{1}{2} - 2\right)$$

3. (B)

$$\cos^{-1}(2x^2 - 1) = 2\pi - 2\cos^{-1}x \quad (\text{as } x < 0)$$

$$\cos^{-1}(2x^2 - 1) - 2\sin^{-1}x = 2\pi - 2\cos^{-1}x - 2\sin^{-1}x \\ = 2\pi - 2(\cos^{-1}x + \sin^{-1}x)$$

$$= 2\pi - 2 \cdot \frac{\pi}{2} = \pi$$

4. (C)

$$\sin^{-1}(x-1) \Rightarrow -1 \leq x-1 \leq 1 \Rightarrow 0 \leq x \leq 2$$

$$\cos^{-1}(x-3) \Rightarrow -1 \leq x-3 \leq 1 \Rightarrow 2 \leq x \leq 4$$

$$\tan^{-1}\left(\frac{x}{2-x^2}\right) \Rightarrow x \in \mathbb{R}, x \neq \sqrt{2}, -\sqrt{2}$$

$$\therefore x=2$$

$$\sin^{-1}(2-1) + \cos^{-1}(2-3) + \tan^{-1}\frac{2}{2-4} = \cos^{-1}k + \pi$$

$$\Rightarrow \sin^{-1}1 + \cos^{-1}(-1) + \tan^{-1}(-1) = \cos^{-1}k + \pi$$

$$\frac{\pi}{2} + \pi - \frac{\pi}{4} = \cos^{-1}k + \pi$$

$$\Rightarrow \cos^{-1}k = \frac{\pi}{4} \Rightarrow k = \frac{1}{\sqrt{2}}$$

5. (A)

$$\text{Let } \sin^{-1}a = A,$$

$$\sin^{-1}b = B$$

$$\sin^{-1}c = C$$

$$\therefore \sin A = a, \sin B = b, \sin C = c$$

and  $A + B + C = \pi$ , then

$$\sin 2A + \sin 2B + \sin 2C = 4 \sin A \sin B \sin C$$

$$\text{Now } a\sqrt{1-a^2} + b\sqrt{1-b^2} + c\sqrt{1-c^2}$$

$$= \sin A \cos A + \sin B \cos B + \sin C \cos C$$

$$= \frac{1}{2} [\sin 2A + \sin 2B + \sin 2C] = 2 \sin A \sin B \sin C = 2abc$$

6. (C)

$$0 \leq \{x\} < 1 \quad \text{i.e. } -1 < -\{x\} \leq 0$$

$$\therefore \frac{\pi}{2} \leq \cos^{-1}(-\{x\}) < \pi$$

$$\therefore \text{the range is } \left[\frac{\pi}{2}, \pi\right)$$

7. (B)

$$\text{We have } 2 \tan^{-1}(\cos x) = \tan^{-1}(2 \operatorname{cosec} x)$$

$$\Rightarrow \tan(2 \tan^{-1} \cos x) = 2 \operatorname{cosec} x$$

$$\Rightarrow \frac{2 \cos x}{1 - \cos^2 x} = 2 \operatorname{cosec} x \Rightarrow \frac{2 \cos x}{\sin^2 x} = 2 \operatorname{cosec} x$$

$$\Rightarrow \sin x = \cos x \Rightarrow x = \frac{\pi}{4}.$$



## MATHS FOR JEE MAIN & ADVANCED

8. (C)

$$\begin{aligned} \log_{\frac{1}{2}} \sin^{-1} x &> \log_{\frac{1}{2}} \cos^{-1} x \\ \Leftrightarrow \cos^{-1} x &> \sin^{-1} x, \quad 0 < x < 1 \\ \Leftrightarrow \cos^{-1} x &> \frac{\pi}{2} - \cos^{-1} x, \quad 0 < x < 1 \\ \Leftrightarrow \cos^{-1} x &> \frac{\pi}{4}, \quad 0 < x < 1 \\ \Leftrightarrow 0 < x &< \frac{1}{\sqrt{2}} \end{aligned}$$

9. (A)

$$\begin{aligned} S_1 : \sin^{-1} x - \frac{\pi}{2} + \sin^{-1}(-x) &= \frac{\pi}{2} \\ \Rightarrow \sin^{-1}(x) + \sin^{-1}(-x) &= \pi \\ 0 = \pi &\text{ which is not possible} \\ \therefore \text{no solution} & \end{aligned}$$

$$\begin{aligned} S_2 : \sin^{-1}(x^2 + 4x + 3) + \cos^{-1}(x^2 + 6x + 8) &= \frac{\pi}{2} \\ &= \sin^{-1}(x^2 + 4x + 3) + \cos^{-1}(x^2 + 4x + 3) \\ \Rightarrow x^2 + 6x + 8 &= x^2 + 4x + 3 \\ \Rightarrow 2x &= -5 \quad \Rightarrow x = -\frac{5}{2} \\ \nexists x^2 + 4x + 3 &= (x+2)^2 - 1 \in [-1, 1] \text{ at } x = -\frac{5}{2} \\ \& x^2 + 6x + 8 = (x+3)^2 - 1 \in [-1, 1] \text{ at } x = -\frac{5}{2} \\ \therefore x &= -\frac{5}{2} \\ S_3 : \sin^{-1}\{\cos(\sin^{-1}x)\} + \cos^{-1}\{\sin(\cos^{-1}x)\} &= \frac{\pi}{2} \\ \{ \text{As } \cos(\sin^{-1}x) = \sin(\cos^{-1}x) = \sqrt{1-x^2} \} & \end{aligned}$$

$$\begin{aligned} S_4 : 2 \left[ \tan^{-1} \frac{1+2}{1-2} + \pi + \tan^{-1} 3 \right] \\ = 2[\pi - \tan^{-1} 3 + \tan^{-1} 3] = 2\pi \end{aligned}$$

10. (C)

$$\sin^{-1} \sin 5 = \sin^{-1} \sin(5 - 2\pi) = 5 - 2\pi$$

$$\left( \text{As } -\frac{\pi}{2} \leq 5 - 2\pi \leq \frac{\pi}{2} \right)$$

$$\therefore \sin^{-1} \sin 5 > x^2 - 4x$$

$$\Rightarrow 5 - 2\pi > x^2 - 4x \quad \Rightarrow x^2 - 4x + 2\pi - 5 < 0$$

sign sum of  $(x^2 - 4x + 2\pi - 5)$



$$2 - \sqrt{9 - 2\pi} < x < 2 + \sqrt{9 - 2\pi}$$

Integral values of x are 1, 2, 3

Number of integral value of x = 3

11. Let  $\tan^{-1} x = \theta$ .

$$\text{Then } -\frac{\pi}{2} < \theta < \frac{\pi}{2}, \quad \theta \neq \pm \frac{\pi}{4} \text{ and } x = \tan \theta$$

$$\tan^{-1} x + \tan^{-1} \frac{2x}{1-x^2} = \theta + \tan^{-1} \frac{2 \tan \theta}{1 - \tan^2 \theta}$$

$$= \theta + \tan^{-1} (\tan 2\theta), \text{ where } -\pi < 2\theta < \pi, 2\theta \neq \pm \frac{\pi}{2}$$

$$= \begin{cases} \theta + \pi + 2\theta & \text{when } -\pi < 2\theta < -\frac{\pi}{2} \\ \theta + 2\theta & \text{when } -\frac{\pi}{2} < 2\theta < \frac{\pi}{2} \\ \theta - \pi + 2\theta & \text{when } \frac{\pi}{2} < 2\theta < \pi \end{cases}$$

$$= \begin{cases} \pi + 3\theta & \text{when } -\frac{\pi}{2} < \theta < -\frac{\pi}{4} \\ 3\theta & \text{when } -\frac{\pi}{4} < \theta < \frac{\pi}{4} \\ -\pi + 3\theta & \text{when } \frac{\pi}{4} < \theta < \frac{\pi}{2} \end{cases}$$

$$= \begin{cases} \pi + 3 \tan^{-1} x & \text{when } x < -1 \\ 3 \tan^{-1} x & \text{when } -1 < x < 1 \\ -\pi + 3 \tan^{-1} x & \text{when } 1 < x \end{cases}$$

12. (A,C)

$$\text{Domain of } f(x) = \bullet n \cos^{-1} x$$

$$\text{is } x \in [-1, 1]$$

$$\therefore [\alpha] = -1 \quad \text{or} \quad 0$$

$$13. \text{ If } -1 \leq x < 0, \text{ then } -\frac{\pi}{2} \leq \sin^{-1} x < 0$$

$$\text{Also } 0 < 2 \cot^{-1}(y^2 - 2y) < 2\pi$$



Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035

+91-9350679141

$$\therefore -\frac{\pi}{2} < \sin^{-1}x + 2\cot^{-1}(y^2 - 2y) < 2\pi$$

$\therefore$  there is no solution in this case.

thus x can not be negative .....(i)

$$\text{Now if } x \geq 0, \text{ then } 0 \leq \sin^{-1}x \leq \frac{\pi}{2}$$

$$\Rightarrow \frac{3\pi}{4} \leq \cot^{-1}(y^2 - 2y) < \pi$$

$$\Rightarrow y^2 - 2y \leq -1 \Rightarrow y = 1$$

$$\text{since for } y=1, \text{ we have } 2\cot^{-1}(y^2 - 2y) = 2\cot^{-1}(-1) = \frac{3\pi}{2}$$

$$\therefore \sin^{-1}x = \frac{\pi}{2} \text{ i.e. } x=1$$

$\therefore$  the solution is  $x = 1, y = 1$

14. (A, B, C)

$$(A) \sin\left(\tan^{-1}3 + \tan^{-1}\frac{1}{3}\right) = \sin\frac{\pi}{2} = 1$$

$$(B) \cos\left(\frac{\pi}{2} - \sin^{-1}\frac{3}{4}\right) = \cos\left(\cos^{-1}\frac{3}{4}\right) = \frac{3}{4}$$

$$(C) \sin\left(\frac{1}{4}\sin^{-1}\frac{\sqrt{63}}{8}\right)$$

$$\text{Let } \sin^{-1}\frac{\sqrt{63}}{8} = \theta$$

$$\text{so } \sin\theta = \frac{\sqrt{63}}{8} \quad \text{if } \cos\theta = \frac{1}{8}$$

$$\text{we have } \cos\frac{\theta}{2} = \sqrt{\frac{1+\cos\theta}{2}} = \frac{3}{4}$$

$$\sin\frac{\theta}{4} = \sqrt{\frac{1-\cos\frac{\theta}{2}}{2}} = \frac{1}{2\sqrt{2}}$$

$$\text{Now } \log_2 \sin\left(\frac{1}{4}\sin^{-1}\frac{\sqrt{63}}{8}\right) = \log_2 \frac{1}{2\sqrt{2}} = -\frac{3}{2}$$

$$(D) \cos^{-1}\frac{\sqrt{5}}{3} = \theta$$

$$\cos\theta = \frac{\sqrt{5}}{3}$$

$$\therefore \tan\frac{\theta}{2} = \frac{3-\sqrt{5}}{2} \text{ which is irrational}$$

$$15. \sin^{-1}x + \sin^{-1}(1-x) = \cos^{-1}x$$

$$\Rightarrow \frac{\pi}{2} - \cos^{-1}x + \frac{\pi}{2} - \cos^{-1}(1-x) \\ = \cos^{-1}x$$

$$\Rightarrow 2\cos^{-1}x = \pi - \cos^{-1}(1-x)$$

$$\Rightarrow \cos^{-1}(2x^2 - 1) = \cos^{-1}(x-1) \Rightarrow 2x^2 - 1 = x - 1$$

$$\Rightarrow x(2x-1) = 0 \Rightarrow x = 0, \frac{1}{2}$$

$$17. \text{ Range of } f \text{ is } \left\{\frac{\pi}{2}\right\} \text{ and domain of } f \text{ is } \{0\}.$$

Hence if domain of  $f$  is singleton then range has to be a singleton.

If S-2 and S-1 are reverse then the answer will be B.

18. (A)

(Moderate)

$$\sin^{-1}x = \tan^{-1}\frac{x}{\sqrt{1-x^2}} > \tan^{-1}x > \tan^{-1}y$$

$$\{x > y, \frac{x}{\sqrt{1-x^2}} > x\}$$

Statement-II is true

$$e < \pi$$

$$\frac{1}{\sqrt{e}} > \frac{1}{\sqrt{\pi}}$$

by Statement-II

$$\sin^{-1}\left(\frac{1}{\sqrt{e}}\right) > \tan^{-1}\left(\frac{1}{\sqrt{e}}\right) > \tan^{-1}\left(\frac{1}{\sqrt{\pi}}\right)$$

Statement-I is true



## MATHS FOR JEE MAIN & ADVANCED

**19. (A)**

$$f(x) = \sin^{-1} \left( \frac{2x}{1+x^2} \right) = \pi - 2 \tan^{-1} x, x \geq 1$$

$$f'(x) = -\frac{2}{1+x^2} \Rightarrow f'(2) = -\frac{2}{5}$$

Statement-I is True, Statement-II is True ; Statement-II is a correct explanation for statement-I.

**20. (A)**

$$\operatorname{cosec}^{-1} x > \sec^{-1} x$$

$$\operatorname{cosec}^{-1} x > \frac{\pi}{2} - \operatorname{cosec}^{-1} x$$

$$\operatorname{cosec}^{-1} x > \frac{\pi}{4}$$

$$1 \leq x < \sqrt{2} \quad \text{and} \quad \left( \frac{1}{2} + \frac{1}{\sqrt{2}} \right) \in [1, \sqrt{2})$$

Statement 2 is true and explains statement 1

**21. (A)  $\rightarrow$  (r), (B)  $\rightarrow$  (s), (C)  $\rightarrow$  (p), (D)  $\rightarrow$  (q)**

(A) The difference  $= 2 - (-2) = 4$

(B) Let  $f(x) = x^2 - 4x + 3$

$$f'(x) = 2x - 4 = 0 \Rightarrow x = 2$$

$$f(1) = 0, f(2) = -1, f(3) = 0$$

$$\therefore |\text{greatest value} - \text{least value}| = 1$$

$$(C) \tan^{-1} \frac{1-x}{1+x} = \tan^{-1} 1 - \tan^{-1} x$$

$$\therefore \text{greatest value} = \frac{\pi}{4}$$

$$(D) \therefore \text{greatest value} = \frac{\pi}{2}, \text{least value} = \frac{\pi}{3}$$

$$\therefore \text{difference} = \frac{\pi}{6}$$

**22. (A)  $\rightarrow$  (q, s), (B)  $\rightarrow$  (r, s, t), (C)  $\rightarrow$  (r, s), (D)  $\rightarrow$  (p, q)**

$$(A) \text{ Given } \sin^{-1} x - \cos^{-1} x = \frac{\pi}{6}$$

$$\text{Also, } \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$$

$$\text{Solving } x = \frac{\sqrt{3}}{2}$$

$$(B) \sec^2(\tan^{-1} 2) + \operatorname{cosec}^2(\cot^{-1} 3)$$

$$= 1 + (\tan(\tan^{-1} 2))^2 + 1 + (\cot(\cot^{-1} 3))^2 \\ = 15$$

$$(C) \text{ Given eqn. is } \frac{\pi}{2} - 2 \cos^{-1} x = \sin^{-1}(3x - 2)$$

$$\text{or } 3x - 2 = \cos(2 \cos^{-1} x) = 2 \cos^2(\cos^{-1} x) - 1 \\ = 2x^2 - 1$$

$$\Rightarrow 2x^2 - 3x + 1 = 0 \Rightarrow x = 1 \text{ or } \frac{1}{2}$$

$$(D) \sin 5 = \sin(5 - 2\pi)$$

$$\Rightarrow \sin^{-1}(\sin 5) = \sin^{-1}(\sin(5 - 2\pi)) \\ = 5 - 2\pi$$

**24.**

**1. (B)**

$$\sin^{-1} \left( \frac{4x}{x^2 + 4} \right) + 2 \tan^{-1} \left( -\frac{x}{2} \right)$$

$$= \sin^{-1} \left( \frac{\frac{2x}{2}}{\left( \frac{x}{2} \right)^2 + 1} \right) - 2 \tan^{-1} \frac{x}{2}$$

$$= 2 \tan^{-1} \frac{x}{2} - 2 \tan^{-1} \frac{x}{2} = 0$$

$$\text{Here } \left| \frac{x}{2} \right| \leq 1$$

$$|x| \leq 2 \Rightarrow -2 \leq x \leq 2$$

**2. (A)**

$$\cos^{-1} \frac{6x}{1+9x^2} = -\frac{\pi}{2} + 2 \tan^{-1} 3x$$

$$\Rightarrow \frac{\pi}{2} - \sin^{-1} \frac{6x}{1+9x^2} = -\frac{\pi}{2} + 2 \tan^{-1} 3x$$

$$\Rightarrow \sin^{-1} \frac{6x}{1+9x^2} = \pi - 2 \tan^{-1} 3x$$

$$\Rightarrow \sin^{-1} \frac{2 \cdot 3x}{1+(3x)^2} = \pi - 2 \tan^{-1} 3x$$

Above is true when  $3x > 1$



Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035

+91-9350679141

## INVERSE TRIGONOMETRIC FUNCTION

$$\Rightarrow x > \frac{1}{3}$$

$$x \in \left( \frac{1}{3}, \infty \right)$$

**3. (C)**

$$(x-1)(x^2+1) > 0$$

$$\Rightarrow x > 1$$

$$\therefore \sin \left[ \frac{1}{2} \tan^{-1} \left( \frac{2x}{1-x^2} \right) - \tan^{-1} x \right]$$

$$= \sin \left[ \frac{1}{2}(-\pi + 2 \tan^{-1} x) - \tan^{-1} x \right] = \sin \left( -\frac{\pi}{2} \right) = -1$$

**25.**

**1. (B)**

$$A = (\tan^{-1} x)^3 + (\cot^{-1} x)^3$$

$$A = (\tan^{-1} x + \cot^{-1} x)^3 - 3 \tan^{-1} x \cot^{-1} x (\tan^{-1} x + \cot^{-1} x)$$

$$\Rightarrow A = \left( \frac{\pi}{2} \right)^3 - 3 \tan^{-1} x \cot^{-1} x \cdot \frac{\pi}{2}$$

$$\Rightarrow A = \frac{\pi^3}{8} - \frac{3\pi}{2} \tan^{-1} x \left( \frac{\pi}{2} - \tan^{-1} x \right)$$

$$\Rightarrow A = \frac{\pi^3}{32} + \frac{3\pi}{2} \left( \tan^{-1} x - \frac{\pi}{4} \right)^2$$

as  $x > 0$

$$\frac{\pi^3}{32} \leq A < \frac{\pi^3}{8}$$

**2. (C)**

$$B = (\sin^{-1} t)^2 + (\cos^{-1} t)^2$$

$$B = (\sin^{-1} t + \cos^{-1} t)^2 - 2 \sin^{-1} t \cos^{-1} t$$

$$B = \frac{\pi^2}{4} - 2 \sin^{-1} t \left( \frac{\pi}{2} - \sin^{-1} t \right)$$

$$B = \frac{\pi^2}{8} + 2 \left( \sin^{-1} t - \frac{\pi}{4} \right)^2$$

$$B_{\max} = \frac{\pi^2}{8} + 2 \cdot \frac{\pi^2}{16} = \frac{\pi^2}{4}$$

**3. (A)**

$$\lambda = \frac{\pi^3}{32} \quad \mu = \frac{\pi^2}{4}$$

$$\frac{\lambda}{\mu} = \frac{\pi}{8}$$

$$\frac{\lambda - \mu\pi}{\mu} = \frac{\pi}{8} - \pi = \frac{-7\pi}{8}$$

$$\cot^{-1} \cot \left( \frac{\lambda - \mu\pi}{\mu} \right) = \cot^{-1} \cot \left( -\frac{7\pi}{8} \right) = \frac{\pi}{8}$$

**26. (1)**

$$(\tan^{-1} x)^2 + (\cot^{-1} x)^2 = (\tan^{-1} x + \cot^{-1} x)^2 - 2\tan^{-1} x$$

$$\left( \frac{\pi}{2} - \tan^{-1} x \right)^2 = \frac{\pi^2}{4} - \pi \tan^{-1} x + 2(\tan^{-1} x)^2 = \frac{5\pi^2}{8}$$

$$\therefore \tan^{-1} x = \frac{2\pi}{3}, -\frac{\pi}{4}$$

$$\tan^{-1} x = -\frac{\pi}{4} \quad \{ \tan^{-1} x \neq \frac{2\pi}{3} \}$$

$\therefore x = -1$  is the solution

**27. (1)**

$$\tan^{-1} \left[ \frac{3 \sin 2\alpha}{5 + 3 \cos 2\alpha} \right] + \tan^{-1} \left[ \frac{\tan \alpha}{4} \right]$$

$$= \tan^{-1} \left( \frac{6 \tan \alpha}{8 + 2 \tan^2 \alpha} \right) + \tan^{-1} \left( \frac{\tan \alpha}{4} \right)$$

$$= \tan^{-1} \left( \frac{\frac{3 \tan \alpha}{4 + \tan^2 \alpha} + \frac{\tan \alpha}{4}}{1 - \frac{3 \tan^2 \alpha}{16 + 4 \tan^2 \alpha}} \right)$$

$$\left\{ Q \frac{3 \tan^2 \alpha}{16 + 4 \tan^2 \alpha} < 1 \right\}$$

$$= \tan^{-1} (\tan \alpha) = \alpha$$

$$28. \tan \left( \frac{\pi}{4} + \alpha \right) \quad \text{when } \alpha = \tan^{-1} \left( \frac{\frac{1}{4} + \frac{1}{5}}{1 - \frac{1}{20}} \right);$$

$$\alpha = \tan^{-1} \left( \frac{9}{19} \right) = \frac{1 + \frac{9}{19}}{1 - \frac{9}{19}} = \frac{28}{10} = \frac{14}{5} = \frac{a}{b}$$

$$\Rightarrow 14 - 5 = 9$$



Add. 41-42A, Ashok Park Main, New Rohtak Road, New Delhi-110035  
+91-9350679141

29.  $\cos^{-1}(2x) + \cos^{-1}(3x) = \pi - \cos^{-1}(x) = \cos^{-1}(-x)$

$$\cos^{-1}[(2x)(3x) - \sqrt{1-4x^2}\sqrt{1-9x^2}] = \cos^{-1}(-x)$$

$$6x^2 - \sqrt{1-4x^2} \cdot \sqrt{1-9x^2} = -x$$

$$(6x^2 + x)^2 = (1 - 4x^2)(1 - 9x^2)$$

$$\Rightarrow x^2 + 12x^3 = 1 - 13x^2$$

$$\Rightarrow 12x^3 + 14x^2 - 1 = 0$$

$$\therefore a = 12; b = 14; c = 0$$

$$\Rightarrow -a + b + c = 2$$

