

SETS

- Given the sets $A = \{1, 2, 3\}$, $B = \{3, 4\}$, $C = \{4, 5, 6\}$, then $A \cup (B \cap C)$ is
 (A) $\{3\}$ (B) $\{1, 2, 3, 4\}$
 (C) $\{1, 2, 4, 5\}$ (D) $\{1, 2, 3, 4, 5, 6\}$
- If A and B are any two sets, then $A \cup (A \cap B)$ is equal to
 (A) A (B) B
 (C) A^c (D) B^c
- If A and B are two given sets, then $A \cap (A \cap B)^c$ is equal to
 (A) A (B) B
 (C) \varnothing (D) $A \cap B^c$
- If the sets A and B are defined as $A = \{(x, y) : y = \frac{1}{x}, 0 \neq x \in R\}$, $\bigcup_{i=1}^{30} A_i = \bigcup_{j=1}^n B_j$, then
 (A) $A \cap B = A$ (B) $A \cap B = B$
 (C) $A \cap B = \varnothing$ (D) None of these
- Let $A = \{x : x \in R, |x| < 1\}$; $B = \{x : x \in R, |x-1| \geq 1\}$ and $A \cup B = R - D$, then the set D is
 (A) $[x : 1 < x \leq 2]$ (B) $[x : 1 \leq x < 2]$
 (C) $[x : 1 \leq x \leq 2]$ (D) None of these
- If the sets A and B are defined as $A = \{(x, y) : y = e^x, x \in R\}$; $B = \{(x, y) : y = x, x \in R\}$, then
 (A) $B \subseteq A$ (B) $A \subseteq B$
 (C) $A \cap B = \varnothing$ (D) $A \cup B = A$
- If $X = \{4^n - 3n - 1 : n \in N\}$ and $Y = \{9(n-1) : n \in N\}$, then $X \cup Y$ is equal to
 (A) X (B) Y
 (C) N (D) None of these
- Let $n(U) = 700$, $n(A) = 200$, $n(B) = 300$ and $n(A \cap B) = 100$, then $n(A^c \cap B^c) =$
 (A) 400 (B) 600
 (C) 300 (D) 200
- In a town of 10,000 families it was found that 40% family buy newspaper A , 20% buy newspaper B and 10% families buy newspaper C , 5% families buy A and B , 3% buy B and C and 4% buy A and C . If 2% families buy all the three newspapers, then number of families which buy A only is
 (A) 3100 (B) 3300
 (C) 2900 (D) 1400
- In a city 20 percent of the population travels by car, 50 percent travels by bus and 10 percent travels by both car and bus. Then persons travelling by car or bus is
 (A) 80 percent (B) 40 percent
 (C) 60 percent (D) 70 percent
- If $Q = \left\{x : x = \frac{1}{y}, \text{ where } y \in N\right\}$, then
 (A) $0 \in Q$ (B) $1 \in Q$
 (C) $2 \in Q$ (D) $\frac{2}{3} \in Q$
- Which set is the subset of all given sets
 (A) $\{1, 2, 3, 4, \dots\}$ (B) $\{1\}$
 (C) $\{0\}$ (D) $\{\}$
- Let $S = \{0, 1, 5, 4, 7\}$. Then the total number of subsets of S is
 (A) 64 (B) 32
 (C) 40 (D) 20
- The number of non-empty subsets of the set $\{1, 2, 3, 4\}$ is
 (A) 15 (B) 14
 (C) 16 (D) 17
- The smallest set A such that $A \cup \{1, 2\} = \{1, 2, 3, 5, 9\}$ is
 (A) $\{2, 3, 5\}$ (B) $\{3, 5, 9\}$
 (C) $\{1, 2, 5, 9\}$ (D) None of these
- If $A \cap B = B$, then
 (A) $A \subset B$ (B) $B \subset A$
 (C) $A = \varnothing$ (D) $B = \varnothing$

17. If A and B are two sets, then $A \cup B = A \cap B$ if
 (A) $A \subseteq B$ (B) $B \subseteq A$
 (C) $A = B$ (D) None of these
18. Let A and B be two sets. Then
 (A) $A \cup B \subseteq A \cap B$ (B) $A \cap B \subseteq A \cup B$
 (C) $A \cap B = A \cup B$ (D) None of these
19. Let $A = \{(x, y) : y = e^x, x \in R\}$,
 $B = \{(x, y) : y = e^{-x}, x \in R\}$. Then
 (A) $A \cap B = \varnothing$ (B) $A \cap B \neq \varnothing$
 (C) $A \cup B = R^2$ (D) None of these
20. If $A = \{2, 3, 4, 8, 10\}$, $B = \{3, 4, 5, 10, 12\}$,
 $C = \{4, 5, 6, 12, 14\}$ then $(A \cap B) \cup (A \cap C)$
 is equal to
 (A) $\{3, 4, 10\}$ (B) $\{2, 8, 10\}$
 (C) $\{4, 5, 6\}$ (D) $\{3, 5, 14\}$
21. If A and B are two sets then $(A - B) \cup (B - A)$
 $\cup (A \cap B)$ is equal to
 (A) $A \cup B$ (B) $A \cap B$
 (C) A (D) B'
22. Let A and B be two sets then $(A \cup B)' \cup (A' \cap B)$
 is equal to
 (A) A' (B) A
 (C) B' (D) None of these
23. Let U be the universal set and $A \cup B \cup C = U$.
 Then $\{(A - B) \cup (B - C) \cup (C - A)\}'$ is equal to
 (A) $A \cup B \cup C$ (B) $A \cup (B \cap C)$
 (C) $A \cap B \cap C$ (D) $A \cap (B \cup C)$
24. If $n(A) = 3$, $n(B) = 6$ and $A \subseteq B$. Then the
 number of elements in $A \cup B$ is equal to
 (A) 3 (B) 9
 (C) 6 (D) None of these
25. Let A and B be two sets such that
 $n(A) = 0.16$, $n(B) = 0.14$, $n(A \cup B) = 0.25$.
 Then $n(A \cap B)$ is equal to
 (A) 0.3 (B) 0.5
 (C) 0.05 (D) None of these
26. If A and B are disjoint, then $n(A \cup B)$ is equal to
 (A) $n(A)$ (B) $n(B)$
 (C) $n(A) + n(B)$ (D) $n(A) \cdot n(B)$
27. If A and B are not disjoint sets, then $n(A \cup B)$
 is equal to
 (A) $n(A) + n(B)$ (B) $n(A) + n(B) - n(A \cap B)$
 (C) $n(A) + n(B) + n(A \cap B)$ (D) $n(A)n(B)$
28. In a battle 70% of the combatants lost one eye,
 80% an ear, 75% an arm, 85% a leg, $x\%$ lost
 all the four limbs. The minimum value of x is
 (A) 10 (B) 12
 (C) 15 (D) None of these
29. Out of 800 boys in a school, 224 played
 cricket, 240 played hockey and 336 played
 basketball. Of the total, 64 played both
 basketball and hockey; 80 played cricket and
 basketball and 40 played cricket and hockey;
 24 played all the three games. The number of
 boys who did not play any game is
 (A) 128 (B) 216
 (C) 240 (D) 160
30. A survey shows that 63% of the Americans like
 cheese whereas 76% like apples. If $x\%$ of the
 Americans like both cheese and apples, then
 (A) $x = 39$ (B) $x = 63$
 (C) $39 \leq x \leq 63$ (D) None of these
31. 20 teachers of a school either teach
 mathematics or physics. 12 of them teach
 mathematics while 4 teach both the subjects.
 Then the number of teachers teaching physics
 only is
 (A) 12 (B) 8
 (C) 16 (D) None of these
32. Of the members of three athletic teams in a
 school 21 are in the cricket team, 26 are in the
 hockey team and 29 are in the football team.
 Among them, 14 play hockey and cricket, 15
 play hockey and football, and 12 play football
 and cricket. Eight play all the three games.
 The total number of members in the three
 athletic teams is
 (A) 43 (B) 76
 (C) 49 (D) None of these

33. In a class of 100 students, 55 students have passed in Mathematics and 67 students have passed in Physics. Then the number of students who have passed in Physics only is
 (A) 22 (B) 33
 (C) 10 (D) 45
34. If A and B are two sets, then $A \times B = B \times A$ if
 (A) $A \subseteq B$ (B) $B \subseteq A$
 (C) $A = B$ (D) None of these
35. If A and B be any two sets, then $(A \cap B)'$ is equal to
 (A) $A' \cap B'$ (B) $A' \cup B'$
 (C) $A \cap B$ (D) $A \cup B$
36. If $(1, 3)$, $(2, 5)$ and $(3, 3)$ are three elements of $A \times B$ and the total number of elements in $A \times B$ is 6, then the remaining elements of $A \times B$ are
 (A) $(1, 5)$; $(2, 3)$; $(3, 5)$
 (B) $(5, 1)$; $(3, 2)$; $(5, 3)$
 (C) $(1, 5)$; $(2, 3)$; $(5, 3)$
 (D) None of these
37. $A = \{1, 2, 3\}$ and $B = \{3, 8\}$, then $(A \cup B) \times (A \cap B)$ is
 (A) $\{(3, 1), (3, 2), (3, 3), (3, 8)\}$
 (B) $\{(1, 3), (2, 3), (3, 3), (8, 3)\}$
 (C) $\{(1, 2), (2, 2), (3, 3), (8, 8)\}$
 (D) $\{(8, 3), (8, 2), (8, 1), (8, 8)\}$
38. If $A = \{2, 3, 5\}$, $B = \{2, 5, 6\}$, then $(A - B) \times (A \cap B)$ is
 (A) $\{(3, 2), (3, 3), (3, 5)\}$
 (B) $\{(3, 2), (3, 5), (3, 6)\}$
 (C) $\{(3, 2), (3, 5)\}$
 (D) None of these
39. In a class of 30 pupils, 12 take needle work, 16 take physics and 18 take history. If all the 30 students take at least one subject and no one takes all three then the number of pupils taking 2 subjects is
 (A) 16 (B) 6
 (C) 8 (D) 20
40. The number of elements in the set $\{(a, b) : 2a^2 + 3b^2 = 35, a, b \in Z\}$, where Z is the set of all integers, is
 (A) 2 (B) 4
 (C) 8 (D) 12
41. If $A = \{1, 2, 3, 4\}$; $B = \{a, b\}$ and f is a mapping such that $f : A \rightarrow B$, then $A \times B$ is
 (A) $\{(a, 1), (3, b)\}$
 (B) $\{(a, 2), (4, b)\}$
 (C) $\{(1, a), (1, b), (2, a), (2, b), (3, a), (3, b), (4, a), (4, b)\}$
 (D) None of these
42. If $A = \{1, 2, 3, 4, 5\}$, $B = \{2, 4, 6\}$, $C = \{3, 4, 6\}$, then $(A \cup B) \cap C$ is
 (A) $\{3, 4, 6\}$ (B) $\{1, 2, 3\}$
 (C) $\{1, 4, 3\}$ (D) None of these
43. If $A = \{x, y\}$ then the power set of A is
 (A) $\{x^x, y^y\}$
 (B) $\{\phi, x, y\}$
 (C) $\{\phi, \{x\}, \{2y\}\}$
 (D) $\{\phi, \{x\}, \{y\}, \{x, y\}\}$
44. A set contains $2n+1$ elements. The number of subsets of this set containing more than n elements is equal to
 (A) 2^{n-1} (B) 2^n
 (C) 2^{n+1} (D) 2^{2n}
45. Which of the following is a true statement
 (A) $\{a\} \in \{a, b, c\}$ (B) $\{a\} \subseteq \{a, b, c\}$
 (C) $\phi \in \{a, b, c\}$ (D) None of these
46. If $A = \{x : x \text{ is a multiple of } 4\}$ and $B = \{x : x \text{ is a multiple of } 6\}$ then $A \subset B$ consists of all multiples of
 (A) 16 (B) 12
 (C) 8 (D) 4

47. A class has 175 students. The following data shows the number of students obtaining one or more subjects. Mathematics 100, Physics 70, Chemistry 40; Mathematics and Physics 30, Mathematics and Chemistry 28, Physics and Chemistry 23; Mathematics, Physics and Chemistry 18. How many students have offered Mathematics alone

- (A) 35 (B) 48
(C) 60 (D) 22

48. Consider the following relations :

- (1) $A - B = A - (A \cap B)$
(2) $A = (A \cap B) \cup (A - B)$
(3) $A - (B \cup C) = (A - B) \cup (A - C)$

which of these is/are correct

- (A) 1 and 3 (B) 2 only
(C) 2 and 3 (D) 1 and 2

49. If two sets A and B are having 99 elements in common, then the number of elements common to each of the sets $A \times B$ and $B \times A$ are

- (A) 2^{99} (B) 99^2
(C) 100 (D) 18

50. Given $n(U) = 20$, $n(A) = 12$, $n(B) = 9$, $n(A \cap B) = 4$, where U is the universal set, A and B are subsets of U , then $n((A \cup B)^c) =$

- (A) 17 (B) 9
(C) 11 (D) 3