


Single Choice

-
- Protium Deuterium Tritium


Orthohydrogen Parahydrogen
(Isomers of hydrogen)

oxidation no. of hydrogen in hydrides is -1 .

CHEMISTRY FOR JEE MAIN & ADVANCED



23. In the reaction with calcium hydrogen acts as an oxidising agent.




25. Hydrogen does not combine with helium.

26. Saline hydride means salt like hydride. Alkali metals and alkaline earth metals and some highly +ve members of lanthanide series can transfer electron easily to hydrogen atoms. Ex. NaH, KH, CaH_2 (Si, B, and Al do not form salt like hydride).

27. $\text{Ti H}_{1.5-1.8}$ is an interstitial hydride.


28. In CaH_2 H has oxidation state $(-1) [+2 + 2x = 0, x = -1]$

29. Transition elements like Ni, Pt, Pd adsorb hydrogen.



33. Hydrogen is liberated at anode $2\text{H}^- \longrightarrow \text{H}_{2(\text{g})} + 2\text{e}^-$

34. Temporary hardness of water is due to the presence of $\text{Ca}(\text{HCO}_3)_2$ and $\text{Mg}(\text{HCO}_3)_2$ in water



37. Hard water contains soluble salts CaCl_2 , MgCl_2 , CaSO_4 , MgSO_4 , $\text{Ca}(\text{HCO}_3)_2$, $\text{Mg}(\text{HCO}_3)_2$

38. Freezing point of heavy water is 3.8°C .

39. Slowing down the speed of high energy neutrons.




41. Heavy water contains heavy hydrogen (^2H), formula of heavy water is $^2\text{H}_2\text{O}$.

42. Hydrated silicates of Al and Na is called permutit $\text{Na}_2\text{Al}_2\text{Si}_2\text{O}_8 \cdot x\text{H}_2\text{O}$

43. Cation exchange resin exchanges Ca^{2+} , Mg^{2+} ions from water, and water becomes soft.

44. Repeated electrolysis of 3% aqueous solution of NaOH.

45. Hard water passed through cation exchange resin which releases H^+ and then passed through anion exchanges resin which releases OH^-



OH^- neutralise the H^+ released in the cation exchange in (eq i)





46. Hard water when passed through resin containing $\text{R}-\text{COOH}$ groups it becomes free from Ca^{2+} ions.

49. Freezing point of D_2O and H_2O are 276.8 K and 273 K respectively.



50. Anhydrous CoCl (Blue) changes pink with water.

51. Ionic compounds are more soluble in soft water than heavy water. Soft water has high dielectric constant (78.39), while that of heavy water is 78.06. Due to higher polar character of soft water it is an excellent solvent for ionic compounds. Distillation of ionic compounds takes place because of ion-dipole interactions. Solubility of covalent compounds is due to the formation of hydrogen bonds with water molecules.

52. Heavy water is composed of heavy hydrogen (Deuterium) and oxygen. The formula of heavy water is $^2\text{H}_2\text{O}$ (D_2O).




56. Gaseous H_2O_2 has dihedral angle equal to 111.5° .




HYDROGEN AND ITS COMPOUNDS


60. H_2O_2 slowly decomposes to give nascent oxygen, which decolourises the coloured substances




nascent oxygen



61. Fe^{2+} ion oxidises to Fe^{3+} ion by H_2O_2



62. PbS reacts with H_2O_2 to give white PbSO_4



Black

White

63. 10 V means 3.035% H_2O_2 , hence 20 V means 6.070% H_2O_2

64. 30 vol = $3.035 \times 3 = 9.105\%$ H_2O_2

100 ml sol contain 9.105 gms H_2O_2

Hence 1000 ml solution contain 91.05 gms H_2O_2

Strength of $\text{H}_2\text{O}_2 = 91.05 \text{ gms/lit}$, basicity of $\text{H}_2\text{O}_2 = 2$




$$\text{Equivalent wt of } \text{H}_2\text{O}_2 = \left(\frac{34}{2} \right) = 17$$

$$\text{Normality} = \frac{\text{Strength}}{\text{Eq. wt.}} = \frac{91.05}{17} = 5.3558 = 5.36 \quad \text{Ans.}$$



66. H_2O_2 acts as reducing agent in acidic medium





67. H_2O_2 can not be dried over conc. H_2SO_4 because it oxidises by H_2SO_4




68. Hydrogen peroxide can be used as an oxidant, reductant and an acid. It oxidise Fe^{2+} into Fe^{3+} slowly in acidic medium but in basic medium it oxides very fast.



Hydrogen peroxide is also acidic in nature.



69. On industrial scale H_2O_2 is prepared by auto oxidation of 2-ethylanthraquinol.



2-ethyl anthru quinone

70. H_2O_2 can be used as antiseptic, bleaching agent and propellant.

EXERCISE - 2

Part # I : Multiple Choice

- (a) Hydrogen has both oxidising and reducing property while halogen have oxidising property.

(b) Hydrogen has both the + ve and - ve nature while halogen have - ve nature only.
- Metals react with hydrogen to form hydrides and thus oxidised

$$\text{Ca} + \text{H}_2 \longrightarrow \text{CaH}_2$$

$$2\text{Li} + \text{H}_2 \longrightarrow 2\text{LiH}$$
- Hydrides of 3rd gp and 17th gp are good acids. Also HN_3 is a good hydride.

$$\text{B}_2\text{H}_6 + 2\text{NH}_3 \longrightarrow 2[\text{BH}_3 \cdot \text{NH}_3]$$
1. $\text{Zn}_{(\text{s})} + \text{H}_2\text{SO}_{4(\text{l})} \longrightarrow \text{ZnSO}_{4(\text{aq})} + \text{H}_{2(\text{g})}$.

2. $\text{Zn}_{(\text{s})} + 2\text{NaOH} \longrightarrow \text{Na}_2\text{ZnO}_{2(\text{aq})} + \text{H}_{2(\text{g})}$

3. $\text{Cu}_{(\text{s})} + 2\text{H}_2\text{SO}_{4(\text{conc.})} \longrightarrow \text{CuSO}_{4(\text{aq})} + 2\text{H}_2\text{O}_{(\text{l})} + \text{SO}_{2(\text{g})}$

4. $2\text{F}_{2(\text{g})} + 2\text{H}_2\text{O}_{(\text{l})} \longrightarrow 4\text{HF} + \text{O}_2$
- (A) $2\text{MnO}_4^- + 6\text{H}^+ + 5\text{H}_2\text{O} \longrightarrow 2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 5\text{O}_2$
(reduction in acidic medium)

$$2\text{MnO}_4 + 3\text{H}_2\text{O}_2 \longrightarrow 2\text{MnO}_2 + 3\text{O}_2 + 2\text{H}_2\text{O} + 2\text{OH}^-$$

(reduction in basic medium)

(B) $2\text{Fe}^{2+}_{(\text{aq})} + 2\text{H}^+_{(\text{aq})} + \text{H}_2\text{O}_2 \longrightarrow 2\text{Fe}^{3+}_{(\text{aq})} + 2\text{H}_2\text{O}_{(\text{l})}$
(oxidation in acidic medium)

$$2\text{Fe}^{2+} + \text{H}_2\text{O}_2 \longrightarrow 2\text{Fe}^{3+} + 2\text{OH}^-$$



(C) $\text{Mn}^{2+} + \text{H}_2\text{O}_2 \longrightarrow \text{Mn}^{4+} + 2\text{OH}^-$
(oxidation in basic medium)

(D) $2\text{H}_2\text{O}_2 + 2\text{KI} \longrightarrow 2\text{KOH} + \text{I}_2 + 2\text{H}_2\text{O}$
(oxidation in acidic medium)


$$\text{I}_2 + \text{H}_2\text{O} + 2\text{OH}^- \longrightarrow 2\text{I}^- + 2\text{H}_2\text{O} + \text{O}_2$$

(reduction in basic medium)




14. A solution of $K_2Cr_2O_7$ in H_2SO_4 is oxidised to blue chromic acid by H_2O_2 and dissolve in ether to give blue coloured solution.



15. Hydrogen is non-metal while alkali metals have metallic character.
16. Tritium is radioactive.
17. Lattice energy of Al_2O_3 is very high so it require more energy to reduce, and giving high energy to hydrogen is explosive.
18. 5th gp elements V, Nb, Ta forms interstitial hydrides with hydrogen 7, 8, 9 gps of periodic table do not form hydrides but they absorb hydrogen, and it is called absorption. This inability of 7, 8, 9 gps of periodic table is referred to as hydride gap of d-block.
19. H_2O is an electron rich hydride it contain unbonded two lone pairs of electrons.
20. $10V = 3.035\%$
 \rightarrow 100 ml sol of H_2O_2 contain 3.035 gm
 \therefore 1000 ml sol of H_2O_2 contain $3.035 \times 10 = 30.35$ gm/lit.
21. H_2O_2 can not be concentrated by simple distillation because it decompose at its boiling point. It is concentrated under reduced pressure.
22. Aqueous solution of hydrogen peroxide is weakly acidic
 $H_2O_{2(aq)} \rightleftharpoons H^{\oplus} + HO_2^{\ominus} \quad K_a = 1.5 \times 10^{-12}$ at $25^\circ C$
 30% solution of H_2O_2 has $pH = 4$.
23. Heavy water is used as moderator in nuclear reactors.
24. Lime water $[Ca(OH)_2]$ is used to remove the temporary hardness of water. $Ca(OH)_2$ reacts with bicarbonates of Ca and Mg to form insoluble carbonate.
 $Ca(HCO_3)_{2(aq)} + Ca(OH)_2 \longrightarrow 2CaCO_3 \downarrow + 2H_2O$
 $Mg(HCO_3)_2 + Ca(OH)_2 \longrightarrow CaCO_3 \downarrow + MgCO_3 \downarrow + 2H_2O$
26. Due to hydrogen bonding interactions water molecules are associated together so water has high density than ice.




32. Water gas is the mixture of $[CO + H_2]$




34. Copper does not reacts with dil HCl.

36. Oxidation No. of Pb in PbO_2 is +4 while that is in PbO is +2 hence PbO_2 reduced to PbO by H_2O_2




40. Due to dipole-dipole interaction ice has cage like structure with a number of vacant spaces in the crystal lattice so the density of ice is lower than water.



$$\therefore 11.2 \text{ Vol } H_2O_2 = \frac{3.035 \times 11.2}{10} = 33.99 \text{ g,}$$


$$\text{molarity} = \frac{\text{strength}}{\text{mole wt}} = \frac{33.99}{34} = 1 \text{ M}$$

44. Ortho and para hydrogens are isomers of hydrogen.
 46. Both $-OH$ bonds in H_2O_2 do not lie in the same plane.



48. Water gas contains CO and H_2 in the ratio about. 4 : 5

49. High electron density on small N atom repel lone pair, making it more available for protonation. From P to Bi size of atom increases and so lone pair becomes less available for protonation.



51. $M_{\text{MgSO}_4} = 120$, $120 \text{ gms MgSO}_4 = 100 \text{ gms CaCO}_3$


$$0.24 \text{ gms MgSO}_4 = \frac{100}{120} \times .24$$

$$= \frac{2}{10} = .2 \text{ gms}$$

Again $\rightarrow 10^3 \text{ parts water} = .2 \text{ gm MgSO}_4$



$$\therefore 10^6 = \frac{.2}{10^3} \times 10^6 = .2 \times 1000$$

$$= 200 \text{ ppm.}$$




53. Enthalpy of vaporisation

PH_3	<	AsH_3	<	NH_3
b.b(k)	185.5	210.6		238.5



55. Silver does not react with dil or conc HCl.



$$.01 \text{ mole } 34 \times 4 \quad 303 \text{ gms}$$

$$2.39 \text{ gms } 136 \text{ gms}$$

$$2.39 \text{ gms PbS require } 136 \text{ gms of } \text{H}_2\text{O}_2$$


$$\text{We know } 10 \text{ V } \text{H}_2\text{O}_2 = 3.035 \text{ gms of } \text{H}_2\text{O}_2$$

$$\text{Hence } 3.035 \text{ gms } \text{H}_2\text{O}_2 = 1 \text{ ml of } 10 \text{ vol}$$

$$136 \text{ gms } \text{H}_2\text{O}_2 = \frac{1}{3.035} \times 136 \text{ ml} = 44.8 \text{ ml. Ans.}$$



60. S_1 : Tritium liberate β -radiations



61. S_1 : $\text{CaH}_2(\text{melt}) \rightarrow \text{Ca}^{2+}(\text{melt}) + 2\text{H}^-(\text{melt})$




$$\text{S}_3: \text{D}_2\text{O} = 41.61 \text{ kJ mol}^{-1}$$

$$\text{H}_2\text{O} = 40.66 \text{ kJ mol}^{-1}$$

62. S_3 : Form soluble complex which do not cause hindrance in the formation of lather.




Part # II : Assertion & Reason

1. NaH is crystalline hydride, it liberate H_2 at anode when electric current is passed through molten NaH
at anode $2\text{H}^- \rightarrow \text{H}_2 + 2\text{e}^-$





3. D_2 - $443.35 \text{ kJ mol}^{-1}$ and H_2 - $435.88 \text{ kJ mol}^{-1}$.

4. $\dot{\text{N}}\text{H}$ is electron rich hydride. NH_3 contain one lone pair of electron on nitrogen.




Complex salts

9. The bleaching action of H_2O_2 is due to oxidation




12. Demineralised water is free from Ca^{2+} and Mg^{2+} ions. Permutit exchange sodium ions with Ca^{2+} and Mg^{2+} ions, to give soft water.



EXERCISE - 3


Part # I : Matrix Match Type




Part # II : Comprehension

Comprehension # 1 :

Sol (1 to 2)



(A) (B)



(B) (A) (A) (C)

3. Interstitial hydrides are non-stoichiometric hydrides and thus deficient in hydrogen. Transition and innertransition elements at elevated temp. absorb hydrogen into the interstices of their lattices to yield metal like hydrides.




EXERCISE - 4

Part # I : AIEEE/JEE-MAIN

1. $[\text{CO} + \text{H}_2]_{(\text{g})} + \text{H}_2\text{O}_{(\text{g})} \xrightarrow[\text{Catalyst}]{673\text{K}} \text{CO}_{2(\text{g})} + 2\text{H}_{2(\text{g})}$,
 $\text{CO}_2 + 2\text{NaOH} \rightarrow \text{Na}_2\text{CO}_3 + \text{H}_2\text{O}$
2. A reducing agent loses electrons during redox reaction
Hence (b, d) is correct.
3. hydrogen bond is dipole-dipole interaction
4. It acts as oxidizing as well as reducing agent.
5. Intra molecular H bond is not present
6. Nitrate

Part # II : IIT-JEE ADVANCED

1. Ice is less dense than water due to open crystal structure because of H-bonding.
The basicity of 1° amines is more than 3° amines as after they donate lone pair to H^+ , they can form H-bonding with H_2O molecules and get easily stabilized by solvation. Where as in tertiary amines, the stabilization by solvation is very less.
The dimerisation of acetic acid in benzene is due to H-bonding.



Hence ans is : A, B, D

2. $\text{Zn} + 2\text{NaOH} \rightarrow \text{Na}_2\text{ZnO}_2 + \text{H}_2$

MOCK TEST

1. In electronegativity of hydrogen is in between metals and non metals so it behaves as both electro +ve and electro - ve. It can lose electron to form H^+ ion and it can also gain electron to form H^- ion.
2. (A) $\text{Cu} + \text{HCl} \rightarrow$ No reaction
(B) $3\text{Fe} + 4\text{H}_2\text{O} \xrightarrow[\text{magnetic oxide}]{\text{steam}} \text{Fe}_3\text{O}_4 + 4\text{H}_2 \uparrow$
(C) $\text{Mg} + 2\text{H}_2\text{O} \rightarrow \text{Mg}(\text{OH})_2 + \text{H}_2$
(D) $\text{Na} + \text{C}_2\text{H}_5\text{OH} \rightarrow \text{C}_2\text{H}_5\text{ONa} + \text{H}$
4. Ionisation energy of hydrogen is (1312 kJ mol^{-1}) which is too high than that of alkali metals and low than that of halogens.
5. Hydrogen accepts electron to form anion and get inert gas configuration like halogens.
 $\text{H} + \text{e}^- \rightarrow \text{H}^-$ (hydride ion)
 $\text{X} + \text{e}^- \rightarrow \text{X}^-$
halogen halide ion

6. $\text{TiH}_{1.5-1.8}$ is a interstitial hydride.
7. Hydrogen does not combine with helium.
8. $\text{H}^-_{(\text{aq})} + \text{H}-\text{OH}_{(\text{l})} \rightarrow \text{H}_{2(\text{g})} + \text{OH}^-_{(\text{aq})}$
 $\text{CaH}_2 + 2\text{H}_2\text{O} \rightarrow 2\text{H}_2 + \text{Ca}(\text{OH})_2$
9. Cation exchange resin exchanges Ca^{2+} , Mg^{2+} ions from water, and water becomes soft.
10. Hydrated silicates of Al and Na is called permutit $\text{Na}_2\text{Al}_2\text{Si}_2\text{O}_8 \cdot x\text{H}_2\text{O}$
11. H_2O_2 when oxidised in acidic or basic medium it produces O_2
 $2\text{MnO}_4^- + 6\text{H}^+ + 5\text{H}_2\text{O}_2 \rightarrow 2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 5\text{O}_2$ (acidic medium)
 $2\text{MnO}_4^- + 3\text{H}_2\text{O}_2 \rightarrow 2\text{MnO}_2 + 3\text{O}_2 + 2\text{H}_2\text{O}$ (basic medium)
12. Gaseous H_2O_2 has dihedral angle equal to 111.5° .
13. Fe^{2+} ion oxidises to Fe^{3+} ion by H_2O_2
 $2\text{Fe}^{2+} + \text{H}_2\text{O}_2 + 2\text{H}^+ \rightarrow 2\text{Fe}^{3+} + 2\text{H}_2\text{O}$
15. H_2O_2 acts as reducing agent in acidic medium
 $\text{H}_2\text{O}_2 \rightarrow 2\text{H}^+ + \text{O}_2 + 2\text{e}^-$
16. H_2O_2 can be used as antiseptic, bleaching agent and propellant.
20. NaH is crystalline hydride, it liberate H_2 at anode when electric current is passed through molten NaH
at anode $2\text{H}^- \rightarrow \text{H}_2 + 2\text{e}^-$
at cathode $2\text{Na}^+ + 2\text{e}^- \rightarrow 2\text{Na}$
22. $2\text{CaSO}_4 + \text{Na}_2[\text{Na}_4(\text{PO}_3)_6] \rightarrow \text{Na}_2[\text{Ca}_2(\text{PO}_3)_6] + 2\text{Na}_2\text{SO}_4$
 $2\text{MgSO}_4 + \text{Na}_2[\text{Na}_4(\text{PO}_3)_6] \rightarrow [\text{Mg}_2(\text{PO}_3)_6] + 2\text{Na}_2\text{SO}_4$
Complex salts
27. $\text{H}_2(\text{g}) + \text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g}) \xrightarrow[673 \text{ K}]{\text{FeCrO}_4} 2\text{H}_2(\text{g}) + \text{CO}_2(\text{g})$
28. (i) $\text{CaNCN} + 3\text{H}_2\text{O} \rightarrow \text{CaCO}_3 + 2\text{NH}_3$
(ii) $\text{Al}_4\text{C}_3 + 12\text{H}_2\text{O} \rightarrow 4\text{Al}(\text{OH})_3 + 3\text{CH}_4$
(iii) $\text{Cr}_2\text{O}_7^{2-} + 2\text{H}^+ + 4\text{H}_2\text{O}_2 \rightarrow 2\text{CrO}_5 + 5\text{H}_2\text{O}$
29. (i) Storage of H_2
(ii) Catalytic agents for hydrogenation.
30. (i) High enthalpy of combustion
(ii) No pollutant as product of combustion.