## Exercise-1

> Marked questions are recommended for Revision.

#### SUBJECTIVE QUESTIONS

#### Section (A) : Sine rule, Cosine rule, Napier's Analogy, Projection rule

A-1. In a  $\triangle ABC$ , prove that :

- $a \sin (B C) + b \sin (C A) + c \sin (A B) = 0$ (i)
- $\frac{a^2 \sin(B-C)}{\sin A} + \frac{b^2 \sin(C-A)}{\sin B} + \frac{c^2 \sin(A-B)}{\sin C} = 0$ (ii)
- $2(bc \cos A + ca \cos B + ab \cos C) = a^2 + b^2 + c^2$ (iii)

$$(iv)$$
  $(a - b)^2 \cos^2 \frac{b}{2} + (a + b)^2 \sin^2 \frac{b}{2} = c^2$ 

(v) 
$$b^2 \sin 2C + c^2 \sin 2B = 2bc \sin A$$

(vi) 
$$\frac{\sin B}{\sin C} = \frac{c - a \cos B}{b - a \cos C}$$

- Find the real value of x such that  $x^2 + 2x$ , 2x + 3 and  $x^2 + 3x + 8$  are lengths of the sides of a triangle. A-2.
- The angles of a  $\triangle ABC$  are in A.P. (order being A, B, C) and it is being given that b : c =  $\sqrt{3}$  :  $\sqrt{2}$ , then A-3. find  $\angle A$ .

**A-4.** If 
$$\cos A + \cos B = 4 \sin^2 \left(\frac{C}{2}\right)$$
, prove that sides a, c, b of the triangle ABC are in A.P.

If in a  $\triangle ABC$ ,  $\frac{\sin A}{\sin C} = \frac{\sin(A - B)}{\sin(B - C)}$ , then prove that  $a^2$ ,  $b^2$ ,  $c^2$  are in A.P. A-5.

In a triangle ABC, prove that for any angle  $\theta$ , b cos (A –  $\theta$ ) + a cos (B +  $\theta$ ) = c cos  $\theta$ . A-6.

A-7. With usual notations, if in a  $\triangle$  ABC,  $\frac{b+c}{11} = \frac{c+a}{12} = \frac{a+b}{13}$ , then prove that  $\frac{\cos A}{7} = \frac{\cos B}{19} = \frac{\cos C}{25}$ .

- Let a, b and c be the sides of a  $\triangle ABC$ . If  $a^2$ ,  $b^2$  and  $c^2$  are the roots of the equation A-8.  $x^3 - Px^2 + Qx - R = 0$ , where P, Q & R are constants, then find the value of  $\frac{\cos A}{2} + \frac{\cos B}{2} + \frac{\cos C}{2}$  in terms of P, Q and R.
- A-9. If in a triangle ABC, the altitude AM be the bisector of  $\angle BAD$ , where D is the mid point of side BC, then prove that  $(b^2 - c^2) = a^2/2$ .
- If in a triangle ABC,  $\angle C = 60^\circ$ , then prove that  $\frac{1}{a+c} + \frac{1}{b+c} = \frac{3}{a+b+c}$ A-10.
- A-11. A In a triangle ABC,  $\angle C = 60^{\circ}$  and  $\angle A = 75^{\circ}$ . If D is a point on AC such that the area of the  $\triangle ABD$  is  $\sqrt{3}$  times the area of the  $\triangle$ BCD, find the  $\angle$ ABD.
- A-12. In a scalene triangle ABC, D is a point on the side AB such that  $CD^2 = AD$ . DB, if sinA. sinB =  $sin^2 \frac{C}{2}$ then prove that CD is internal bisector of  $\angle C$ .
- A-13. In triangle ABC, D is on AC such that AD = BC, BD = DC,  $\angle DBC = 2x$ , and  $\angle BAD = 3x$ , all angles are in degrees, then find the value of x.

#### Section (B) Trigonometric ratios of Half Angles, Area of triangle and circumradius

- **B-1.** In a  $\triangle$ ABC, prove that
  - (i)  $2\left[a\sin^{2}\frac{C}{2} + c\sin^{2}\frac{A}{2}\right] = c + a b.$ (ii)  $\frac{\cos^{2}\frac{A}{2}}{a} + \frac{\cos^{2}\frac{B}{2}}{b} + \frac{\cos^{2}\frac{C}{2}}{c} = \frac{s^{2}}{abc}$
  - (iii)  $4\left(bc.\cos^2\frac{A}{2} + ca.\cos^2\frac{B}{2} + ab.\cos^2\frac{C}{2}\right) = (a + b + c)^2$
  - (iv) (b-c)  $\cot \frac{A}{2} + (c-a) \cot \frac{B}{2} + (a-b) \cot \frac{C}{2} = 0$
  - (v)  $4\Delta (\cot A + \cot B + \cot C) = a^2 + b^2 + c^2$
  - (vi)  $\left(\frac{2abc}{a+b+c}\right) \cdot \cos\frac{A}{2} \cdot \cos\frac{B}{2} \cdot \cos\frac{C}{2} = \Delta$
- **B-2.** If the sides a, b, c of a triangle are in A.P., then find the value of  $\tan \frac{A}{2} + \tan \frac{C}{2}$  in terms of  $\cot(B/2)$ .
- **B-3.** If in a  $\triangle$  ABC, a = 6, b = 3 and cos(A B) = 4/5, then find its area.
- **B-4.** If in a triangle ABC,  $\angle A = 30^{\circ}$  and the area of triangle is  $\frac{\sqrt{3} a^2}{4}$ , then prove that either B = 4C or C = 4B.

#### Section (C) Inradius and Exradius

- **C-1.** In any  $\triangle ABC$ , prove that
  - (i)  $\operatorname{Rr}(\sin A + \sin B + \sin C) = \Delta$  (ii)  $\operatorname{a} \cos B \cos C + \operatorname{b} \cos C \cos A + \operatorname{c} \cos A \cos B = \frac{\Delta}{R}$
  - (iii)  $\frac{1}{ab} + \frac{1}{bc} + \frac{1}{ca} = \frac{1}{2Rr}$ . (iv)  $\cos \cos^2 \frac{A}{2} + \cos^2 \frac{B}{2} + \cos^2 \frac{C}{2} = 2 + \frac{r}{2R}$

(v)  $a \cot A + b \cot B + c \cot C = 2(R + r)$ 

**C-2.** In any  $\triangle$ ABC, prove that (i) r. r<sub>1</sub> .r<sub>2</sub> .r<sub>3</sub> =  $\Delta^2$ 

- (ii) **a**  $r_1 + r_2 r_3 + r = 4R \cos C.$  (iii)  $\frac{1}{r^2} + \frac{1}{r_1^2} + \frac{1}{r_2^2} + \frac{1}{r_3^2} = \frac{a^2 + b^2 + c^2}{\Delta^2}$ (iv)  $\left(\frac{1}{r} + \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3}\right)^2 = \frac{4}{r} \left(\frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3}\right)$  (v) **a**  $\frac{bc - r_2 r_3}{r_1} = \frac{ca - r_3 r_1}{r_2} = \frac{ab - r_1 r_2}{r_3} = r$
- **C-3.** Show that the radii of the three escribed circles of a triangle are roots of the equation  $x^3 x^2(4R + r) + x s^2 r s^2 = 0$ .
- **C-4.** The radii r<sub>1</sub>, r<sub>2</sub>, r<sub>3</sub> of escribed circles of a triangle ABC are in harmonic progression. If its area is 24 sq. cm and its perimeter is 24 cm, find the lengths of its sides.
- **C-5.** If the area of a triangle is 100 sq.cm,  $r_1 = 10$  cm and  $r_2 = 50$  cm, then find the value of (b a).

#### Section (D) Miscellaneous

**D-1.** If  $\alpha$ ,  $\beta$ ,  $\gamma$  are the respective altitudes of a triangle ABC, prove that

(i)	1 _	1	1	$\cot A + \cot B + \cot C$	(ii)	1 1 1 _	2ab	$\cos^2 C$
	$\frac{1}{\alpha^2}$ + $\frac{1}{\beta^2}$ + $\frac{1}{\gamma^2}$	$\gamma^2$	Δ	(11)	$\frac{\alpha}{\alpha} + \frac{\beta}{\beta} - \frac{\gamma}{\gamma} =$	$(a + b + c) \Delta$	2 2	

- **D-2.** If in an acute angled ∆ABC, line joining the circumcentre and orthocentre is parallel to side AC, then find the value of tan A.tan C.
- **D-3.** A regular hexagon & a regular dodecagon are inscribed in the same circle. If the side of the dodecagon is  $(\sqrt{3} 1)$ , if the side of the hexagon is  $\sqrt[4]{k}$ , then find value of k.
- **D-4.** If D is the mid point of CA in triangle ABC and  $\Delta$  is the area of triangle, then show that  $\tan(\angle ADB) = \frac{4\Delta}{a^2 c^2}$ .

## Exercise-2

 $\boldsymbol{\varkappa}$  Marked questions are recommended for Revision.

### **PART-I (OBJECTIVE QUESTIONS)**

#### Section (A) : Sine rule, Cosine rule, Napier's Analogy, Projection rule

A-1.	In a ∆ABC, A : B : C = 3 (A) 2b	3 : 5 : 4. Then a + b + c · (B) 2c	√2 is equal to (C) 3b	(D) 3a			
A-2*.	In a triangle ABC, the	altitude from A is not les	s than BC and the altitu	de from B is not less than AC.			
	(A) right angled	(B) isosceles	(C) obtuse angled	(D) equilateral			
A-3.	If in a $\triangle$ ABC, $\frac{\cos A}{a} = -$	$\frac{\cos B}{b} = \frac{\cos C}{c}$ , then the	$\frac{dsB}{ds} = \frac{cosC}{c}$ , then the triangle is :				
	(A) right angled	(B) isosceles	(C) equilateral	(D) obtuse angled			
A-4.	In a $\triangle ABC = \frac{bc \sin^2}{\cos A + \cos^2}$	$\frac{A}{B\cos C}$ is equal to					
	(A) b <sup>2</sup> + c <sup>2</sup>	(B) bc	(C) a <sup>2</sup>	(D) a² + bc			
A-5æ	Given a triangle ∆ABC	such that sin <sup>2</sup> A + sin <sup>2</sup> C =	1001.sin <sup>2</sup> B. Then the va	lue of $\frac{2(\tan A + \tan C).\tan^2 B}{\tan A + \tan B + \tan C}$ is			
	(A) $\frac{1}{2000}$	(B) <u>1</u> 1000	(C) $\frac{1}{500}$	(D) $\frac{1}{250}$			
A-6.	If in a triangle ABC, (a - (A) $k < 0$	+ b + c) (b + c – a) = k. b (B) k > 6	c, then : (C) 0 < k < 4	(D) k > 4			
Δ-7	In a triangle ABC, a: b:	c = 4.5.6 Then 34 + B	equals to :	、 ,			
~	(A) 4C	(B) 2π	(C) $\pi$ – C	(D) π			

Solu	tion of Triangle									
A-8.æ	The distance between t	the middle point of BC ar	nd the foot of the perpend	dicular from A is :						
	(A) $\frac{-a^2 + b^2 + c^2}{2a}$	(B) $\frac{b^2 - c^2}{2 a}$	(C) $\frac{b^2 + c^2}{\sqrt{bc}}$	(D) $\frac{b^2 + c^2}{2 a}$						
A-9*.	If in a triangle ABC, cos (A) isosceles	s A cos B + sin A sin B si (B) right angled	n C = 1, then the triangle (C) equilateral	e is (D) None of these						
A-10.๖	Triangle ABC is right an	gle at A. The points P ar	nd Q are on hypotenuse	BC such that BP = PQ = QC.						
	If AP = 3 and AQ = 4, the second sec	hen length BC is equal to	) (C) 4.5	(D) 7						
	(A) 373	(B) 5 <del>4</del> 5	(C) 443							
A-11.	In $\triangle ABC$ , bc = 2b <sup>2</sup> cosA + 2c <sup>2</sup> cosA – 4bc cos <sup>2</sup> A, then $\triangle ABC$ is (A) isosceles but not necessarily equilaterial (B) equilateral (C) right angled but not neccessarily isosceles (D) right angled isosceles									
Sectio	on (B) Trigonometric	ratios of Half Angles	, Area of triangle and	circumradius						
B-1.	If in a triangle ABC, righ (A) a = 2, c = 3	nt angle at B, s – a = 3 a (B) a = 3, c = 4	nd s – c = 2, then (C) a = 4, c = 3	(D) a = 6, c = 8						
B-2.	If in a triangle ABC, b c	$\cos^2\frac{A}{2} + a\cos^2\frac{B}{2} = \frac{3}{2}c$	then a. c. b are :							
:	(A) in A.P.	2 2 2 (B) in G.P.	(C) in H.P.	(D) None						
B-3.æ	If H is the orthocentre of a triangle ABC, then the radii of the circle circumscribing the triangles BHC,									
	(A) R, R, R	(B) √2R, √2R, √2R	(C) 2R, 2R, 2R	(D) $\frac{R}{2}$ , $\frac{R}{2}$ , $\frac{R}{2}$						
R-4	In a $\wedge$ ABC if b + c = 3a	a then $\cot \frac{B}{H} \cdot \cot \frac{C}{H}$ has	the value equal to:							
D 4.	(A) 4	(B) 3	(C) 2	(D) 1						
	(,	(_) 0	(°) -							
B-5.	In a $\triangle ABC$ , A = $\frac{2\pi}{3}$ , b -	$-c = 3 \sqrt{3} cm and area$	$(\Delta ABC) = \frac{9\sqrt{3}}{2} \text{ cm}^2$ . The	en 'a' is						
	(A) 6 √3 cm	(B) 9 cm	(C) 18 cm	(D) 7 cm						
B-6.*	The diagonals of a par	rallelogram are inclined	to each other at an ang	le of 45°, while its sides a and						
	b(a > b) are inclined to	each other at an angle o	f 30º, then the value of	a is						
	(A) 2cos36º	(B) $\sqrt{\frac{3+\sqrt{5}}{4}}$	(C) $\frac{3+\sqrt{5}}{4}$	(D) $\frac{\sqrt{5}+1}{2}$						
B-7.	If in a $\triangle ABC$ , $\triangle = a^2 - (b A) (A) (A) (A) (A) (A) (A) (A) (A) (A) $	o – c)², then tan A is equ (B) 8/15	al to (C) 8/17	(D) 1/2						
B-8*.	If in a $\triangle ABC$ , a = 5, b =	4 and cos (A – B) = $\frac{31}{32}$	, then							
	(A) c = 6		(B) $\sin A = \left(\frac{5\sqrt{7}}{16}\right)$							
	(C) area of $\triangle ABC = \frac{15\sqrt{4}}{4}$	7	(D) c = 8							

- **B-9.** If R denotes circumradius, then in  $\triangle ABC$ ,  $\frac{b^2 c^2}{2a R}$  is equal to
  - (A)  $\cos (B C)$  (B)  $\sin (B C)$  (C)  $\cos B \cos C$  (D)  $\sin(B + C)$

B-10\*. Which of the following holds good for any triangle ABC?

(A) 
$$\frac{\cos A}{a} + \frac{\cos B}{b} + \frac{\cos C}{c} = \frac{a^2 + b^2 + c^2}{2abc}$$
(B) 
$$\frac{\sin A}{a} + \frac{\sin B}{b} + \frac{\sin C}{c} = \frac{3}{2R}$$
(C) 
$$\frac{\cos A}{a} = \frac{\cos B}{b} = \frac{\cos C}{c}$$
(D) 
$$\frac{\sin 2A}{a^2} = \frac{\sin 2B}{b^2} = \frac{\sin 2C}{c^2}$$

**B-11.** A triangle is inscribed in a circle. The vertices of the triangle divide the circle into three arcs of length 3, 4 and 5 units. Then area of the triangle is equal to:

(A) 
$$\frac{9\sqrt{3}(1+\sqrt{3})}{\pi^2}$$
 (B)  $\frac{9\sqrt{3}(\sqrt{3}-1)}{\pi^2}$  (C)  $\frac{9\sqrt{3}(1+\sqrt{3})}{2\pi^2}$  (D)  $\frac{9\sqrt{3}(\sqrt{3}-1)}{2\pi^2}$ 

B-12. In a ∆ABC, a = 1 and the perimeter is six times the arithmetic mean of the sines of the angles. Then measure of ∠ A is

- (A)  $\frac{\pi}{3}$  (B)  $\frac{\pi}{2}$  (C)  $\frac{\pi}{6}$  (D)  $\frac{\pi}{4}$
- B-13\*. Three equal circles of radius unity touches one another. Radius of the circle touching all the three circles is :

(A) 
$$\frac{2-\sqrt{3}}{\sqrt{3}}$$
 (B)  $\frac{\sqrt{3}-\sqrt{2}}{\sqrt{2}}$  (C)  $\frac{2+\sqrt{3}}{\sqrt{3}}$  (D)  $\frac{\sqrt{3}+\sqrt{2}}{\sqrt{2}}$ 

B-14. Triangle ABC is isosceles with AB = AC and BC = 65 cm. P is a point on BC such that the perpendicular distances from P to AB and AC are 24 cm and 36 cm, respectively. The area of triangle ABC (in sq. cm is)
(A) 1254
(B) 1950
(C) 2535
(D) 5070

#### Section (C) Inradius and Exradius

C-1	In a A ABC, the value of	$a\cos A + b\cos B + c\cos \theta$	C is equal to:	
•		a+b+c		
	(A) $\frac{r}{R}$	(B) $\frac{R}{2r}$	(C) $\frac{R}{r}$	(D) $\frac{2r}{R}$

C-2. In a triangle ABC, if a : b : c = 3 : 7 : 8, then R : r is equal to (A) 2 : 7 (B) 7 : 2 (C) 3 : 7 (D) 7 : 3

**C-3\*.** If  $r_1 = 2r_2 = 3r_3$ , then

(Δ)	a _	4	(B) <sup>a</sup> –	5	$(c)^{a}$ –	3	(ח)	а_	5
(~)	b _	5	(D) — – b	4	(C) — – c	5	(D)	с –	3

**C-4\*.** In a  $\triangle$ ABC, following relations hold good. In which case(s) the triangle is a right angled triangle? (A)  $r_2 + r_3 = r_1 - r$  (B)  $a^2 + b^2 + c^2 = 8 R^2$  (C)  $r_1 = s$  (D)  $2 R = r_1 - r$ 

**C-5.** The perimeter of a triangle ABC right angled at C is 70, and the inradius is 6, then |a - b| equals (A) 1 (B) 2 (C) 8 (D) 9

- **C-6.** In a triangle ABC, if  $\frac{a-b}{b-c} = \frac{s-a}{s-c}$ , then  $r_1$ ,  $r_2$ ,  $r_3$  are in: (A) A.P. (B) G.P. (C) H.P. (D) none of these
- **C-7.** If the incircle of the ∆ ABC touches its sides at L, M and N as shown in the figure and if x, y, z be the circumradii of the triangles MIN, NIL and LIM respectively, where I is the incentre, then the product xyz is equal to :



**C-.12\*.** With usual notations, in a  $\Delta$  ABC the value of  $\Pi$  (r<sub>1</sub> – r) can be simplified as:

(A) 
$$abc \prod tan \frac{A}{2}$$
 (B)  $4 r R^2$  (C)  $\frac{(abc)^2}{R(a+b+c)^2}$  (D)  $4 R r^2$ 

- **C-13. STATEMENT-1** : In a triangle ABC, the harmonic mean of the three exradii is three times the inradius. **STATEMENT-2** : In any triangle ABC,  $r_1 + r_2 + r_3 = 4R$ .
  - (A) STATEMENT-1 is true, STATEMENT-2 is true and STATEMENT-2 is correct explanation for STATEMENT-1
  - (B) STATEMENT-1 is true, STATEMENT-2 is true and STATEMENT-2 is not correct explanation for STATEMENT-1
  - (C) STATEMENT-1 is true, STATEMENT-2 is false
  - (D) STATEMENT-1 is false, STATEMENT-2 is true
  - (E) Both STATEMENTS are false

#### Section (D) Miscellaneous

D-1. ▲ If in a triangle ABC, the line joining the circumcentre and incentre is parallel to BC, then cos B + cos C is equal to : (A) 0 (B) 1 (C) 2 (D) 1/2

**D-2.** In a ∆ABC, if AB = 5 cm, BC = 13 cm and CA = 12 cm, then the distance of vertex 'A' from the side BC is (in cm)

- (A)  $\frac{25}{13}$  (B)  $\frac{60}{13}$  (C)  $\frac{65}{12}$  (D)  $\frac{144}{13}$
- **D-3.** If AD, BE and CF are the medians of a  $\triangle ABC$ , then  $(AD^2 + BE^2 + CF^2) : (BC^2 + CA^2 + AB^2)$  is equal to (A) 4 : 3 (B) 3 : 2 (C) 3 : 4 (D) 2 : 3

D-4\*. In a triangle ABC, with usual notations the length of the bisector of internal angle A is :

(A) $\frac{2bc \cos{\frac{A}{2}}}{b+c}$	(B) $\frac{2bc \sin \frac{A}{2}}{b+c}$	(C) $\frac{\text{abc} \cos \sec \frac{\pi}{2}}{2\text{R} (b+c)}$	(D) $\frac{2\Delta}{b+c} \cdot \csc \frac{A}{2}$
		()	

**D-5.** Let f, g, h be the lengths of the perpendiculars from the circumcentre of the  $\triangle$  ABC on the sides BC, CA and AB respectively. If  $\frac{a}{f} + \frac{b}{g} + \frac{c}{h} = \lambda \frac{a b c}{f g h}$ , then the value of ' $\lambda$ ' is: (A) 1/4 (B) 1/2 (C) 1 (D) 2

**D-6.** In an acute angled triangle ABC, AP is the altitude. Circle drawn with AP as its diameter cuts the sides AB and AC at D and E respectively, then length DE is equal to

(A) 
$$\frac{\Delta}{2R}$$
 (B)  $\frac{\Delta}{3R}$  (C)  $\frac{\Delta}{4R}$  (D)  $\frac{\Delta}{R}$ 

**D-7.**  $AA_1$ ,  $BB_1$  and  $CC_1$  are the medians of triangle ABC whose centroid is G. If points A,  $C_1$ , G and  $B_1$  are concyclic, then (A)  $2b^2 = a^2 + c^2$  (B)  $2c^2 = a^2 + b^2$  (C)  $2a^2 = b^2 + c^2$  (D)  $3a^2 = b^2 + c^2$ 

- **D-8.** If ' $\square$ ' is the length of median from the vertex A to the side BC of a  $\triangle$ ABC, then (A)  $4\square^2 = b^2 + 4ac \cos B(B) 4\square^2 = a^2 + 4bc \cos A(C) 4\square^2 = c^2 + 4ab \cos C(D) 4\square^2 = b^2 + 2c^2 - 2a^2$
- **D-9\*.** The product of the distances of the incentre from the angular points of a  $\triangle$  ABC is:

(A) 
$$4 R^2 r$$
 (B)  $4 Rr^2$  (C)  $\frac{(a b c) R}{s}$  (D)  $\frac{(a b c) r}{s}$ 

**D-10.** In a triangle ABC,  $B = 60^{\circ}$  and  $C = 45^{\circ}$ . Let D divides BC internally in the ratio 1 : 3,

then value of 
$$\frac{\sin \angle BAD}{\sin \angle CAD}$$
 is  
(A)  $\sqrt{\frac{2}{3}}$  (B)  $\frac{1}{\sqrt{3}}$  (C)  $\frac{1}{\sqrt{6}}$  (D)  $\frac{1}{3}$ 

**D-11\*.** In a triangle ABC, points D and E are taken on side BC such that BD = DE = EC. If angle ADE = angle AED =  $\theta$ , then:

(A) 
$$\tan \theta = 3 \tan B$$
 (B)  $3 \tan \theta = \tan C$  (C)  $\frac{6 \tan \theta}{\tan^2 \theta - 9} = \tan A$  (D) angle B = angle C

**D-12.** STATEMENT-1 : If R be the circumradius of a  $\triangle ABC$ , then circumradius of its excentral  $\triangle I_1I_2I_3$  is 2R.

**STATEMENT-2**: If circumradius of a triangle be R, then circumradius of its pedal triangle is  $\frac{\kappa}{2}$ 

- (A) STATEMENT-1 is true, STATEMENT-2 is true and STATEMENT-2 is correct explanation for STATEMENT-1
- (B) STATEMENT-1 is true, STATEMENT-2 is true and STATEMENT-2 is not correct explanation for STATEMENT-1
- (C) STATEMENT-1 is true, STATEMENT-2 is false
- (D) STATEMENT-1 is false, STATEMENT-2 is true
- (E) Both STATEMENTS are false

### **PART-II (COMPREHENSION)**

#### Comprehension # 1 (Q. No. 1 to 4)

The triangle DEF which is formed by joining the feet of the altitudes of triangle ABC is called the Pedal Triangle.

Answer The Following Questions :



- **1.** Angle of triangle DEF are (A)  $\pi$  – 2A,  $\pi$  – 2B and  $\pi$  – 2C (C)  $\pi$  – A,  $\pi$  – B and  $\pi$  – C
- 2\*. Sides of triangle DEF are
  (A) b cosA, a cosB, c cosC
  (C) R sin 2A, R sin 2B, R sin 2C

(B) a cosA, b cosB, c cosC

(B)  $\pi$  + 2A,  $\pi$  + 2B and  $\pi$  + 2C

(D)  $2\pi - A$ ,  $2\pi - B$  and  $2\pi - C$ 

- (D) a cotA, b cotB, c cotC
- 3. Circumraii of the triangle PBC, PCA and PAB are respectively
  (A) R, R, R
  (B) 2R, 2R, 2R
  (C) R/2, R/2, R/2
  (D) 3R, 3R, 3R
  4\*. Which of the following is/are correct
  - (A)  $\frac{\text{Perimeter}}{\text{Perimeter}} \text{ of } \Delta \text{DEF} = \frac{r}{R}$ (B) Area of  $\Delta \text{DEF} = 2 \Delta \cos A \cos B \cos C$ (C) Area of  $\Delta \text{AEF} = \Delta \cos^2 A$ (D) Circum-radius of  $\Delta \text{DEF} = \frac{R}{2}$

#### Comprehension # 2 (Q. 5 to 8)

The triangle formed by joining the three excentres  $I_1$ ,  $I_2$  and  $I_3$  of  $\Delta$  ABC is called the excentral or excentric triangle and in this case internal angle bisector of triangle ABC are the altitudes of triangles  $I_1I_2I_3$ 

5.24	Incentre I of $\triangle$ ABC is the (A) Circumcentre	ne of the excentral (B) Orthocentre	$\Delta I_1 I_2 I_3.$ (C) Centroid		(D) None of these
6.24	Angles of the $\Delta I_1 I_2 I_3$ and (A) $\frac{\pi}{2} - \frac{A}{2}$ , $\frac{\pi}{2} - \frac{B}{2}$ and (C) $\frac{\pi}{2} - A$ , $\frac{\pi}{2} - B$ and	$\frac{\pi}{2} - \frac{C}{2}$ $\frac{\pi}{2} - C$	(B) $\frac{\pi}{2} + \frac{A}{2}$ , $\frac{\pi}{2}$ (D) None of the	$\frac{1}{2} + \frac{B}{2}$ and we se	$\frac{\pi}{2} + \frac{C}{2}$

Solu	tion of Triangle			
7.2	Sides of the $\Delta I_1 I_2 I_3$ are			
	(A) $\operatorname{Rcos} \frac{A}{2}$ , $\operatorname{Rcos} \frac{B}{2}$ ar	nd Rcos $\frac{C}{2}$	(B) 4R $\cos\frac{A}{2}$ , 4R $\cos\frac{A}{2}$	$\frac{B}{2}$ and 4R cos $\frac{C}{2}$
	(C) $2R\cos\frac{A}{2}$ , $2R\cos\frac{B}{2}$	and $2R\cos{\frac{C}{2}}$	(D) None of these	
8.2	Value of $II_1^2 + I_2I_3^2 = II_2^2$ (A) $4R^2$	$I_{1}^{2} + I_{3}I_{1}^{2} = II_{3}^{2} + I_{1}I_{2}^{2} =$ (B) 16R <sup>2</sup>	(C) 32R <sup>2</sup>	(D) 64R <sup>2</sup>

## PART-III (MATCH THE COLUMN)

Column–II

**1.** Match the column

#### Column– I

2.

(A)	In a $\triangle ABC$ , $2B = A + C$ and $b^2 = ac$ .	(p)	8							
	Then the value of $\frac{a^2(a+b+c)}{3abc}$ is equal to									
(B)	In any right angled triangle ABC, the value of $\frac{a^2 + b^2 + c^2}{R^2}$	(q)	1							
	is always equal to (where R is the circumradius of $\Delta ABC$ )									
(C)	In a $\triangle ABC$ if a = 2, bc = 9, then the value of $2R\Delta$ is equal to	(r)	5							
(D)	In a $\triangle ABC$ , a = 5, b = 3 and c = 7, then the value of 3 cos C + 7 cos B is equal to	(s)	9							
Match the column										
Colum	n – I	Colum	n – II							
(A)	In a $\triangle ABC$ , a = 4, b = 3 and the medians $AA_1$ and $BB_1$ are mutually perpendicular, then square of area of the $\triangle ABC$ is equal to	(p)	27							
(B)æ	In any $\triangle ABC$ , minimum value of $\frac{r_1 r_2 r_3}{r^3}$ is equal to	(q)	7							
(C)	In a $\triangle ABC$ , a = 5, b = 4 and tan $\frac{C}{2} = \sqrt{\frac{7}{9}}$ , then side 'c' is equal to	(r)	6							
(D) کھ	In a $\triangle ABC$ , $2a^2 + 4b^2 + c^2 = 4ab + 2ac$ , then value of (8 cos B) is equal to	(s)	11							

## **Exercise-3**

> Marked Questions may have for Revision Questions.

### PART - I : JEE (ADVANCED) / IIT-JEE PROBLEMS (PREVIOUS YEARS)

#### \* Marked Questions may have more than one correct option.

- 1. Given an isosceles triangle, whose one angle is 120° and radius of its incircle is  $\sqrt{3}$  unit. Then the area of the triangle in sq. units is [IIT-JEE-2006, Main.,(3, -1)/184] (A) 7 + 12  $\sqrt{3}$  (B) 12 - 7  $\sqrt{3}$  (C) 12 + 7  $\sqrt{3}$  (D)  $4\pi$
- **2.\*** Internal bisector of  $\angle A$  of triangle ABC meets side BC at D. A line drawn through D perpendicular to AD intersects the side AC at E and the side AB at F. If a, b, c represent sides of  $\triangle ABC$ , then

[IIT-JEE-2006, Main.,(5, -1)/184](A) AE is HM of b and c (B) AD =  $\frac{2bc}{b+c} \cos \frac{A}{2}$ (C) EF =  $\frac{4bc}{b+c} \sin \frac{A}{2}$ (D) the triangle AEF is isosceles

**3.** Let ABC and ABC' be two non-congruent triangles with sides AB = 4,  $AC = AC' = 2\sqrt{2}$  and angle  $B = 30^{\circ}$ . Find the absolute value of the difference between the areas of these triangles.

#### [IIT-JEE 2009, Paper-2, (4, -1), 80]

4\*. In a triangle ABC with fixed base BC, the vertex A moves such that  $\cos B + \cos C = 4 \sin^2 \frac{A}{2}$ . If a, b and c denote the lengths of the sides of the triangle opposite to the angles A, B and C respectively, then [IIT-JEE 2009, Paper-1, (4, -1)/ 80] (A) b + c = 4a (B) b + c = 2a (C) locus of points A is an ellipse (D) locus of point A is a pair of straight lines

If the angle A, B and C of a triangle are in arithmetic progression and if a, b and c denote the lengths of the sides opposite to A, B and C respectively, then the value of the expression and if a sin 2C + c/a sin 2A is
 [IIT-JEE 2010, Paper-1, (3, -1), 84]

(A)  $\frac{1}{2}$  (B)  $\frac{\sqrt{3}}{2}$  (C) 1 (D)  $\sqrt{3}$ 

**6.** Let ABC be a triangle such that  $\angle ACB = \frac{\pi}{6}$  and let a, b and c denote the lengths of the sides opposite to A, B and C respectively. The value(s) of x for which  $a = x^2 + x + 1$ ,  $b = x^2 - 1$  and c = 2x + 1 is (are) (A)  $-(2 + \sqrt{3})$  (B)  $1 + \sqrt{3}$  (C)  $2 + \sqrt{3}$  (D)  $4\sqrt{3}$ 

#### [IIT-JEE 2010, Paper-1, (3, 0), 84]

**7.** Consider a triangle ABC and let a, b and c denote the lengths of the sides opposite to vertices A, B and C respectively. Suppose a = 6, b = 10 and the area of the triangle is  $15\sqrt{3}$ . If  $\angle ACB$  is obtuse and if r denotes the radius of the incircle of the triangle, then r<sup>2</sup> is equal to **[IIT-JEE 2010, Paper-2, (3, 0), 79]** 

Let PQR be a triangle of area  $\Delta$  with a = 2, b =  $\frac{7}{2}$  and c =  $\frac{5}{2}$ , where a, b and c are the lengths of the 8. sides of the triangle opposite to the angles at P, Q and R respectively. Then  $\frac{2 \sin P - \sin 2P}{2 \sin P + \sin 2P}$  equals [IIT-JEE 2012, Paper-2, (3, -1), 66] (C)  $\left(\frac{3}{4\Delta}\right)^2$  (D)  $\left(\frac{45}{4\Delta}\right)^2$ (B)  $\frac{45}{4\Lambda}$ (A)  $\frac{3}{4\Lambda}$ In a triangle PQR, P is the largest angle and  $\cos P = \frac{1}{3}$ . Further the incircle of the triangle touches the 9.\*> sides PQ, QR and RP at N, L and M respectively, such that the lengths of PN, QL and RM are consecutive even integers. Then possible length(s) of the side(s) of the triangle is (are) [JEE (Advanced) 2013, Paper-2, (3, -1)/60] (A) 16 (B) 18 (C) 24 (D) 22 10. In a triangle the sum of two sides is x and the product of the same two sides is y. If  $x^2 - c^2 = y$ , where c is the third side of the triangle, then the ratio of the in-radius to the circum-radius of the triangle is [JEE (Advanced) 2014, Paper-2, (3, -1)/60] (A)  $\frac{3y}{2x(x+c)}$  (B)  $\frac{3y}{2c(x+c)}$  (C)  $\frac{3y}{4x(x+c)}$  (D)  $\frac{3y}{4c(x+c)}$ In a triangle XYZ, let x, y, z be the lengths of sides opposite to the angles X, Y, Z, respectively, and 11\*. 2s = x + y + z. If  $\frac{s - x}{4} = \frac{s - y}{3} = \frac{s - z}{2}$  and area of incircle of the triangle XYZ is  $\frac{8\pi}{3}$ , then (A) area of the triangle XYZ is  $6\sqrt{6}$ [JEE (Advanced) 2016, Paper-1, (4, -2)/62] (B) the radius of circumcircle of the triangle XYZ is  $\frac{35}{\epsilon} \sqrt{6}$ (C)  $\sin \frac{X}{2} \sin \frac{Y}{2} \sin \frac{Z}{2} = \frac{4}{25}$ (D)  $\sin^2\left(\frac{X+Y}{2}\right) = \frac{3}{5}$ In a triangle PQR, let  $\angle$  PQR = 30° and the sides PQ and QR have lengths 10  $\sqrt{3}$  and 10, respectively. 12\*. Then, which of the following statement(s) is (are) TRUE? (A) ∠QPR = 45° [JEE(Advanced) 2018, Paper-1,(4, -2)/60] (B) The area of the triangle PQR is  $25\sqrt{3}$  and  $\angle QRP = 120^{\circ}$ (C) The radius of the incircle of the triangle PQR is  $10\sqrt{3}$  – 15 (D) The area of the circumcircle of the triangle PQR is  $100\pi$ 

## PART - II : JEE (MAIN) / AIEEE PROBLEMS (PREVIOUS YEARS)

1.	The sum of the radii of inscribed and circumscribed circles for an n sided regular polygon of side 'a', is : [AIEEE – 2003 (3, 0), 225]									
	(1) a $\cot\left(\frac{\pi}{n}\right)$	(2) $\frac{a}{2} \cot\left(\frac{\pi}{2n}\right)$	(3) a cot $\left(\frac{\pi}{2n}\right)$	(4) $\frac{a}{4} \cot\left(\frac{\pi}{2n}\right)$						
2.	If in a triangle ABC, a c	$\cos^2\left(\frac{C}{2}\right) + c\cos^2\left(\frac{A}{2}\right) =$	$=\frac{3b}{2}$ , then the sides a, b and c :							
	(1) are in A.P.	(2) are in G.P.	(3) are in H.P.	[AIEEE – 2003 (3, 0), 225] (4) satisfy a + b = c.						
3.2	In a triangle ABC, med	ians AD and BE are drav	vn. If AD = 4, $\angle$ DAB = $\frac{\pi}{6}$	and $\angle ABE = \frac{\pi}{3}$ , then the area						
	of the $\triangle ABC$ is :		-	[AIEEE – 2003 (3, 0), 225]						
	(1) $\frac{8}{3}$	(2) $\frac{16}{3}$	(3) $\frac{32}{3\sqrt{3}}$	(4) $\frac{64}{3}$ .						
4.	The sides of a triangle	are sin $\alpha$ , cos $\alpha$ and $\sqrt{1+\alpha}$	$\sin \alpha \cos \alpha$ for some 0 <	$\alpha < \frac{\pi}{2}$ . Then the greatest angle						
	of the triangle is : (1) 60º	(2) 90°	(3) 120º	[AIEEE – 2004 (3, 0), 225] (4) 150°						
5.	In a triangle ABC, let 2	$\angle C = \pi/2$ , if r is the inrad	lius and R is the circum	radius of the triangle ABC, then						
	2(r+R) equals : (1) c + a	(2) a + b + c	(3) a + b	[ <b>AIEEE - 2005 (3, 0), 225]</b> (4) b + c						
6.	If in a ∆ABC, the altitue are in : (1) HP (3) AP	des from the vertices A,I	<ul> <li>3,C on opposite sides are in H.P., then sinA, sinB, sinC [AIEEE - 2005 (3, 0), 225]</li> <li>(2) Arithemetico-Geometric Progression</li> <li>(4) GP</li> </ul>							
7.	For a regular polygon, statement among the for	let r and R be the radii blowing is	of the inscribed and the	e circumscribed circles. A false [AIEEE - 2010 (4, -1), 144]						
	(1) There is a regular p	polygon with $\frac{r}{R} = \frac{1}{\sqrt{2}}$ .	(2) There is a regular polygon with $\frac{r}{R} = \frac{2}{3}$ .							
	(3) There is a regular p	polygon with $\frac{r}{R} = \frac{\sqrt{3}}{2}$ .	(4) There is a regular p	polygon with $\frac{r}{R} = \frac{1}{2}$ .						
8.2	ABCD is a trapezium so then AB is equal to :	uch that AB and CD are	parallel and BC $\perp$ CD. If	∠ADB = θ , BC = p and CD = q, [AIEEE - 2013, (4, –1),120]						
	(1) $\frac{(p^2 + q^2)\sin\theta}{p\cos\theta + q\sin\theta}$	(2) $\frac{p^2 + q^2 \cos \theta}{p \cos \theta + q \sin \theta}$	$(3) \ \frac{p^2 + q^2}{p^2 \cos \theta + q^2 \sin \theta}$	(4) $\frac{(p^2 + q^2)\sin\theta}{(p\cos\theta + q\sin\theta)^2}$						
9.	With the usual notation	, in $\triangle ABC$ , if $\angle A + \angle B =$	120°, a = $\sqrt{3}$ + 1 and b =	= $\sqrt{3}$ –1, then the ratio $\angle A$ : $\angle B$ ,						
	is :	[JEE(N	Main) 2019, Online (10-0	)1-19),P-2 (4, – 1), 120]						
	(1) 9 : 7	(2) 7 : 1	(3) 3 : 1	(4) 5 : 3						
10.	In a triangle, the sum o y. If $x^2 - c^2 = y$ , wher triangle is	f lengths of two sides is a e c is the length of the [JEE(N	x and the product of the third side of the triangl <b>/ain) 2019, Online (11-(</b>	lengths of the same two sides is e, then the circumradius of the 01-19),P-1 (4, – 1), 120]						
	(1) $\frac{c}{\sqrt{3}}$	(2) $\frac{3}{2}y$	(3) $\frac{c}{3}$	(4) $\frac{y}{\sqrt{3}}$						

Solu	Solution of Triangle												
	Answers												
	EXERCISE - 1												
Secti	Section (A) :												
					_								
A-2.	x > 5	A-3.	75°	A-8.	$\frac{P}{2\sqrt{R}}$	A-11.	30°	A-13.	10°				
Section (B)													
B-2.	$\frac{2}{3}\cot{\frac{4}{3}}$	<u>3</u> 2	B-3.	9 sq. u	nit								
Section (C) :													
C-4.	6, 8, 10	0 cm	C-5.	8									
Section (D) :													
D-2.	3		D-3.	√2									
						EXERC	SISE - 2						
Secti	on (A)	:											
A-1.	(C)	A-2*.	(AB)	A-3.	(C)	A-4.	(C)	A-5.	(D)	A-6.	(C)	A-7.	(D)
A-8.	(B)	A-9.	(AB)	A-10.	(A)	A-11.	(A)						
Secti	on (B)	:											
B-1.	(B)	B-2.	(A)	B-3.	(A)	B-4.	(C)	B-5.	(B)	B-6.	(AD)	B-7.	(B)
B-8*.	(ABC)	B-9.	(B)	B-10*.	(AB)	B-11.	(A)	B-12.	(C)	B-13*.	(AC)	B-14.	(C)
Secti	on (C)	:											
C-1.	(A)	C-2.	(B)	C-3*.	(BD)	C-4*.	(ABCD)	) <b>C-5.</b>	(A)	C-6.	(A)	C-7.	(C)
C-8.	(B)	C-9.	(D)	C-10.	(D)	C-11*.	(AD)	C12*.	(ACD)	C-13.	(C)		

<u>Solu</u> Secti	Solution of Triangle Section (D)												
0000	011 (2)												
D-1.	(B)	D-2.	(B)	D-3.	(C)	D-4*.	(ACD)	D-5.	(A)	D-6.	(D)		
D-7.	(C)	D-8.	(B)	D-9*.	(BD)	D-10.	(C)	D-11*.	(ACD)	D-12.	(A)		
	PART-II												
1.	(A)	2*.	(BC)	3.	(A)	4*.	(ABCD	) 5.	(B)	6.	(A)	7.	(B)
8.	(B)												
	PART-III												
1.	$(A) \to$	(q),	$(B) \to$	(p),	$(C) \rightarrow$	(s),	$(D) \rightarrow$	(r)					
2.	$(A) \to$	(s),	$(B) \to$	(p),	$(C) \rightarrow$	(r),	$(D) \rightarrow$	(q)					
						EXER	CISE - :	3					
						PAF	RT - I						
1.	(C)	2.*	(ABCE	<b>D</b> )	3.	4	4*.	(BC)	5.	(D)	6.	(B)	<b>7.</b> 3
8.	(C)	9.*	(BD)	10.	(B)	11.	(ACD)		12.	(BCD)			
						PAR	RT - II						
1.	(2)	2.	(1)	3.	(3)	4.	(3)	5.	(3)	6.	(3)	7.	(2)
8.	(1)	9.	(2)	10.	(1)								

## Advance Level Problems (ALP)

A Marked questions are recommended for Revision.

- 1. In  $\triangle ABC$ , P is an interior point such that  $\angle PAB = 10^{\circ} \angle PBA = 20^{\circ}$ ,  $\angle PCA = 30^{\circ}$ ,  $\angle PAC = 40^{\circ}$  then prove that  $\triangle ABC$  is isosceles
- **2.** In a triangle ABC, if a tan A + b tan B = (a + b) tan  $\left(\frac{A+B}{2}\right)$ , prove that triangle is isosceles.
- **3.** In any triangle ABC, if  $2\Delta a b^2c = c^3$ , (where  $\Delta$  is is the area of triangle), then prove that  $\angle A$  is obtuse
- 4. If in a triangle ABC,  $\frac{\cos A + 2\cos C}{\cos A + 2\cos B} = \frac{\sin B}{\sin C}$  prove that the triangle ABC is either isosceles or right angled.
- **5.** In a  $\triangle$  ABC,  $\angle$  C = 60° and  $\angle$  A = 75°. If D is a point on AC such that the area of the  $\triangle$  BAD is  $\sqrt{3}$  times the area of the  $\triangle$  BCD, find the  $\angle$  ABD.
- 6. In a  $\triangle ABC$ , if a, b and c are in A.P., prove that  $\cos A.cot \frac{A}{2}$ ,  $\cos B.cot \frac{B}{2}$ , and  $\cos C.cot \frac{C}{2}$  are in A.P.
- 7. In a triangle ABC, prove that the area of the incircle is to the area of triangle itself is,  $\pi : \cot\left(\frac{A}{2}\right) \cdot \cot\left(\frac{B}{2}\right) \cdot \cot\left(\frac{C}{2}\right).$

8. In 
$$\triangle ABC$$
, prove that  $a^2 (s - a) + b^2 (s - b) + c^2 (s - c) = 4R\Delta \left(1 + 4\sin\frac{A}{2}\sin\frac{B}{2}\sin\frac{C}{2}\right)$ 

**9.** In any  $\triangle ABC$ , prove that

(i) 
$$(r_3 + r_1) (r_3 + r_2) \sin C = 2 r_3 \sqrt{r_2 r_3 + r_3 r_1 + r_1 r_2}$$

(ii) 
$$\frac{\tan \frac{A}{2}}{(a - b)(a - c)} + \frac{\tan \frac{B}{2}}{(b - a)(b - c)} + \frac{\tan \frac{C}{2}}{(c - a)(c - b)} = \frac{1}{\Delta}$$

(iii) 
$$(r+r_1) \tan \frac{B-C}{2} + (r+r_2) \tan \frac{C-A}{2} + (r+r_3) \tan \frac{A-B}{2} = 0$$

(iv)  $r^2 + r_1^2 + r_2^2 + r_3^2 = 16R^2 - a^2 - b^2 - c^2$ .

**10.** In an acute angled triangle ABC,  $r + r_1 = r_2 + r_3$  and  $\angle B > \frac{\pi}{3}$ , then prove that b + 3c < 3a < 3b + 3c

- **11.** If the inradius in a right angled triangle with integer sides is r. Prove that
  - (i) If r = 4, the greatest perimeter (in units) is 90
  - (ii) If r = 5, the greatest area (in sq. units) is 330

**12.** If 
$$\left(1 - \frac{r_1}{r_2}\right) \left(1 - \frac{r_1}{r_3}\right) = 2$$
, then prove that the triangle is right angled

**13.** DEF is the triangle formed by joning the points of contact of the incircle with the sides of the triangle ABC; prove that

(i) its sides are 
$$2r \cos \frac{A}{2}$$
,  $2r \cos \frac{B}{2}$  and  $2r \cos \frac{C}{2}$   
(ii) its angles are  $\frac{\pi}{2} - \frac{A}{2}$ ,  $\frac{\pi}{2} - \frac{B}{2}$  and  $\frac{\pi}{2} - \frac{C}{2}$ 

and

- (iii) its area is  $\frac{2\Delta^3}{(abc)s}$  , i.e.  $\frac{1}{2} \frac{r}{R} \Delta$ .
- **14.** Three circles, whose radii are a, b and c, touch one another externally and the tangents at their points of contact meet in a point, prove that the distance of this point from either of their points of contact is

$$\left(\frac{abc}{a+b+c}\right)^{\frac{1}{2}}.$$

- **15.** OA and OB are the equal sides of an isoscles triangle lying in the first quadrant making angles  $\theta$  and  $\phi$  respectively with x-axis. Show that the gradient of the bisector of acute angle AOB is cosec  $\beta$  cot  $\beta$  where  $\beta = \phi + \theta$ . (Where O is origin)
- **16.** The hypotenuse BC = a of a right-angled triangle ABC is divided into n equal segments where n is odd. The segment containing the midpoint of BC subtends angle  $\alpha$  at A. Also h is the altitude of the triangle

through A. Prove that  $\tan \alpha = \frac{4nh}{a(n^2-1)}$ 

# Answer Key (ALP)

**5.**  $\angle ABD = 30^{\circ}$