### **REDOX REACTION**

### 6.0 INTRODUCTION

Redox reactions show vital role in non renewable energy source. In cell reaction where oxidation and reduction both occurs simultaneously will have redox reaction for interconversion of energy.

## 6.1 Redox Reaction (Oxidation-Reduction) :

Many chemical reaction involve transfer of electrons from one chemical substance to another. These electron-transfer reaction are termed as oxidation-reduction or redox reaction.

Or

Those reaction which involve oxidation and reduction both simultaneously are known as oxidation reduction or redox reactions.

Or

Those reactions in which increase and decrease in oxidation number of same or different atoms occurs are known as redox reactions.

## 6.2 Oxidation State

Ex.

Oxidation state of an atom in a molecule or ion is the hypothetical or real charge present on an atom due to electronegativity difference.

Or

Oxidation state of an element in a compound represents the number of electrons lost or gained during its change from free state into that compound.

## Some important points concerning oxidation number :-

(1) Electronegativity values of no two elements are same-

P > H C > H S > C Cl > N

(2) Oxidation number of an element may be positive or negative.

(3) Oxidation number can be zero, whole number or a fractional value.

Ex.	Ni(CO) <sub>4</sub>	$\Rightarrow$	O.S of Ni = $0$
	N <sub>3</sub> H	$\Rightarrow$	O.S of N = $-1/3$
	HC1	$\Rightarrow$	O.S of $Cl = -1$
(4) O	xidation stat	e of same	e element can be diff

(4) Oxidation state of same element can be different in same or different compounds.

$H_2S$	$\Rightarrow$	O.S of $S = -2$
$H_2SO_3$	$\Rightarrow$	O.S of $S = +4$
$H_2SO_4$	$\Rightarrow$	O.S of $S = +6$

## 6.3 Some helping rules for calculating oxidation number :

## (A) In case of covalent bond –

(i) For hmoatomic molecule

A - A $\mathbf{A} = \mathbf{A}$  $A \equiv A$  $\downarrow$  $\downarrow$  $\downarrow$  $\downarrow$  $\downarrow$  $\downarrow$ 0 0 0 0 0 0 (ii) For heteroatomic molecule (EN of B > A) A - AA = A $A \equiv A$  $\downarrow$  $\downarrow$  $\downarrow$  $\downarrow$  $\downarrow$  $\downarrow$ +1 -1+2 -2+3 -3

(iii) The oxidation state of an element in its free state is zero. Example – Oxidation state of Na, Cu, I, Cl, O etc. are zero. (iv) Oxidation state of atoms present in homoatomic molecules is zero. Ex.  $H_2$ ,  $O_2$ ,  $N_2$ ,  $P_4$ ,  $S_8 = zero$ (v) Oxidation state of an element in any of its allotropic form is zero. Ex.  $C_{Diamond}$ ,  $C_{Graphite}$ ,  $S_{Monoclinic}$ ,  $S_{Rhombic} = 0$ (vi) Oxidation state of all the components of an alloy are 0 (vii) In complex compounds, oxidation state of some neutral molecules (ligands) is zero. Ex. CO, NO,  $NH_3$ ,  $H_2O$ . (viii) Oxidation state of fluorine in all its compounds is -1 (ix) Oxidation state of I A & II A group elements are +1 and +2 respectively. (x) Oxidation stee of hydrogen in most of its compounds is +1 except in metal hydrides (-1) Ex. NaH LiH  $CaH_2$ MgH<sub>2</sub>  $\downarrow \downarrow$  $\downarrow \downarrow$  $\downarrow \downarrow$  $\downarrow \downarrow$ +1 - 1+1 - 1+2 - 1+2 - 1(xi) Oxidation state of oxygen in most of its compounds is -2except in-(a) Peroxides  $(O_2^{-2}) \rightarrow Oxidation state (O) = -1$ Ex.  $H_2O_2$ ,  $BaO_2$ (b) Super oxides  $(O_2^{-1}) \rightarrow Oxidation state (O) = -1/2$ Ex.  $KO_2$  $\downarrow$ -1/3(d) OF<sub>2</sub> (Oxygen difluoride) Ex. F - O - F $\downarrow$ +2(e)  $O_2F_2$  (Dioxygen difluoride)  $\downarrow$ +1(xii) Oxidation state of monoatomic ions is equal to the charge present on the ion. Ex.  $Mg^{+2} \rightarrow Oxidation state = +2$ (xiii) The algebric sum of oxidation state of all the atoms present in a polytomic neutral molecule is 0. Ex.  $H_2SO_4$ If O.S. of S is x then 2(+1) + x + 4(-2) = 0x - 6 = 0=+6Ex.  $H_2SO_3$ If O.S. of S is x then 2(+1) + x + 3(-2) = 0x - 4 = 0= +4(xiv) The algebraic sum of oxidation state of all the atoms in a polyatomic complex ion is equal to the charge present on the ion.

Ex.  $\underline{S}O_4^{-2}$ 

Power by: VISIONet Info Solution Pvt. Ltd	
Website : www.edubull.com	

### Edubull

If O.S. of S is x then x + 4(-2) = -2x - 6 = 0=+6 $H\underline{C}O_3^-$ Ex. If O.S. of C is x then +1 + x + 3(-2) = -1x - 4 = 0= +4**(B)** In case of co-ordinate bond (EN of B > A) :  $A \rightarrow A$  $B \rightarrow B$  $A \rightarrow B$  $B \rightarrow A$  $\downarrow$  $\downarrow$  $\downarrow$  $\downarrow \downarrow$  $\downarrow$  $\downarrow$  $\downarrow$ +2 -2 +2 -2 +2 -2 (neglected) **(C)** In case of Ionic bond : Charge on cation = O.S. of cation Charge on anion = O.S. of anion Ex.  $NaCl \rightarrow$  $Na^+$ Cl + $\downarrow$  $\downarrow$ -1 +1 $Mg^{+2}$  $MgCl_2 \rightarrow$  $2Cl^{-}$ +  $\downarrow$  $\downarrow$ +2-1 Illustrations Illustration 1. Oxidation number of cobalt in [Co(NH<sub>3</sub>)<sub>6</sub>]Cl<sub>2</sub>Br is – (2) Zero (3) + 3(1) + 6(5) + 2Solution Let the oxidation number of Co be x Oxidation number of NH<sub>3</sub> is zero Oxidation number of Cl is -1 Oxidation number of Br is -1 Hence,  $x + 6(0) - (1 \times 2) - 1 = 0$ 

So, the oxidation number of cobalt in the given complex compound is +3.

**Illustration 2.** The order of increasing oxidation state of S in S<sub>8</sub>,  $S_2O_8^{-2}$ ,  $S_2O_3^{-2}$ ,  $S_4O_6^{-2}$  is given below-

(1) S <sub>8</sub>	$< S_2O$	$S_8^{-2} < S_2 O_3^{-2} <$	$S_4O_6^{-2}$	(2) $S_2O_8^{-2} <$	$S_2O_3^{-2} < S_4O_6^{-2} < S_8$
(3) $S_2$	$O_8^{-2} < C_8^{-2}$	$S_8 < S_4 O_6^{-2} <$	$S_2O_3^{-2}$	(4) $S_8 < S_2O$	$S_3^{-2} < S_4 O_6^{-2} < S_2 O_8^{-2}$
Solution	The c	oxidation num	ber of S are s	hown below along	g with the compounds
	$S_8$	$S_2O_8^{-2}$	$S_2O_3^{-2}$	$\mathbf{S}_4\mathbf{O}_6^{-2}$	
	0	+6	+2	+2.5	
	Henc	e the order of	increasing ox	didation state of S	is –
	$S_8 < 1$	$S_2O_3^{-2} < S_4O_6^{-2}$	$^{2} < S_{2}O_{8}^{-2}$		
Illustration 3	<b>3.</b> The c	oxidation num	ber of Cl in N	VOClO <sub>4</sub> is -	
(1) + 1	1	(2)	+9	(3) +7	(4) +5

x = +3

*.*..

Down by VISIONat Info Solution But I td	
rower by: visionet into Solution rvt. Liu	
Website : www.edubull.com	Mob no. : +91-9350679141

# Edubull

Solution	The compound	may be written as N	$O^+ClO_4^-$ .	
	For $ClO_4^-$ , Let	oxidation number of	Cl = a	
	;	$a + 4 \times (-2) = -1$		
		a = +7		
	Hence, the oxic	lation number of CI i	In NOCIO <sub>4</sub> 18 $+$ /	
<b>Illustrat</b> i	<b>ion 4.</b> The two possib	le oxidation number $(2) + 3 - 5$	of N atoms in $NH_4NC$ (3) -3. +5	$P_3$ are respectively- (4) $-3$ , $-5$
Solution	There are two	N atoms in $NH_4NC$	$D_3$ , but one N atom ha	as negative oxidation number
	(attached to H)	and the other has po	sitive oxidation numbe	er of (attached to O). Therefore
	evaluation shou	Ild be made separates $1 + 1 + 1 = 1$	ly as –	
	Oxidation num a + 4x(+1) = +1	ber of IN 18 IN II <sub>4</sub>	UXIdation initial $2(-2) = -2$	mber of N in $NO_3$
	$a + 4 \times (+1) - +$ a = -3	l	a + 3(-2) : $a = +5$	
	Here the two or $H_{\rm Here}$	xidation number are -	-3 and $+5$ respectively.	
,				
Illustrat	ion 5. The oxidation r	$\begin{array}{c} \text{umber of S in } H_2S_2U \\ (2) \\ \end{array}$	$D_8$ is -	$(A) \rightarrow A$
Solution	I) $+\delta$ In H <sub>2</sub> S <sub>2</sub> O <sub>8</sub> two	(2) - 0 O atoms form peroxi	(3) +0 ide linkage i.e.	(4) + 4
Dollar	Q (	Q		
	H-U-S-U-U-,	S-O-н ↓		
	Ŏ	Ŏ		
	$2 \times 1 + 2a + 6(-$	-2) + 2(-1) = 0		
	$\therefore$ $a = +o$ Thus the oxidat	tion number of S in F	$J_2S_2O_2$ is $\pm 6$	
	Thus the origin		120208 13 10	
Illustrati	ion 6. The oxidation r	number of S in (CH <sub>3</sub> )	$O_2$ SO is -	
()	1) 1	(2) 2	(3) 0	(4) 3
Solution	Dyidation num	on number of S is a ber of $CH_2 = +1$		
	Oxidation num	ber of $O = -2$		
	2(+1) + a + (-2)	()=0		
	a = 0			
	Hence the oxid	ation number of S in	dimethyl sulphoixae is	s zero.
		BEGINNE	ER'S BOX-1	
<b>1.</b> Ir	which of the followi	ng compounds, the c	oxidation state of I atom	n is highest ?
(1	$1) \mathrm{KI}_3$	(2) $KIO_4$	(3) $KIO_3$	$(4) \text{ IF}_5$
<b>2.</b> T	he oxidation number	of phosphorus in Ba	$(H_2PO_2)_2$ is -	
(1	1) +3	(2) + 2	(3) + 1	(4) –1
•	i i di seconda			
<b>3.</b> U	$\mathbf{V}$ idation number of $\mathbf{N}$	$\sqrt{11}$ in N1(CO <sub>4</sub> ) 1s -	( <b>3</b> ) <b>8</b>	(A) <b>7</b>
(.		(2) न	(5) 0	(4) 2
Power by: V	ISIONet Info Solution Pvt. Lt	d		
Website : wv	vw.edubull.com	Mob no. : +91-9	350679141	

4.	Positive oxidation state of an element indicates that it is -					
	(1) Elementry form	(2) Oxidised	(3) Reduced	(4) Only reductant		
5.	Predict the highest an (1) a[0, +3] b[0, +2] (3) a[+4, 0] b[+4, 0]	d lowest oxidation stat	e of (a) Ti and (b) Tl i (2) a[+3, 0] b[+4, 0] (4) a[+4, +2] b[+3, +	n combined state.		
6.	The oxidation state of (1) Zero	f oxygen atom in potas $(2) - 1/2$	sium superoxide is :- (3) -1	(4) -2		
6.4	APPLICATIONS O (A) To compare to Strength of ac Strength of ba	<b>F OXIDATION NUM</b> the strength of acid and id $\infty$ Oxidation Numbers se $\infty \frac{1}{Oxidation number}$	<b>IBER :</b> nd base :- er er			
	Example : or	der of acidic strength in	n HClO, HClO <sub>2</sub> , HClO	93, HClO4 will be.		
	Solution :		Oxidation nur	nber of chlorine		
		HClO (Hypo chlorous	s a <mark>cid) +1</mark>			
		HClO <sub>2</sub> (Chlorous acid	d) +3			
		HClO <sub>3</sub> (Chloric acid)	+5			
		$HCIO_4$ (Perchloric ac	id) +7			
	Θ Streng	th of acid $\propto$ Oxidation	Number			
	So the order w	vill be :-				
	HClO	$H > HClO_3 > HClO_2 > HClO_2$	ICIO			

## (B) To determine the oxidising and reducing nature of the substances :-

Oxidising agents are the substances which accept electrons in a chemical reaction i.e., electron acceptors are oxidizing agent.

Reducing agents are the substances which donate electrons in a chemical reaction i.e., electron donors are reducing agent.

Highest O.S.	+4	+5	+5	+6	+7	+6	+7	+8	+8	+2	+1
Elements	С	Ν	Р	S	Cl	Cr	Mn	Os	Ru	0	Η
Lowest O.S.	-4	-3	-3	-2	-1	0	0	0	0	-2	-1

(a) If effective element in a compound is present in maximum oxidation state then the compound acts as oxidizing agent.

Ex.	KMnO <sub>4</sub>	$K_2Cr_2O_7$	$H_2SO_4$	$SO_3$	$H_3PO_4$	HNO <sub>3</sub>	$HClO_4$
	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$
	+7	+6	+6	+6	+5	+5	+7

(b) If effective element in a compound is present in minimum oxidation state then the compound acts as reducing agent.

Ex.	$PH_3$	$NH_3$	$CH_4$
	$\downarrow$	$\downarrow$	$\downarrow$
	-3	-3	_4

(c) If effective element in a compound is present in intermediate oxidation state then the compound can act as well as reducing agent.

Ex.	$HNO_2$	$H_3PO_3$	$SO_2$	$H_2O_2$
	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$
	+3	+3	+4	-1

#### **(C)** To calculate the equivalent weight of compounds :

The equivalent weight of an oxidising agent or reducing agent is that weight which accepts one mole electrons in a chemical reaction or looses one mole electron in chemical reaction.

(a) Equivalent weight of oxidant =  $\frac{\text{Molecular weight}}{\text{No.of electrons gained by one mole}}$ 

## **Example :** In acidic medium

 $6e^{-} + Cr_2O_7^{2-} + 14H^+ \longrightarrow 2Cr^{3+} + 7H_2O$ 

Here atoms which undergoes reduction is Cr. Its O.S. is decreasing from +6 to +3

Equivalent weight of 
$$K_2Cr_2O_7 = \frac{\text{Molecular weight of } K_2Cr_2O_7}{3 \times 2} = \frac{M}{6}$$

Note :- [6 in denominator indicates that 6 electron were gained by  $Cr_2O_7^{2-}$  as it is clear from the given balanced equation]

(b) Equivalent weight of a reducant =  $\frac{\text{Molecular weight}}{\text{No.of electrons lost by one mole}}$ 

 $C_2O_4^{2-} \longrightarrow 2CO_2 + 2e^-$ In acidic medium,

Here, Total electrons lost = 2 So, equivalent weight =  $\frac{M}{2}$ 

(c) In different conditions a compound may have different equivalent wts. Because, it depends upon the number of electron gained or lost by that compound in that reaction. **Example :** 

(i)  $MnO_4^- \longrightarrow Mn^{+2}$  (acidic medium)

Here 5 electrons are taken so equivalent weight =  $\frac{M}{5} = \frac{158}{5} = 31.6$ 

(ii)  $MnO_4^- \longrightarrow MnO_2$  (neutral medium) or (Weak alkaline medium) (+7) (+4)

Here, only 3 electrons are gained, so eq. wt. =  $\frac{M}{3} = \frac{158}{3} = 52.7$ 

Note : When only alkaline medium is given consider it as weak alkaline medium.

(iii)  $MnO_4^- \longrightarrow MnO_4^{-2}$  (strong alkaline medium)

(+7) (+6)

Here, only one electron is gained, so eq. wt. =  $\frac{M}{1} = 158$ 

Note : KMnO<sub>4</sub> acts as an oxidant in every medium although with different strength which follows the order –

Acidic medium > neutral medium > alkaline medium

Power by: VISIONet Info Solution Pvt. Ltd
Website : www.edubull.com

While, K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> acts as an oxidant only in acidic medium as follows

$$Cr_2O_7^{2-} \longrightarrow 2Cr^{3+}$$

$$(2\times 6) \longrightarrow (2\times 3)$$

Here, 6 electrons are gained so eq. wt. =  $\frac{M}{6} = \frac{294}{6} = 49$ 

(D) To determine the possible molecular formula of compound : Since the sum of oxidation number of all the atoms present in a compound is zero, so the validity of the formula can be confirmed.

### **GOLDEN KEY POINTS**

## SOME OXIDIZING AGENTS / REDUCING AGENTS WITH EQUIVALENT WEIGHT

Species	Changed to	Reaction	Electrons exchanged or change in O.N.	Eq. wt.
$MnO_4^-(O.A.)$	Mn <sup>+2</sup> in acidic medium	$\frac{MnO_4^- + 8H^+ + 5e^- \rightarrow Mn^{2+} + }{4H_2O}$	5	$E = \frac{M}{5}$
$MnO_{4}^{-}(O.A.)$	$MnO_2$ in neutral medium	$MnO_4^- + 3e^- + 2H_2O \rightarrow MnO_2 + 4OH^-$	3	$E = \frac{M}{3}$
$MnO_4^-(O.A.)$	${\rm MnO_4^{2-}}$ in basic medium	$MnO_4^- + e^- \rightarrow MnO_4^{2-}$	1	$E = \frac{M}{1}$
$Cr_2O_7^{2-}(O.A.)$	Cr <sup>3+</sup> in acidic medium	$CrO_7^{2-} + 14H^+ + 2e^- \rightarrow 2Cr^{3+} + 7H_2O$	6	$E = \frac{M}{6}$
MnO <sub>2</sub> (O.A.)	Mn <sup>2+</sup> in acidic medium	$\frac{\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^- \rightarrow \text{Mn}^{2+} + 2\text{H}_2\text{O}}{2\text{H}_2\text{O}}$	2	$E = \frac{M}{2}$
Cl <sub>2</sub> (O.A.) in bleaching powder	Cl⁻	$Cl_2 + 2e^- \rightarrow 2Cl^-$	2	$E = \frac{M}{2}$
CuSO <sub>4</sub> (O.A.) in iodometric titration	Cu <sup>+</sup>	$Cu^{2+} + e^- \rightarrow Cu^+$	1	$E = \frac{M}{1}$
$S_2O_3^{2-}$ (R.A.)	$S_4O_6^{2-}$	$2S_2O_3^{2-} \rightarrow S_4O_6^{2-} + 2e^-$	2 (for two molecules)	$E = \frac{2M}{2} = M$
H <sub>2</sub> O <sub>2</sub> (O.A.)	H <sub>2</sub> O	$\mathrm{H}_{2}\mathrm{O}_{2} + 2\mathrm{H}^{+} + 2\mathrm{e}^{-} \rightarrow 2\mathrm{H}_{2}\mathrm{O}$	2	$E = \frac{M}{2}$
H <sub>2</sub> O <sub>2</sub> (R.A.)	O <sub>2</sub>	$H_2O_2 \rightarrow O_2 + 2H^+ + 2e^-$ (O.N. of oxygen in $H_2O_2$ is -1 per atom)	2	$E = \frac{M}{2}$
$Fe^{2+}$ (R.A.)	Fe <sup>3+</sup>	$Fe^{2+} \rightarrow Fe^{3+} + e^{-}$	1	$E = \frac{M}{1}$

# Illustrations

**Illustration 7.** To find the n-factor in the following chemical changes.

(i) KMnO<sub>4</sub>  $\xrightarrow{H^+}$  Mn<sup>2+</sup>

(ii) KMnO<sub>4</sub>  $\xrightarrow{H_2O}$  Mn<sup>4+</sup>

Power by: VISIONet Info Solution Pvt. Ltd Website : www.edubull.com

### Edubull

(iii) KMnO<sub>4</sub>  $\xrightarrow{OH (concentrated basic medium)} Mn^{6+}$  (iv) K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>  $\xrightarrow{H^+} Cr^{3+}$ (v)  $C_2 O_4^{2-} \longrightarrow CO_2$ (vi)  $FeSO_4 \longrightarrow Fe_2O_3$ (vii)  $Fe_2O_3 \longrightarrow FeSO_4$ (i) In this reaction, KMnO<sub>4</sub> which is an oxidizing agent, itself gets reduced to  $Mn^{2+}$ Solution under acidic conditions.  $n = |1 \times (+7) - 1 \times (+2)| = 5$ (ii) In this reaction,  $KMnO_4$  gets reduced to  $Mn^{4+}$  under neutral or slightly (weakly) basic conditions.  $n = |1 \times (+7) - 1 \times (+4)| = 3$ (iii) In this reaction,  $KMnO_4$  gets reduced to  $Mn^{6+}$  under basic conditions.  $n = |1 \times (+7) - 1 \times (+6)| = 1$ (iv) In this reaction,  $K_2Cr_2O_7$  which acts as an oxidizing agent reduced to  $Cr^{3+}$  under acidic conditions. (It does not react under basic conditions.)  $n = |2 \times (+6) - 2 \times (+3)| = 6$ (v) In this reaction  $C_2 O_4^{2-}$  (oxalate ion) gets oxidized to  $CO_2$  when it is reacted with an oxidizing agent.  $n = |2 \times (+3) - 2 \times (+4)| = 2$ (vi) In this reaction, ferrous ions get oxidized to ferric ions.  $n = |2 \times (+2) - 1 \times (+3)| = 1$ (vii) In this reaction, ferric ions are getting reduced to ferrous ions.  $n = |2 \times (+3) - 2 \times (+2)| = 2$ Illustration 8. Suppose that there are three atoms A, B, C and their oxidation number are 6, -1, -2,

respectively. Then the molecular formula of compound will be.

Solution Since, the charge on a free compound is zero. So

> $+6 = (-1 \times 4) + (-2)$ +6 = -6Or  $+6 = (-1 \times 2) + (-2 \times 2)$ = -2 + (-4) = -6

So molecular formula,  $AB_4C$  or  $AB_2C_2$ 

## **BEGINNER'S BOX-2**

Molecular weight of KMnO<sub>4</sub> in acidic medium and neutral medium will be respectively -1.

(1) 7  $\times$  equivalent weight and 2  $\times$  equivalent weight

(2) 5  $\times$  equivalent weight and 3  $\times$  equivalent weight

(3) 4  $\times$  equivalent weight and 5  $\times$  equivalent weight

(4)  $2 \times$  equivalent weight and  $4 \times$  equivalent weight

2. In acidic medium, equivalent weight of  $K_2Cr_2O_7$  (Molecular weight = M) is -(3) M/6 (1) M/3(2) M/4(4) M/2

#### **OXIDATION AND REDUCTION** 6.5

There are two concepts of oxidation and reduction.

## (A) Classical / old concept :-

	OXIDATION	REDUCTION
(1)	Addition of O <sub>2</sub>	Addition of H <sub>2</sub>

Power by: VISIONet Info Solution Pvt. Ltd	
Website : www.edubull.com	Mob no. : +91-9350679141

	$2Mg + O_2 \rightarrow 2MgO$	$N_2 + Cl_2 \rightarrow 2NH_3$
	$C + O_2 \rightarrow CO_2$	$H_2 + Cl_2 \rightarrow HCl$
(2)	Removal of H <sub>2</sub>	Removal of O <sub>2</sub>
	$H_2S + Cl_2 \rightarrow 2HCl + S$ (oxidation of $H_2S$ )	$CuO + C \rightarrow Cu + CO$ (reduction of CuO)
	$4HI + O_2 \rightarrow 2I_2 + 2H_2O$ (oxidation of HI)	$H_2O + C \rightarrow CO + H_2$ (reduction of $H_2O$ )
(3)	Addition of electronegative element	Addition of electropositive element
	$Fe + S \rightarrow FeS$ (oxidation of Fe)	$CuCl_2 + Cu \rightarrow Cu_2Cl_2$ (reduction of $CuCl_2$ )
	$SnCl_2 + Cl_2 \rightarrow SnCl_4$ (oxidation of $SnCl_2$ )	$HgCl_2 + Hg \rightarrow Hg_2Cl_2$ (reduction of $HgCl_2$ )
(4)	Removal of electropositive element	Removal of electronegative element
	$2NaI + H_2O_2 \rightarrow 2NaOH + I_2$ (oxidation of NaI)	$2\text{FeCl}_3 + \text{H}_2 \rightarrow 2\text{FeCl}_2 + 2\text{HCl}$ (reduction of
		FeCl <sub>3</sub> )

## (B) Electronic / Modern Concept

(1) De-electronation	Electroation
(2) Oxidation process are those process in	Reduction process are those process in which
which one or more e <sup>-</sup> s are lost by an atom, ion	one or more e <sup>-</sup> s are gained by an atom, ion or
or molecule.	molecule.
(3) Example :-	
(a) $Zn \rightarrow Zn^{+2} + 2e^{-1}$	(a) $Cu^{+2} 2e^{-} \rightarrow Cu$
$M \rightarrow M^{n+} + ne^-$	$M^{n+} + ne^- \rightarrow M$
(b) $\text{Sn}^{+2} \rightarrow \text{Sn}^{+4} + (4-2)e^{-1}$	(b) $Fe^{+3} + (3-2)e^{-} \rightarrow Fe^{+2}$
$M^{+n_1} \rightarrow M^{+n_2} (n_2 - n_1)e^-$	$M^{+x_1} (x_1 - x_2)e^- \rightarrow M^{+x_2}$
(c) $Cl^- \rightarrow Cl + e^-$	(c) $O + 2e^- \rightarrow O^{2-}$
$A^{-n} \rightarrow A + ne^{-}$	$A + xe \rightarrow A^{-x}$
(d) $MnO_4^{-2} \rightarrow MnO_4^{-} + (2-1)e^{-1}$	(d) $[Fe(CN)_4]^{3-} (4-3)e^- \rightarrow [Fe(CN)_4]^{-4}$
$A^{-n_1} \rightarrow A^{-n_2} (n_1 - n_2)e^-$	$\mathbf{A}^{-\mathbf{n}_1} (\mathbf{n}_1 - \mathbf{n}_2) \mathbf{e}^- \rightarrow - \mathbf{A}^{-\mathbf{n}_2}$

# 6.6 TYPES OF REDOX REACTIONS

(A) Intermolecular redox reaction :- When oxidation and reduction takes place separately in different compounds, then the reaction is called intermolecular redox reaction.

 $SnCl_{2} + 2FeCl_{3} \longrightarrow SnCl_{4} 2FeCl_{2}$   $Sn^{+2} \longrightarrow Sn^{+4} (Oxidation)$  $Fe^{+3} \longrightarrow Fe^{+2} (Reduction)$ 

(B) Inteamolecular redox reaction :- During the chemical reactions place in single compound then there action is called intermolecular redox reaction.



(C) **Disproportion reaction :-** when reduction an oxidation take place in the same element of the same compound then the reaction is called disproportionation reaction.

 $H_2O_2 \longrightarrow H_2O + 1/2O_2$ 

Power by: VISIONet Info Solution Pvt. Ltd	
Website : www.edubull.com	Mob no. : +91-9350679141



(D) Comproportionation reaction : Reverse of disproportionation reaction known as comproportionation reaction Ex.  $ClO^- + Cl^- \rightarrow Cl_2 + OH^-$ 

### **BEGINNER'S BOX-3**

 Oxidation is defined as-(1) Gain of electrons
 (3) Loss of electron

- (2) Decreases in positive valency
- (4) Addition of electropositive element
- Reduction is defined as 
   (1) Increases in positive valency
   (3) Loss of protons
- (2) Gain of electrons
- (4) Decreases in negative valency

3. In the reaction  $MnO_4^- + SO_3^{-2} + H^+ \longrightarrow SO_4^{-2} + Mn^{2+} + H_2O$ 

- (1)  $MnO_4^-$  and  $H^+$  both are reduced
- (2)  $MnO_4^-$  is reduced and  $H^+$  is oxidised
- (3)  $MnO_4^-$  is reduced and  $SO_3^{-2}$  is oxidised
- (4)  $MnO_4^-$  is oxidised and  $SO_3^{-2}$  is reduced
- 4. The charge on cobalt in  $[Co(CN)_6]^{-3}$  is (1) -6 (2) -3 (3) +3 (4) +6
- 5. Which of the following halogen always show only one oxidating state its compounds? (1) Cl (2) F (3) Br (4) I
- 6. Which of the following reaction do not involve oxidation-reduction ?
  (1) 2Rb +2H<sub>2</sub>O → 2RbOH + H<sub>2</sub>
  (2) 2CuI<sub>2</sub> → 2CuI + I<sub>2</sub>
  (3) NH<sub>4</sub>Cl + NaOH → NaCl + NH<sub>3</sub> + H<sub>2</sub>O
  - (4)  $3Mg + N_2 \longrightarrow Mg_3N_2$
- 7. The fast reaction between water and sodium is the example of (1) Oxidation
  (2) Reduction
  (3) Intermolecular redox
  (4) Intramolecular redox
- 8. Choose the redox reaction from the following :-
  - $(1) \operatorname{Cu} + 2\operatorname{H}_2\operatorname{SO}_4 \longrightarrow \operatorname{CuSO}_4 + \operatorname{SO}_2 + 2\operatorname{H}_2\operatorname{O}$

(2)  $BaCl_2 + H_2SO_4 \longrightarrow BaSO_4 + 2HCl$ 

- $(3) 2NaOH + H_2SO_4 \longrightarrow Na_2SO_4 + 2H_2O$
- (4)  $KNO_3 + H_2SO_4 \longrightarrow 2HNO_3 + K_2SO_4$
- 9. Which of the following not a redox reaction ?

Power by: VISIONet Info Solution Pvt. Ltd	
Website : www.edubull.com	Mob no. : +91-9350679141

- (1)  $MnO_4^- \longrightarrow MnO_2 + O_2$
- (2)  $Cl_2 + H_2O \longrightarrow HCl + HClO$
- (3)  $2CrO_4^{-2} + 2H^+ \longrightarrow 2CrO_7^{2-} + H_2O$
- (4)  $MnO_4^- + 8H^+ + 5Ag \longrightarrow Mn^{+2} + 4H_2O + 5Ag^+$
- 10. In the reaction  $6Li + N_2 \longrightarrow 2LI_3N$ (1) Li undergoes reduction (3) N undergoes oxidation
- (2) Li undergoes oxidation
- (4) Li is oxidation
- 11.  $H_2O_2 + H_2O_2 \longrightarrow 2H_2O + O_2$  is an example of dispropotionation because :-
  - (1) Oxidation number of oxygen only decreases
  - (2) Oxidation number of oxygen only increases
  - (3) Oxidation number of oxygen decreases as well as increase
  - (4) Oxidation number of oxygen neither decreases nor increases

## 6.7 BALANCING OF REDOX REACTION :-

- (A) Oxidation number change method
- (B) Ion electron method.

## (A) Oxidation number change method :-

This method was given by Jonson. In a balanced redox reaction, total increase in oxidation number must be equal to total decreases in oxidation number. This equivalence provides the basis for balancing redox reactions.

The general procedure involves the following steps :-

(i) Select the atom in oxidising agent whose oxidation number decreases and indicate the gain of electrons.

(ii) Select the atom in reducing agent whose oxidation number increases and indicate the loss of electrons.

(iii) Now cross multiply i.e. multiply oxidizing agent by the number of loss of electrons and reducing agent by number of gain of electrons.

(iv) Balance the number of atoms on both sides whose oxidation numbers change in the reaction.

(v) In order to balance oxygen atom, add  $H_2O$  molecules to the side deficient in oxygen.

(vi) Then balance the number of H atoms by adding  $H^+$  ions to the side deficient in hydrogen.

	Illustrations
Illustration	9. Balance the following reaction by the oxidation number method :-
	$Cu + HNO_3 \longrightarrow Cu(NO_3)_2 + NO_2 + H_2O$
Solution	Write the oxidation number of all the atoms.
	0 +1+5-2 +2+5-2 +4-2 +1-2
	$Cu + HNO_3 \longrightarrow Cu(NO_3)_2 + NO_2 + H_2O$
	There is change in oxidation number of Cu and N.
	0 +2+5-2
	Cu $\longrightarrow$ Cu(NO <sub>3</sub> ) <sub>2</sub> (1) (Oxidation no. is increased by 2)
	+5 +4
	$HNO_3 \longrightarrow NO_2$ (2) (Oxidation no. is decreased by 1)
Power by: VISIO	Net Info Solution Pvt. Ltd
Website : www.ee	Mob no. : +91-9350679141

To make increase and decrease equal, eq. (2) is multiplied by 2.  $Cu + 2HNO_3 \longrightarrow Cu(NO_3)_2 + 2NO_2 + H_2O$ Balancing nitrates ions, hydrogen and oxygen, the following equation is obtained.  $Cu + 4HNO_3 \longrightarrow Cu(NO_3)_2 + 2NO_2 + 2H_2O$ This is the balance dequation. Illustration 10. Balance the following reaction by the oxidation number method :-  $MnO_4^- + Fe^{+2} \longrightarrow Mn^{+2} + Fe^{+3}$ Solution Write the oxidation number of all the atoms. +7 - 2  $MnO_4^- + Fe^{+2} \longrightarrow Mn^{+2} + Fe^{+3}$ Change in oxidation number has occurred in Mn and Fe. +7

 $\begin{array}{c} MnO_{4}^{-} \longrightarrow Mn^{+2} \qquad \dots \dots \dots (1) \text{ (Decrement in oxidation no. by 5)} \\ Fe^{+2} \longrightarrow Fe^{+3} \qquad \dots \dots \dots (2) \text{ (Increment in oxidation no. by 1)} \\ To make increase and decrease equal, eq. (2) is multiplied by 5. \end{array}$ 

 $MnO_4^- + 5Fe^{+2} \longrightarrow Mn^{+2} + 5Fe^{+3}$ 

To balance oxygen,  $4H_2O$  are added to R.H.S. and to balance hydrogen,  $8H^+$  are added to L.H.S  $MnO_4^- + 5Fe^{+2} + 8H^+ \longrightarrow Mn^{+2} + 5Fe^{+3} + 4H_2O$ 

This is the balanced equation.

## (B) Ion-Electron method :-

This method was given by Jette and La Mev in 1972.

The following steps are followed while balancing redox reaction (equations) by this method.

(i) Write the equation in ionic from.

(ii) Split the redox equation into two half reactions, one representing oxidation and the other representing reduction.

(iii) Balance these half reactions separately and then add by multiplying with suitable coefficients so that the electrons are cancelled. Balancing is done using following substeps.

(a) Balance all other atoms except H and O.

(b) Then balance oxygen atoms by adding  $H_2O$  molecules to the side deficient in oxygen. The number of  $H_2O$  molecules added is equal to the deficiency of oxygen atoms.

(c) Balance hydrogen atoms by adding  $H^+$  ions equal to the deficiency in the side which is deficient in hydrogen atoms.

(d) Balance the charge by adding electrons tot the die which is rich in +ve charge. i.e. deficient in electrons. Number of electrons added is equal to the deficiency.

(e) Multiply the half equations with suitable coefficients to equalize the number of electrons.

(iv) Add these half equations to get an equation which is balanced with respect to charge and atoms.

(v) If the medium of reaction is basic,  $OH^-$  ions are added to both sides of balanced equation , which is equal to number of  $H^+$  ions in Balanced Equation.

# Illustrations

Illustration 11. Balance the following reaction by ion-electron method in acidic medium :-

$$\operatorname{Cr}_2\operatorname{O}_7^{2-} + \operatorname{C}_2\operatorname{O}_4^{2-} \longrightarrow \operatorname{Cr}^{3+} + \operatorname{CO}_2$$

Solution

 $Cr_2O_7^{2-} + C_2O_4^{2-} \longrightarrow Cr^{3+} + CO_2$ Write both the half reaction. (a)  $Cr_2O_7^{2-} \longrightarrow Cr^{3+}$  (Reduction half reaction)  $C_2O_4^{2-} \longrightarrow CO_2$  (Oxidation half reaction) Atoms other than H and O are balanced (b)  $Cr_2O_7^{2-} \longrightarrow 2Cr^{3+}$  $C_2O_4^{2-} \longrightarrow 2CO_2$ Balance O-atoms by the addition of  $H_2O$  to another side (c)  $Cr_2O_7^{2-} \longrightarrow 2Cr^{3+} + 7H_2O$  $C_2O_4^{2-} \longrightarrow 2CO_2$ Balance H-atoms by the addition of H<sup>+</sup> to another side (d)  $Cr_2O_7^{2-}$  14 H<sup>+</sup>  $\longrightarrow$  2Cr<sup>3+</sup> + 7H<sub>2</sub>O  $C_2 O_4^{2-} \longrightarrow 2CO_2$ Now, balance the charge by the addition of electron (e<sup>-</sup>) (e)  $Cr_2O_7^{2-}$  14H<sup>+</sup> + 6e<sup>-</sup>  $\longrightarrow$  2Cr<sup>3+</sup> + 7H<sub>2</sub>O  $C_2O_4^{2-} \longrightarrow 2CO_2 + 2e^-$ .....(2) Multiply equations by a constant to get the same number of electrons on both (f) side. In the above case second equation is multiplied by 3 and then added to first equation  $Cr_{*}O_{*}^{2-} + 14H^{+} + 6e^{-} \longrightarrow 2Cr^{3+} + 7H_{2}O_{*}$ 

$$3C_2O_4^{2-} \longrightarrow 6CO_2 + 6e^-$$

$$Cr_2O_7^{2-} + 3C_2O_4^{2-} + \longrightarrow 2Cr^{3+} + 6CO_2 + 7H_2O$$

Illustration 12. Balance the following reaction by ion-electron method :-

$$Cr(OH)_3 + IO_3^- \xrightarrow{OH^-} \Gamma + CrO_4^{2-}$$

Sol

Website : www.edubull.com

Solution 
$$\operatorname{Cr}(\operatorname{OH})_3 + \operatorname{IO}_3^- \xrightarrow{\operatorname{OH}^-} \Gamma + \operatorname{CrO}_4^{2^-}$$
  
(a) Separate the two half reactions.  
 $\operatorname{Cr}(\operatorname{OH})_3 \longrightarrow \operatorname{CrO}_4^{2^-}$  (Oxidation half reaction)  
 $\operatorname{IO}_3^- \longrightarrow \Gamma$  (Reduction of half reaction)  
(b) Balance O-atoms by adding H<sub>2</sub>O.  
 $\operatorname{H}_2\operatorname{O} + \operatorname{Cr}(\operatorname{OH})_3 \longrightarrow \operatorname{CrO}_4^{2^-}$   
 $\operatorname{IO}_3^- \longrightarrow \Gamma + 3\operatorname{H}_2\operatorname{O}$   
(c) Balance H-atoms by adding H<sup>+</sup> to side having deficiency and add equal no of  
 $\operatorname{OH}^-$  ions to the side ( $\Theta$  medium is known)  
 $\operatorname{H}_2\operatorname{O} + \operatorname{Cr}(\operatorname{OH})_3 \longrightarrow \operatorname{CrO}_4^{2^-} + 5\operatorname{H}^+$   
 $5\operatorname{OH}^- + \operatorname{H}_2\operatorname{O} + \operatorname{Cr}(\operatorname{OH})_3 \longrightarrow \operatorname{CrO}_4^{2^-} + 5\operatorname{H}^+ + 5\operatorname{OH}^-$ 

Mob no. : +91-9350679141

or 
$$5OH^- + Cr(OH)_3 \longrightarrow CrO_4^{2-} + 4H_2O$$
  
 $IO_3^- + 6H^+ \longrightarrow \Gamma + 3H_2O$   
 $IO_3^- + 6H^+ + 6OH^- \longrightarrow \Gamma + 3H_2O + 6OH^-$   
or  $IO_3^- + 3H_2O \longrightarrow \Gamma + 6OH^-$   
(d) Balance the charges by adding electrons  
 $5OH^- + Cr(OH)_3 \longrightarrow CrO_4^{2-} + 4H_2O + 3e^-$   
 $IO_3^- + 3H_2O + 6e^- \longrightarrow \Gamma + 6OH^-$   
(e) Multiply first equation by 2 and add to second to give  
 $100 H^- + 2Cr(OH)_3 \longrightarrow 2CrO_4^{2-} + 8H_2O + 6e^-$   
 $IO_3^- + 3H_2O + 6e^- \longrightarrow \Gamma + 6OH^-$   
 $4OH^- + 2Cr(OH)_3 + IO_3^- \longrightarrow 5H_2O + 2CrO_4^{2-} + \Gamma -$ 

## 6.8 LAW OF EQUIVALENCE

The law states that one equivalent of an element combine with one equivalent of the other, and in a chemical reaction equivalent and mill equivalent of reactants react in equivalent to give same no. of equivalent or milli equivalents of products separately.

According :

(i)  $aA + bB \rightarrow mM + nN$ m. eq of A = m. eq of B = m. eq of M = m. eq of N(ii) In a compound  $M_xN_y$ m. eq of  $M_xN_y = m$ . eq of M = m. eq of N

## **GOLDEN KEY POINTS**

• FOR ACID-BASE (NEUTRALIZATION REACTION) OR REDOX REACTION :  $N_1V_1 = N_2V_2$  is always true. But  $M_1V_1 = M_2V_2$  (may or may not be true) But  $M_1 \times n_1 \times V_1 = M_2 \times n_2 \times V_2$  (always true when n terms represent n-factor)

## Illustrations

- **Illustration 13.** Calculate the normality of a solution containing 15.8 g of KMnO<sub>4</sub> in 50 mL acidic solution .
- Solution

Normality (N) = 
$$\frac{W \times 1000}{E \times VmL}$$

Where 
$$W = 15.8$$
 g,  $V = 50$  mL  $E = \frac{\text{molar mass}}{100}$ 

$$\frac{\text{ar mass of KMnO}_4}{\text{Valence factor}} = \frac{158}{5} = 31.6$$

So, N = 10

- **Illustration 14.** Calculate the normality of a solution containing 50 mL of 5 M solution  $K_2Cr_2O_7$  in acidic medium.
- **Solution** Normality (N) = Molarity  $\times$  Valence factor = 5  $\times$  6 = 30 N

Power by: VISIONet Info Solution Pvt. Ltd
Website : www.edubull.com

**Illustration 15.** Find the number of moles of  $KMnO_4$  needed to oxidize one mole  $Cu_2S$  in acidic medium . The reaction is  $KMnO_4 + Cu_2S \longrightarrow Mn^{2+} + Cu^{2+} + SO_2$ 

Solution From law of equivalence

Equivalents of Cu<sub>2</sub>S = equivalents of KMnO<sub>4</sub> Moles of Cu<sub>2</sub>S × v.f. = moles of KMnO<sub>4</sub> × v.f.  $1\times 8 = n_2 \times 5$  $n_2 = \frac{8}{5} = 1.6$ 

**Illustration 16.** The number of moles of oxalate ions oxidized by one mole of  $MnO_4^-$  ion in acidic medium.

(1) 
$$\frac{5}{2}$$
 (2)  $\frac{2}{5}$  (3)  $\frac{3}{5}$  (4)  $\frac{5}{3}$   
Solution Equivalent of  $C_2O_4^{2-}$  = equivalents of  $MnO_4^{-}$   
 $x \text{ (mole)} \times 2 = 1 \times 5 \text{ ; } x = \frac{5}{2}$ 

**Illustration 17.** What volume of 6 M HCl and 2 M HCl should be mixed to get two litre of 3 M HCl? **Solution** Let, the volume of 6 M HCl required to obtain 2 L of 3M HCl = x L

> $\therefore$  Volume of 2 M HCl required = (2 - x)L $M_1V_1$ + $M_2V_2$  $M_3V_3$ 6M HCl 2M HCl 3M HCl  $6 \times (x) + 2 \times (2 - x) = 3 \times 2$ 6x + 4 - 6x = 6 $\Rightarrow 4x = 2$  $\Rightarrow$ .... x = 0.5 LHence, volume of 6 M HCl required = 0.5 LVolume of 2 M HCl required = 1.5 L

**Illustration 18.** In a reaction vessel, 1.184 g of NaOH is required to be added for completing the reaction. How many milli litre of 0.15 M NaOH should be added for this requirement ? **Solution** Amount of NaOH present in 1000 mL of 0.15 M NaOH =  $0.15 \times 40 = 6$  g

$$\therefore \quad 1 \text{ ml of this solution contain NaOH} = \frac{6}{1000} \times 10^{-3} \text{ g}$$
  
$$\therefore 1.184 \text{ g of NaOH will be present in} = \frac{1}{6 \times 10^{-3}} \times 1.184 = 197.33 \text{ mL}$$

**Illustration 19.** What weight of Na<sub>2</sub>CO<sub>3</sub> of 85% purity would be required to prepare 45.6 mL of 0.253 NH<sub>2</sub>SO<sub>4</sub>?

Solution

Meq. of 
$$Na_2CO_3 = Meq.$$
 of  $H_2SO_4 = 45.6 \times 0.235$ 

$$\frac{W_{Na_2CO_3}}{E_{Na_2CO_3}} \times 1000 = 45.6 \times 0.235 \qquad \Rightarrow \frac{W_{Na_2CO_3}}{106/2} \times 1000 = 45.6 \times 0.235$$

$$W_{Na_2CO_3} = 0.5679 \text{ g}$$

For 85 g of pure  $Na_2CO_3$ , weighed sample = 100 g

For 0.5679 g of pure Na<sub>2</sub>CO<sub>3</sub>, weighed sample =  $\frac{100}{85} \times 0.5679 = 0.6681$  g *.*.

**Illustration 20.** The number of moles of KMnO<sub>4</sub> that will be required to react with 2 mol of ferrous oxalate is

 $(1) \frac{6}{5}$  $(3) \frac{4}{5}$ (2)  $\frac{2}{5}$ (4) 1 $Mn^{7+} + 5e^- \rightarrow Mn^{2+} ] \times 3$ Solution  $\mathrm{Fe}^{2+} \rightarrow \mathrm{Fe}^{3+} + \mathrm{e}^{-}$  $C_2O_4^{2-} \rightarrow 2CO_2 + 2e^-$ 3 moles of  $KMnO_4 = 5$  moles of  $FeC_2O_4$  $\therefore$  2 mol of ferrous oxalate =  $\frac{6}{5}$  mole of KMnO<sub>4</sub> Hence, (A) is the correct answer.

**Illustration 21.** What volume of 6 M HNO<sub>3</sub> is needed to oxidize 8 g of  $Fe^{2+}$  to  $Fe^{3+}$ , HNO<sub>3</sub> gets converted to NO?

(1) 8 mL (2) 7.936 mL (3) 32 mL (4) 64 mL Meq. of HNO<sub>3</sub> = Meq. of  $Fe^{2+}$ Solution

or 
$$6 \times 3 \times V = \frac{8}{56} \times 1000$$
  
 $V = 7.936 \text{ mL}$   
 $n \text{-factor} = 3$   
 $+5$  2+  
 $(\text{NO}_3^- \rightarrow \text{NO})$ 

Hence, (B) is the correct answer.

**Illustration 22.** Which of the following is/are correct?

- (1) g mole wt. = mol. wt. in g = wt. of  $6.02 \times 10^{23}$  molecules (2) mole = N<sub>A</sub> molecule =  $6.02 \times 10^{23}$  molecules
- (3) mole = g molecules
- (4) None of the above

Solution **Ans.** (A), (B) and (C)

## **BEGINNER'S BOX-4**

1.	In the half reaction : $2ClO_3^- \longrightarrow Cl_2$	
	(1) 5 electrons are gained	(2) 5 electrons are liberated
	(3) 10 electrons are gained	(4) 10 electrons are liberated
2	The number of electrons required to h	palance the following equation:-

of electrons required to balance the following equation:  $NO_3^- + 4H^- + e^- \longrightarrow 2H_2O + NO$  are-(A) 5 (B) 4 (C) 3 (D) 2

3. Which of the following equation is balanced one-

Website : www.edubull.com Mob r	no. : +91-9350679141

(1) 
$$5BiO_{3}^{-} + 22H^{+} + Mn^{2+} \longrightarrow 5Bi^{3+} + 7H_{2}O + MnO_{4}^{-}$$
  
(2)  $5BiO_{3}^{-} + 14H^{+} + 2Mn^{2+} \longrightarrow 5Bi^{3+} + 7H_{2}O + 2MnO_{4}^{-}$   
(3)  $2BiO_{3}^{-} + 4H^{+} + Mn^{2+} \longrightarrow 2Bi^{3+} + 2H_{2}O + MnO_{4}^{-}$   
(4)  $6BiO_{3}^{-} + 12H^{+} + 3Mn^{2+} \longrightarrow 6Bi^{3+} + 6H_{2}O + 3MnO_{4}^{-}$ 

#### **ANSWER KEY BEGINNER'S BOX-1** (2) 2. (3) (2) 1. 3. 4. (2) 5. (4) 6. (1)**BEGINNER'S BOX-2** (3) 1. (2) 2. **BEGINNER'S BOX-3** 3. 7. (3) 2. (2) 4. 5. (2)6. (3) 1. (3) (3) (3) 8. 9. (1) (3) 10. (2) 11. (3) **BEGINNER'S BOX-4** 1. (3) 2. (3) 3. (2)