


Fundamentals of Mathematics-II

He is unworthy of the name of man who is ignorant of the fact that the diagonal of square is incommensurable with its sidePlato

Absolute value function / modulus function :

The symbol of modulus function is $f(x) = |x|$ and is defined as: $y = |x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$.



Properties of modulus : For any $a, b \in \mathbb{R}$

- (i) $|a| \geq 0$
- (ii) $|a| = |-a|$
- (iii) $|a| \geq a, |a| \geq -a$
- (iv) $|ab| = |a| |b|$
- (v) $\left| \frac{a}{b} \right| = \frac{|a|}{|b|}$
- (vi) $|a + b| \leq |a| + |b|$; Equality holds when $ab \geq 0$
- (vii) $|a - b| \geq ||a| - |b||$; Equality holds when $ab \geq 0$

Example # 1 : Solve the following linear equations

$$\begin{array}{ll} (\text{i}) \quad x|x| = 4 & (\text{ii}) \quad |x - 3| + 2|x + 1| = 4 \\ \text{Solution :} \quad (\text{i}) \quad x|x| = 4 & \end{array}$$

If $x > 0$

$$\therefore x^2 = 4 \Rightarrow x = \pm 2$$

$$\therefore x = 2 \quad (\because x \geq 0)$$

If $x < 0 \Rightarrow -x^2 = 4 \Rightarrow x^2 = -4$ which is not possible

$$(\text{ii}) \quad |x - 3| + 2|x + 1| = 4$$

case I : If $x \leq -1$

$$\therefore -(x - 3) - 2(x + 1) = 4$$

$$\Rightarrow -x + 3 - 2x - 2 = 4$$

$$\Rightarrow -3x = 3 \Rightarrow x = -1$$

case II : If $-1 < x \leq 3$

$$\therefore -(x - 3) + 2(x + 1) = 4$$

$$\Rightarrow -x + 3 + 2x + 2 = 4 \Rightarrow x = -1 \text{ which is not possible}$$

case III : If $x > 3$

$$x - 3 + 2(x + 1) = 4$$

$$3x - 1 = 4 \Rightarrow x = 5/3$$

which is not possible $\therefore x = -1 \text{ Ans.}$

Rational function :

A rational function is a function of the form, $y = f(x) = \frac{g(x)}{h(x)}$, where $g(x)$ & $h(x)$ are polynomial functions.

Irrational function :

An irrational function is a function $y = f(x)$ in which the operations of addition, subtraction, multiplication, division and raising to a fractional power are used.

For example $y = \frac{x^3 + x^{1/3}}{2x + \sqrt{x}}$ is an irrational function

(a) The equation $\sqrt{f(x)} = g(x)$, is equivalent to the following system

$$f(x) = g^2(x) \quad \& \quad g(x) \geq 0$$

(b) The inequation $\sqrt{f(x)} < g(x)$, is equivalent to the following system

- $f(x) < g^2(x)$ & $f(x) \geq 0$ & $g(x) \geq 0$
- (c) The inequation $\sqrt{f(x)} > g(x)$, is equivalent to the following system
 $g(x) \leq 0$ & $f(x) \geq 0$ or $g(x) \geq 0$ & $f(x) > g^2(x)$

Example # 2 : Solve : $x + 2 > 2 \sqrt{1-x^2}$

Solution : $4(1-x^2) < (x+2)^2$ and $x+2 \geq 0$ & $1-x^2 \geq 0$

$$x \in \left(-\infty, -\frac{4}{5}\right) \cup (0, \infty) \quad \dots(1)$$

$$x \in [-2, \infty) \quad \dots(2)$$

$$x \in [-1, 1] \quad \dots(3)$$

$$(1) \cap (2) \cap (3)$$

$$\left[-1, -\frac{4}{5}\right) \cup (0, 1]$$

Self Practice Problem :

$$(1) \sqrt{2x^2 + x - 6} < x$$

$$(2) \sqrt{5-x} > x + 1$$


$$(3) x + 3 + \sqrt{x^2 + 4x - 5} > 0$$

$$(4) \sqrt{x} - \sqrt{4-x} \geq 1$$

$$\text{Ans. } (1) \left[\frac{3}{2}, 2\right) \quad (2) (-\infty, 1) \quad (3) (-\infty, -1] \cup [5, \infty) \quad (4) \left[\frac{4+\sqrt{7}}{2}, 4\right]$$

Graphs Related to modulus :

If graph of $y = f(x)$ is



then draw graph of

$$(a) y = -f(x)$$

$$(b) y = f(-x)$$





$$(c) y = f(|x|)$$


$$(d) y = f(-|x|)$$

$$(e) y = |f(x)|$$

$$(f) |y| = f(x)$$


$$(g) |y| = -f(x)$$








Graphical Transformation :

If graph of $y = f(x)$ is




then graph of (a) $y - k = f(x - h)$ (b) $y = kf(x)$, ($k > 0$) (c) $y = f(kx)$, ($k > 0$)






Example # 3 : $y = |x^2 + 4x + 3|$

Solution :




Example # 4 : $|y-1| = \sin x$

Solution :




Example # 5 : $y = -x^2 + 4|x| - 3$

Solution :



Example # 6 : $y = ||x| - 3|$

Solution :



Example # 7 : $y = \sin\left(\frac{x}{3}\right)$


Solution : period is 6π

Example # 8 : $y = |\sin x - 3|$

Solution : Graphical Transformation

Example # 9 : $y = |-e^{\ln|x|}|$

Solution :



Example # 10 : $|y| = x^2 - 3x + 2$

Solution :

